

The Log-Behavior of $\sqrt[n]{p(n)}$ and $\sqrt[n]{p(n)/n}$

William Y.C. Chen¹ and Ken Y. Zheng²

¹Center for Applied Mathematics
Tianjin University
Tianjin 300072, P. R. China

²Center for Combinatorics, LPMC
Nankai University
Tianjin 300071, P. R. China

Email: ¹chenyc@tju.edu.cn, ²kenzheng@aliyun.com

Abstract

Let $p(n)$ denote the partition function and let Δ be the difference operator respect to n . In this paper, we obtain a lower bound for $\Delta^2 \log \sqrt[n-1]{p(n-1)/(n-1)}$, leading to a proof of the conjecture of Sun on the log-convexity of $\{\sqrt[n]{p(n)/n}\}_{n \geq 60}$. From the log-convexity of both $\{\sqrt[n]{p(n)/n}\}_{n \geq 60}$ and $\{\sqrt[n]{n}\}_{n \geq 4}$, we are led to a proof of another conjecture of Sun on the log-convexity of $\{\sqrt[n]{p(n)}\}_{n \geq 26}$. Using the same argument, it can be shown that for any real number α , there exists an integer $n(\alpha)$ such that the sequence $\{\sqrt[n]{p(n)/n^\alpha}\}_{n \geq n(\alpha)}$ is log-convex. Moreover, we show that $\lim_{n \rightarrow +\infty} n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)} = 3\pi/\sqrt{24}$. Finally, by finding an upper bound of $\Delta^2 \log \sqrt[n-1]{p(n-1)}$, we establish an inequality on the ratio $\frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}$.

Keywords: partition function, log-convex sequence, Hardy-Ramanujan-Rademacher formula, Lehmer's error bound

AMS Subject Classifications: 05A20

1 Introduction

The objective of this paper is to study the log-behavior of the sequences $\sqrt[n]{p(n)}$ and $\sqrt[n]{p(n)/n}$, where $p(n)$ denotes the number of partitions of a nonnegative integer n . A positive sequence $\{a_n\}_{n \geq 0}$ is log-convex if it satisfies that for $n \geq 1$,

$$a_n^2 - a_{n-1}a_{n+1} \leq 0,$$

and it is called log-concave if it satisfies that for $n \geq 1$,

$$a_n^2 - a_{n-1}a_{n+1} \geq 0.$$

Let $r(n) = \sqrt[n]{p(n)/n}$ and let Δ be the difference operator respect to n . Sun [11] conjectured that the sequence $\{r(n)\}_{n \geq 60}$ is log-convex. Desalvo and Pak [5] noticed that the log-convexity of $\{r(n)\}_{n \geq 60}$ can be derived from an estimate for $\Delta^2 \log r(n-1)$, see [5, Final Remark 7.7]. They also remarked that their approach to bounding $-\Delta^2 \log p(n-1)$ does not seem to apply to $\Delta^2 \log r(n-1)$. In this paper, we obtain a lower bound for $\Delta^2 \log r(n-1)$, leading to a proof of the log-convexity of $\{r(n)\}_{n \geq 60}$.

Theorem 1.1 *The sequence $\{r(n)\}_{n \geq 60}$ is log-convex.*

The log-convexity of $\{r(n)\}_{n \geq 60}$ implies the log-convexity of $\{\sqrt[n]{p(n)}\}_{n \geq 26}$, because the sequence $\{\sqrt[n]{n}\}_{n \geq 4}$ is log-convex [11]. It is known that $\lim_{n \rightarrow +\infty} \sqrt[n]{p(n)} = 1$. For a combinatorial proof of this fact, see Andrews [1]. Sun [11] proposed the conjecture that $\{\sqrt[n]{p(n)}\}_{n \geq 6}$ is strictly decreasing, which has been proved by Wang and Zhu [12]. The log-convexity of $\{\sqrt[n]{p(n)}\}_{n \geq 26}$ was also conjectured by Sun [11]. It is easy to see that the log-convexity of $\{\sqrt[n]{p(n)}\}_{n \geq 26}$ implies the decreasing property.

It should be noted that there is an alternative way to prove the log-convexity of $\{\sqrt[n]{p(n)}\}_{n \geq 26}$. Chen, Guo and Wang [3] introduced the notion of a ratio log-convex sequence and showed that the ratio log-convexity implies the log-convexity under a certain initial condition. A sequence $\{a_n\}_{n \geq k}$ is called ratio log-convex if $\{a_{n+1}/a_n\}_{n \geq k}$ is log-convex, or, equivalently, for $n \geq k+1$,

$$\log a_{n+2} - 3 \log a_{n+1} + 3 \log a_n - \log a_{n-1} \geq 0.$$

Chen, Wang and Xie [4] showed that for any $r \geq 1$, one can determine a number $n(r)$ such that for $n > n(r)$, $(-1)^{r-1} \Delta^r \log p(n)$ is positive. For $r = 3$, it can be shown that for $n \geq 116$,

$$\Delta^3 \log p(n-1) > 0.$$

Since

$$\Delta^3 \log p(n-1) = \log p(n+2) - 3 \log p(n+1) + 3 \log p(n) - \log p(n-1),$$

it is evident that $\{p(n)\}_{n \geq 116}$ is ratio log-convex. So we are led to the following assertion.

Theorem 1.2 *The sequence $\{\sqrt[n]{p(n)}\}_{n \geq 26}$ is log-convex.*

Moreover, as pointed out by a referee, we may consider the log-behavior of $\sqrt[n]{p(n)}/n^\alpha$ for any real number α . To this end, we obtain the following generalization of Theorem 1.1 and 1.2.

Theorem 1.3 *Let α be a real number. There exists a positive integer $n(\alpha)$ such that the sequence $\{\sqrt[n]{p(n)}/n^\alpha\}_{n \geq n(\alpha)}$ is log-convex.*

We also establish the following inequality on the ratio $\frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}$.

Theorem 1.4 *For $n \geq 2$, we have*

$$\frac{\sqrt[n]{p(n)}}{\sqrt[n+1]{p(n+1)}} \left(1 + \frac{3\pi}{\sqrt{24}n^{5/2}} \right) > \frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}. \quad (1.1)$$

Desalvo and Pak [5] have shown that the limit of $-n^{\frac{3}{2}} \Delta^2 \log p(n)$ is $\pi/\sqrt{24}$. By bounding $\Delta^2 \log \sqrt[n]{p(n)}$, we derive the following limit of $n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)}$:

$$\lim_{n \rightarrow +\infty} n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)} = 3\pi/\sqrt{24}. \quad (1.2)$$

From the above relation (1.2), it can be seen that the coefficient $\frac{3\pi}{\sqrt{24}}$ in (1.1) is the best possible.

2 The Log-convexity of $r(n)$

In this section, we obtain a lower bound of $\Delta^2 \log r(n-1)$ and prove the log-convexity of $\{r(n)\}_{n \geq 60}$. First, we follow the approach of Desalvo and Pak to give an expression of $\Delta^2 \log r(n-1)$ as a sum of $\Delta^2 \tilde{B}(n-1)$ and $\Delta^2 \tilde{E}(n-1)$, where $\Delta^2 \tilde{B}(n-1)$ makes a major contribution to $\Delta^2 \log r(n-1)$ with $\Delta^2 \tilde{E}(n-1)$ being the error term, that is, $\Delta^2 \tilde{B}(n-1)$ converges to $\Delta^2 \log r(n-1)$. The expressions for $B(n)$ and $E(n)$ will be given later. In this setting, we derive a lower bound of $\Delta^2 \tilde{B}(n-1)$. By Lehmer's error bound, we give an upper bound for $|\Delta^2 \tilde{E}(n-1)|$. Combining the lower bound for $\Delta^2 \tilde{B}(n-1)$ and the upper bound for $\Delta^2 \tilde{E}(n-1)$, we are led to a lower bound for $\Delta^2 \log r(n-1)$. By proving the positivity of this lower bound for $\Delta^2 \log r(n-1)$, we reach the log-convexity of $\{r(n)\}_{n \geq 60}$.

The strict log-convexity of $\{r(n)\}_{n \geq 60}$ can be restated as the following relation for $n \geq 61$,

$$\log r(n+1) + \log r(n-1) - 2 \log r(n) > 0,$$

that is, for $n \geq 61$,

$$\Delta^2 \log r(n-1) > 0.$$

For $n \geq 1$ and any positive integer N , the Hardy-Ramanujan-Rademacher formula (see [2, 6, 7, 10]) reads

$$p(n) = \frac{d}{\mu^2} \sum_{k=1}^N A_k^*(n) \left[\left(1 - \frac{k}{\mu}\right) e^{\frac{\mu}{k}} + \left(1 + \frac{k}{\mu}\right) e^{-\frac{\mu}{k}} \right] + R_2(n, N), \quad (2.1)$$

where $d = \frac{\pi^2}{6\sqrt{3}}$, $\mu(n) = \frac{\pi}{6} \sqrt{24n-1}$, $A_k^*(n) = k^{-\frac{1}{2}} A_k(n)$, $A_k(n)$ is a sum of 24th roots of unity with initial values $A_1(n) = 1$ and $A_2(n) = (-1)^n$, $R_2(n, N)$ is the remainder. Lehmer's error bound (see [8, 9]) for $R_2(n, N)$ is given by

$$|R_2(n, N)| < \frac{\pi^2 N^{-2/3}}{\sqrt{3}} \left[\left(\frac{N}{\mu}\right)^3 \sinh \frac{\mu}{N} + \frac{1}{6} - \left(\frac{N}{\mu}\right)^2 \right]. \quad (2.2)$$

Let us give an outline of Desalvo and Pak's approach to proving the log-concavity of $\{p(n)\}_{n > 25}$. Setting $N = 2$ in (2.1), they expressed $p(n)$ as

$$p(n) = T(n) + R(n), \quad (2.3)$$

where

$$\begin{aligned} T(n) &= \frac{d}{\mu(n)^2} \left[\left(1 - \frac{1}{\mu(n)}\right) e^{\mu(n)} + \frac{(-1)^n}{\sqrt{2}} e^{\frac{\mu(n)}{2}} \right], \\ R(n) &= \frac{d}{\mu(n)^2} \left[\left(1 + \frac{1}{\mu(n)}\right) e^{-\mu(n)} - \frac{(-1)^n}{\sqrt{2}} \frac{2}{\mu(n)} + \frac{(-1)^n}{\sqrt{2}} \left(1 + \frac{2}{\mu(n)}\right) e^{-\frac{\mu(n)}{2}} \right] + R_2(n, 2). \end{aligned} \quad (2.4)$$

$$(2.5)$$

They have shown that

$$\left| \Delta^2 \log p(n-1) - \Delta^2 \log T(n-1) \right| = \left| \Delta^2 \log \left(1 + \frac{R(n-1)}{T(n-1)} \right) \right| < e^{-\frac{\pi\sqrt{2n}}{10\sqrt{3}}}. \quad (2.6)$$

and

$$\left| \Delta^2 \log T(n-1) - \Delta^2 \log \frac{d}{\mu(n-1)^2} \left(1 - \frac{1}{\mu(n-1)} \right) e^{\mu(n-1)} \right| < e^{-\frac{\pi\sqrt{2n}}{10\sqrt{3}}}. \quad (2.7)$$

It follows that $\Delta^2 \log \frac{d}{\mu(n-1)^2} \left(1 - \frac{1}{\mu(n-1)} \right) e^{\mu(n-1)}$ converges to $\Delta^2 \log p(n-1)$. Finally, they use $-\Delta^2 \log \frac{d}{\mu(n-1)^2} \left(1 - \frac{1}{\mu(n-1)} \right) e^{\mu(n-1)}$ to estimate $-\Delta^2 \log p(n-1)$, leading to the log-concavity of $\{p(n)\}_{n>25}$.

We shall use an alternative decomposition of $p(n)$. Setting $N = 2$ in (2.1), we can express $p(n)$ as

$$p(n) = \tilde{T}(n) + \tilde{R}(n), \quad (2.8)$$

where

$$\tilde{T}(n) = \frac{d}{\mu(n)^2} \left(1 - \frac{1}{\mu(n)} \right) e^{\mu(n)}, \quad (2.9)$$

$$\begin{aligned} \tilde{R}(n) = \frac{d}{\mu(n)^2} & \left[\left(1 + \frac{1}{\mu(n)} \right) e^{-\mu(n)} + \frac{(-1)^n}{\sqrt{2}} \left(1 - \frac{2}{\mu(n)} \right) e^{\frac{\mu(n)}{2}} \right. \\ & \left. + \frac{(-1)^n}{\sqrt{2}} \left(1 + \frac{2}{\mu(n)} \right) e^{-\frac{\mu(n)}{2}} \right] + R_2(n, 2). \end{aligned} \quad (2.10)$$

Based on the decomposition (2.8) for $p(n)$, one can express $\Delta^2 \log r(n-1)$ as follows:

$$\Delta^2 \log r(n-1) = \Delta^2 \tilde{B}(n-1) + \Delta^2 \tilde{E}(n-1), \quad (2.11)$$

where

$$\tilde{B}(n) = \frac{1}{n} \log \tilde{T}(n) - \frac{1}{n} \log n, \quad (2.12)$$

$$\tilde{y}_n = \tilde{R}(n)/\tilde{T}(n), \quad (2.13)$$

$$\tilde{E}(n) = \frac{1}{n} \log(1 + \tilde{y}_n). \quad (2.14)$$

The following lemma will be used to derive a lower bound and an upper bound of $\Delta^2 \tilde{B}(n-1)$.

Lemma 2.1 *Suppose $f(x)$ has a continuous second derivative for $x \in [n-1, n+1]$. Then there exists $c \in (n-1, n+1)$ such that*

$$\Delta^2 f(n-1) = f(n+1) + f(n-1) - 2f(n) = f''(c). \quad (2.15)$$

If $f(x)$ has an increasing second derivative, then

$$f''(n-1) < \Delta^2 f(n-1) < f''(n+1). \quad (2.16)$$

Conversely, if $f(x)$ has a decreasing second derivative, then

$$f''(n+1) < \Delta^2 f(n-1) < f''(n-1). \quad (2.17)$$

Proof. Set $\varphi(x) = f(x+1) - f(x)$. By the mean value theorem, there exists a number $\xi \in (n-1, n)$ such that

$$f(n+1) + f(n-1) - 2f(n) = \varphi(n) - \varphi(n-1) = \varphi'(\xi).$$

Again, applying the mean value theorem to $\varphi'(\xi)$, there exists a number $\theta \in (0, 1)$ such that

$$\varphi'(\xi) = f'(\xi+1) - f'(\xi) = f''(\xi+\theta).$$

Let $c = \xi + \theta$. Then we get (2.15), which yields (2.16) and (2.17). ■

In order to find a lower bound for $\Delta^2 \log r(n-1)$ and obtain the limit of $n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)}$, we need the following lower and upper bounds for $\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$.

Lemma 2.2 *Let*

$$B_1(n) = \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4 \log(\mu(n-1))}{(n-1)^3}, \quad (2.18)$$

$$B_2(n) = \frac{72\pi}{(n-1)(24n-25)^{3/2}} - \frac{4 \log(\mu(n+1))}{(n+1)^3} + \frac{5}{(n-1)^3}. \quad (2.19)$$

For $n \geq 40$, we have

$$B_1(n) < \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) < B_2(n). \quad (2.20)$$

Proof. By the definition (2.9), we may write

$$\frac{\log \tilde{T}(n)}{n} = \sum_{i=1}^4 f_i,$$

where

$$\begin{aligned} f_1(n) &= \frac{\mu(n)}{n}, \\ f_2(n) &= -\frac{3 \log \mu(n)}{n}, \\ f_3(n) &= \frac{\log(\mu(n)-1)}{n}, \\ f_4(n) &= \frac{\log d}{n}. \end{aligned}$$

Thus

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) = \sum_{i=1}^4 \Delta^2 f_i(n-1). \quad (2.21)$$

Since

$$f_1'''(n) = \frac{\pi}{n(24n-1)^{3/2}} \left(-\frac{216}{n} + \frac{864}{24n-1} + \frac{36}{n^2} - \frac{1}{n^3} \right),$$

we see that for $n \geq 1$, $f_1'''(n) < 0$. Similarly, it can be checked that for $n \geq 4$, $f_2'''(n) > 0$, $f_3'''(n) < 0$, and $f_4'''(n) > 0$. Consequently, for $n \geq 4$, $f_1''(n)$ and $f_3''(n)$ are decreasing, whereas $f_2''(n)$ and $f_4''(n)$ are increasing. Using Lemma 2.1, for each i , we can get a lower bound and an upper bound for $\Delta^2 f_i(n-1)$ in terms of $f_i''(n-1)$ and $f_i''(n+1)$. For example,

$$f_1''(n+1) < \Delta^2 f_1(n-1) < f_1''(n-1).$$

So, by (2.21) we find that

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) > f_1''(n+1) + f_2''(n-1) + f_3''(n+1) + f_4''(n-1), \quad (2.22)$$

and

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) < f_1''(n-1) + f_2''(n+1) + f_3''(n-1) + f_4''(n+1), \quad (2.23)$$

where

$$f_1''(n) = \frac{72\pi}{n(24n-1)^{3/2}} - \frac{12\pi}{n^2(24n-1)^{3/2}} + \frac{\pi}{3n^3(24n-1)^{3/2}}, \quad (2.24)$$

$$f_2''(n) = -\frac{6 \log \mu(n)}{n^3} + \frac{72}{(24n-1)n^2} + \frac{864}{n(24n-1)^2}, \quad (2.25)$$

$$f_3''(n) = -\frac{4\pi^2}{(\mu(n)-1)^2(24n-1)n} + \frac{2 \log(\mu(n)-1)}{n^3} \\ - \frac{4\pi}{(\mu(n)-1)\sqrt{24n-1}n^2} - \frac{24\pi}{(\mu(n)-1)(24n-1)^{3/2}n}, \quad (2.26)$$

$$f_4''(n) = \frac{2 \log d}{n^3}. \quad (2.27)$$

According to (2.24), one can check that for $n \geq 2$,

$$f_1''(n+1) > \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{12\pi}{(n+1)^2(24n+23)^{3/2}}. \quad (2.28)$$

An easy computation shows that for $n \geq 3$,

$$\mu(n) - 1 > \frac{2}{3}\mu(n-2). \quad (2.29)$$

Substituting (2.29) into (2.26) yields that

$$f_3''(n+1) > \frac{2 \log(\mu(n+1)-1)}{(n+1)^3} - \frac{540}{(24n-25)^2(n-1)} - \frac{36}{(24n-25)(n-1)^2}. \quad (2.30)$$

Using (2.25) and (2.30), we find that

$$f_2''(n-1) + f_3''(n+1) \\ > \frac{2 \log(\mu(n+1)-1)}{(n+1)^3} - \frac{6 \log(\mu(n-1))}{(n-1)^3}$$

$$+ \frac{324}{(n-1)(24n-25)^2} + \frac{36}{(n-1)^2(24n-25)} \quad (2.31)$$

Apparently, for $n \geq 2$,

$$\frac{2}{(n+1)^3} - \frac{2}{(n-1)^3} > -\frac{12}{(n-1)^4},$$

so that

$$\begin{aligned} & \frac{2 \log(\mu(n+1)-1)}{(n+1)^3} - \frac{6 \log(\mu(n-1))}{(n-1)^3} \\ & > \frac{2 \log(\mu(n+1)-1)}{(n+1)^3} - \frac{2 \log(\mu(n+1)-1)}{(n-1)^3} - \frac{4 \log(\mu(n-1))}{(n-1)^3} \\ & > -\frac{12 \log(\mu(n+1)-1)}{(n-1)^4} - \frac{4 \log(\mu(n-1))}{(n-1)^3}. \end{aligned} \quad (2.32)$$

Since, for $n \geq 2$,

$$\frac{324}{(n-1)(24n-25)^2} + \frac{36}{(n-1)^2(24n-25)} > \frac{2}{(n-1)^3}, \quad (2.33)$$

utilizing (2.31) and (2.32) yields that for $n \geq 3$,

$$f_2''(n-1) + f_3''(n+1) > -\frac{4 \log(\mu(n-1))}{(n-1)^3} + \frac{2}{(n-1)^3} - \frac{12 \log(\mu(n+1)-1)}{(n-1)^4}. \quad (2.34)$$

Using (2.27), (2.28) and (2.34), we deduce that

$$\begin{aligned} & f_1''(n+1) + f_2''(n-1) + f_3''(n+1) + f_4''(n-1) - B_1(n) \\ & > \frac{2(1+\log d)}{(n-1)^3} - \frac{12\pi}{(n+1)^2(24n+23)^{3/2}} - \frac{12 \log(\mu(n+1)-1)}{(n-1)^4}. \end{aligned} \quad (2.35)$$

Let $C(n)$ be the right hand side of (2.35). To prove (2.22), it is enough to show that $C(n) > 0$ when $n \geq 40$. Since $\log x < x$ for $x > 0$, and for $n \geq 3$

$$\mu(n+1) - 1 < \frac{\pi}{4} \sqrt{24n-24}, \quad (2.36)$$

we get

$$-\frac{12 \log(\mu(n+1)-1)}{(n-1)^4} > -\frac{12(\mu(n+1)-1)}{(n-1)^4} > -\frac{3\sqrt{24}\pi}{(n-1)^{7/2}}. \quad (2.37)$$

Note that for $n \geq 2$,

$$-\frac{12\pi}{(n+1)^2(24n+23)^{3/2}} > -\frac{\sqrt{24}\pi}{48(n-1)^{7/2}}. \quad (2.38)$$

Combining (2.37) and (2.38) gives for $n \geq 2$,

$$C(n) > \frac{2(1+\log d)}{(n-1)^3} - \frac{(3+1/48)\sqrt{24}\pi}{(n-1)^{7/2}}. \quad (2.39)$$

It is straightforward to show that the right hand side of (2.39) is positive if $n \geq 490$. For $40 \leq n \leq 489$, it is routine to check that $C(n) > 0$, and so $C(n) > 0$ for $n \geq 40$. It follows from (2.35) that for $n \geq 40$,

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) > B_1(n).$$

To derive the upper bound for $\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$, we obtain the following upper bounds which can be verified directly. The proofs are omitted. For $n \geq 2$,

$$\begin{aligned} f_1''(n-1) &< \frac{72\pi}{(n-1)[24n-25]^{3/2}}, \\ f_2''(n+1) &< -\frac{6 \log \mu(n+1)}{(n+1)^3} + \frac{9}{2(n-1)^3}, \\ f_3''(n-1) &< -\frac{4\pi^2}{(\mu(n-1))^2(24n-25)(n-1)} + \frac{2 \log(\mu(n-1))}{(n-1)^3} \\ &\quad - \frac{4\pi}{\mu(n-1)\sqrt{24n-25}(n-1)^2} - \frac{24\pi}{\mu(n-1)(24n-25)^{3/2}(n-1)}, \\ f_2''(n+1) + f_3''(n-1) &< \frac{3}{(n-1)^3} + \frac{12 \log(\mu(n+1))}{(n-1)^4} - \frac{4 \log(\mu(n+1))}{(n+1)^3}, \\ f_4''(n+1) &< 0. \end{aligned}$$

Combining the above upper bounds, we conclude that for $n \geq 40$,

$$f_1''(n-1) + f_2''(n+1) + f_3''(n-1) + f_4''(n+1) < B_2(n).$$

This completes the proof. ■

The following lemma gives an upper bound for $|\Delta^2 \tilde{E}(n-1)|$.

Lemma 2.3 For $n \geq 40$,

$$|\Delta^2 \tilde{E}(n-1)| < \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \quad (2.40)$$

Proof. By (2.14), we find that for $n \geq 2$,

$$\Delta^2 \tilde{E}(n-1) = \frac{1}{n-1} \log(1 + \tilde{y}_{n-1}) + \frac{1}{n+1} \log(1 + \tilde{y}_{n+1}) - \frac{2}{n} \log(1 + \tilde{y}_n), \quad (2.41)$$

where

$$\tilde{y}_n = \tilde{R}(n)/\tilde{T}(n).$$

To bound $|\Delta^2 \tilde{E}(n-1)|$, it is necessary to bound \tilde{y}_n . For this purpose, we first consider $\tilde{R}(n)$, as defined by (2.10). Since $d < 1$ and $\mu(n) > 2$, for $n \geq 1$ we have

$$\frac{d}{\mu(n)^2} \left[\left(1 + \frac{1}{\mu(n)}\right) e^{-\mu(n)} + \frac{(-1)^n}{\sqrt{2}} \left(1 - \frac{2}{\mu(n)}\right) e^{\frac{\mu(n)}{2}} + \frac{(-1)^n}{\sqrt{2}} \left(1 + \frac{2}{\mu(n)}\right) e^{-\frac{\mu(n)}{2}} \right]$$

$$< \frac{1}{\mu(n)^2} \left(1 + e^{\frac{\mu(n)}{2}} + 1\right).$$

For $N = 2$ and $n \geq 1$, Lehmer's bound (2.2) reduces to

$$|R_2(n, 2)| < 4 \left(1 + \frac{4}{\mu(n)^3} e^{\frac{\mu(n)}{2}}\right).$$

By the definition of $\tilde{R}(n)$,

$$|\tilde{R}(n)| < \frac{1}{\mu(n)^2} \left(1 + e^{\frac{\mu(n)}{2}} + 1\right) + 4 \left(1 + \frac{4}{\mu(n)^3} e^{\frac{\mu(n)}{2}}\right) < 5 + \frac{9}{\mu(n)^2} e^{\frac{\mu(n)}{2}}. \quad (2.42)$$

Recalling the definition (2.9) of $\tilde{T}(n)$, it follows from (2.42) that for $n \geq 1$,

$$|\tilde{y}_n| < \frac{\mu(n)}{d(\mu(n) - 1)} \left(5\mu(n)^2 e^{-\frac{2\mu(n)}{3}} + 9e^{-\frac{\mu(n)}{6}}\right) e^{-\frac{\mu(n)}{3}}. \quad (2.43)$$

Observe that for $n \geq 2$,

$$\left(5\mu(n)^2 e^{-\frac{2\mu(n)}{3}} + 9e^{-\frac{\mu(n)}{6}}\right)' < 0, \quad (2.44)$$

and

$$\left(\frac{d(\mu(n) - 1)}{\mu(n)}\right)' > 0. \quad (2.45)$$

Since

$$5\mu^2(40)e^{-\frac{2\mu(40)}{3}} + 9e^{-\frac{\mu(40)}{6}} < \frac{d(\mu(40) - 1)}{\mu(40)},$$

using (2.44) and (2.45), we deduce that for $n \geq 40$,

$$5\mu^2(n)e^{-\frac{2\mu(n)}{3}} + 9e^{-\frac{\mu(n)}{6}} < \frac{d(\mu(n) - 1)}{\mu(n)}. \quad (2.46)$$

Now, it is clear from (2.43) and (2.46) that for $n \geq 40$,

$$|\tilde{y}_n| < e^{-\frac{\mu(n)}{3}}. \quad (2.47)$$

In view of (2.47), for $n \geq 40$,

$$|\tilde{y}_n| < e^{-\frac{\mu(40)}{3}} < \frac{1}{5}. \quad (2.48)$$

It is known that $\log(1 + x) < x$ for $0 < x < 1$ and $-\log(1 + x) < -x/(1 + x)$ for $-1 < x < 0$. Thus, for $|x| < 1$,

$$|\log(1 + x)| \leq \frac{|x|}{1 - |x|}, \quad (2.49)$$

see also [5], and so it follows from (2.48) and (2.49) that for $n \geq 40$,

$$|\log(1 + \tilde{y}_n)| \leq \frac{|\tilde{y}_n|}{1 - |\tilde{y}_n|} \leq \frac{5}{4} |\tilde{y}_n|. \quad (2.50)$$

Because of (2.41), we see that for $n \geq 2$,

$$\left| \Delta^2 \tilde{E}(n-1) \right| \leq \frac{1}{n-1} |\log(1+\tilde{y}_{n-1})| + \frac{1}{n+1} |\log(1+\tilde{y}_{n+1})| + \frac{2}{n} |\log(1+\tilde{y}_n)|. \quad (2.51)$$

Applying (2.50) to (2.51), we obtain that for $n \geq 40$,

$$\left| \Delta^2 \tilde{E}(n-1) \right| \leq \frac{5}{4} \left(\frac{|\tilde{y}_{n-1}|}{n-1} + \frac{|\tilde{y}_{n+1}|}{n+1} + \frac{2|\tilde{y}_n|}{n} \right). \quad (2.52)$$

Plugging (2.47) into (2.52), we infer that for $n \geq 40$,

$$\left| \Delta^2 \tilde{E}(n-1) \right| < \frac{5}{4} \left(\frac{e^{-\frac{\mu(n-1)}{3}}}{n-1} + \frac{e^{-\frac{\mu(n+1)}{3}}}{n+1} + \frac{2e^{-\frac{\mu(n)}{3}}}{n} \right). \quad (2.53)$$

But $\frac{1}{n}e^{-\frac{\mu(n)}{3}}$ is decreasing for $n \geq 1$, it follows from (2.53) that for $n \geq 40$,

$$\left| \Delta^2 \tilde{E}(n-1) \right| < \frac{5}{n-1} e^{-\frac{\mu(n-1)}{3}}.$$

This proves (2.40). ■

With the aid of Lemma 2.2 and 2.3, we are ready to prove the log-convexity of $\{r(n)\}_{n \geq 60}$.

Proof of Theorem 1.1. To prove the strict log-convexity of $\{r(n)\}_{n \geq 60}$, we proceed to show that for $n \geq 61$,

$$\Delta^2 \log r(n-1) > 0.$$

Evidently, for $n \geq 40$,

$$\left(-\frac{\log n}{n} \right)''' > 0.$$

By Lemma 2.1,

$$-\Delta^2 \frac{\log(n-1)}{n-1} > \left(-\frac{\log(n-1)}{n-1} \right)'' ,$$

that is,

$$-\Delta^2 \frac{\log(n-1)}{n-1} > -\frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3}. \quad (2.54)$$

It follows from (2.12) that

$$\Delta^2 \tilde{B}(n-1) = \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) - \Delta^2 \frac{\log(n-1)}{n-1}.$$

Applying Lemma 2.2 and (2.54) to the above relation, we deduce that for $n \geq 40$,

$$\Delta^2 \tilde{B}(n-1) > \tilde{B}_1(n) - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3},$$

that is,

$$\Delta^2 \tilde{B}(n-1) > \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3}. \quad (2.55)$$

By (2.11) and Lemma 2.3, we find that for $n \geq 40$,

$$\Delta^2 \log r(n-1) > \Delta^2 \tilde{B}(n-1) - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \quad (2.56)$$

It follows from (2.55) and (2.56) that for $n \geq 40$,

$$\begin{aligned} & \Delta^2 \log r(n-1) \\ & > \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4\log[\mu(n-1)]}{(n-1)^3} - \frac{2\log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \end{aligned}$$

Let $D(n)$ denote the right hand side of the above relation. Clearly, for $n \geq 5505$,

$$\frac{72\pi}{(n+1)(24n+23)^{3/2}} > \frac{3\pi}{\sqrt{24}(n+1)^{5/2}} > \frac{1}{(n-1)^{5/2}}. \quad (2.57)$$

To prove that $D(n) > 0$ for $n \geq 5505$, we wish to show that for $n \geq 5505$,

$$-\frac{4\log[\mu(n-1)]}{(n-1)^3} - \frac{2\log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} > -\frac{1}{(n-1)^{5/2}}. \quad (2.58)$$

Using the fact that for $x > 5504$, $\log x < x^{1/4}$, we deduce that for $n \geq 5505$,

$$\frac{4\log[\mu(n-1)]}{(n-1)^3} < \frac{4\sqrt[4]{\mu(n-1)}}{(n-1)^3} < \frac{4\sqrt[4]{\frac{\pi}{4}\sqrt{24n-24}}}{(n-1)^3} < \frac{6}{(n-1)^{23/8}}, \quad (2.59)$$

and

$$\frac{2\log(n-1)}{(n-1)^3} < \frac{2(n-1)^{1/4}}{(n-1)^3} < \frac{2}{(n-1)^{11/4}}. \quad (2.60)$$

Since $e^x > x^6/720$ for $x > 0$, we see that for $n \geq 2$,

$$\frac{1}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} < \frac{1}{n-1} e^{-\frac{\pi\sqrt{23n}}{18}} < \frac{2094}{n^3(n-1)} < \frac{2094}{(n-1)^4}. \quad (2.61)$$

Combining (2.59), (2.60) and (2.61), we find that for $n \geq 5505$,

$$\begin{aligned} & -\frac{4\log[\mu(n-1)]}{(n-1)^3} - \frac{2\log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} \\ & > -\frac{6}{(n-1)^{23/8}} - \frac{2}{(n-1)^{11/4}} + \frac{3}{(n-1)^3} - \frac{10470}{(n-1)^4} \\ & > -\frac{6}{(n-1)^{23/8}} - \frac{2}{(n-1)^{11/4}} \\ & > -\frac{1}{(n-1)^{5/2}}. \end{aligned}$$

This proves the inequality (2.58). By (2.58) and (2.57), we obtain that $D(n) > 0$ for $n \geq 5505$. Verifying that $\Delta^2 \log r(n-1) > 0$ for $61 \leq n \leq 5504$ completes the proof. \blacksquare

Clearly, Theorem 1.3 is a generalization as well as a unification of Theorem 1.1 and 1.2. In fact, it can be proved in the same manner as the proof of Theorem 1.1.

Proof of Theorem 1.3. Let α be a real number. When $\alpha \leq 0$, it is clear that $\frac{1}{\sqrt[n]{n^\alpha}}$ is log-convex. It follows from Theorem 1.2 that $\sqrt[n]{p(n)/n^\alpha}$ is log-convex for $n \geq 26$.

We now consider the case $\alpha > 0$. A similar argument to the proof of Theorem 1.1 shows that for $n \geq 40$,

$$\begin{aligned} & \Delta^2 \log \sqrt[n-1]{p(n-1)/(n-1)^\alpha} \\ &= \Delta^2 \frac{1}{n-1} \log T(n) + \Delta^2 \frac{1}{n-1} \log(1 + y_{n-1}) - \alpha \Delta^2 \frac{\log(n-1)}{n-1} \\ &> \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2\alpha \log(n-1)}{(n-1)^3} \\ & \quad + \frac{3\alpha}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \end{aligned} \tag{2.62}$$

It is easy to check that for $n \geq \max\left\{\left\lceil\frac{3490}{\alpha}\right\rceil + 2, 5505\right\}$,

$$\frac{3\alpha}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} > \frac{3\alpha}{(n-1)^3} - \frac{10470}{(n-1)^4} > 0,$$

and for $n \geq \max\{[(2\alpha+3)^4] + 2, 5505\}$,

$$-\frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2\alpha \log(n-1)}{(n-1)^3} > -\frac{6}{(n-1)^{23/8}} - \frac{2\alpha}{(n-1)^{11/4}} > -\frac{1}{(n-1)^{5/2}}.$$

Let

$$n(\alpha) = \max\left\{\left\lceil\frac{3490}{\alpha}\right\rceil + 2, [(2\alpha+3)^4] + 2, 5505\right\}.$$

It can be seen that for $n > n(\alpha)$,

$$-\frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2\alpha \log(n-1)}{(n-1)^3} + \frac{3\alpha}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} > -\frac{1}{(n-1)^{5/2}}. \tag{2.63}$$

Combing (2.57) and (2.63), we deduce that the right hand side of (2.62) is positive for $n > n(\alpha)$. So we are led to the log-convexity of the sequence $\{\sqrt[n]{p(n)/n^\alpha}\}_{n \geq n(\alpha)}$. \blacksquare

3 An inequality on the ratio $\frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}$

In this section, we employ Lemma 2.2 and Lemma 2.3 to find the limit of $n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)}$. Then we give an upper bound for $\Delta^2 \log \sqrt[n-1]{p(n-1)}$. This leads to the inequality (1.1).

Theorem 3.1 *Let $\beta = 3\pi/\sqrt{24}$. We have*

$$\lim_{n \rightarrow +\infty} n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)} = \beta. \tag{3.1}$$

Proof. Using (2.8), that is, the $N = 2$ case of the Hardy-Ramanujan-Rademacher formula for $p(n)$, we find that

$$\log \sqrt[n]{p(n)} = \frac{1}{n} \log \tilde{T}(n) + \frac{1}{n} \log(1 + \tilde{y}_n),$$

where $\tilde{T}(n)$ and y_n are given by (2.9) and (2.13). By the definition (2.14) of $\tilde{E}(n)$, we get

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} = \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) + \Delta^2 \tilde{E}(n-1). \quad (3.2)$$

Applying Lemma 2.2, we get that

$$\lim_{n \rightarrow +\infty} (n-1)^{\frac{5}{2}} \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) = \beta. \quad (3.3)$$

From Lemma 2.3, we get

$$\lim_{n \rightarrow +\infty} (n-1)^{\frac{5}{2}} \Delta^2 \tilde{E}(n-1) = 0. \quad (3.4)$$

Using (3.2), (3.3) and (3.4), we deduce that

$$\lim_{n \rightarrow +\infty} n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)} = \beta,$$

as required. ■

To prove Theorem 1.4, we need the following upper bound for $\Delta^2 \log \sqrt[n-1]{p(n-1)}$.

Theorem 3.2 *For $n \geq 2$,*

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} < \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi}. \quad (3.5)$$

Proof. By the upper bound of $\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$ given in Lemma 2.2, the upper bound of $\Delta^2 \tilde{E}(n-1)$ given in Lemma 2.3 and the relation (3.2), we get the following upper bound of $\Delta^2 \log \sqrt[n-1]{p(n-1)}$ for $n \geq 40$,

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} < \frac{72\pi}{(n-1)(24n-25)^{3/2}} + \frac{5}{(n-1)^3} - \frac{4 \log[\mu(n+1)]}{(n+1)^3} + \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}.$$

To prove (3.5), we claim that for $n \geq 2095$,

$$\frac{72\pi}{(n-1)(24n-25)^{3/2}} + \frac{5}{(n-1)^3} - \frac{4 \log[\mu(n+1)]}{(n+1)^3} + \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} < \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi}. \quad (3.6)$$

First, we show that for $n \geq 60$,

$$\frac{72\pi}{(n-1)(24n-25)^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} < \frac{1}{(n-1)^3}. \quad (3.7)$$

For $0 < x \leq \frac{1}{48}$, it can be checked that

$$\frac{1}{(1-x)^{3/2}} < 1 + \frac{3}{2}x + \frac{3}{8}x^{\frac{3}{2}}. \quad (3.8)$$

In the notation $\beta = 3\pi/\sqrt{24}$, we have

$$\frac{72\pi}{(n-1)(24n-25)^{3/2}} = \frac{\beta}{(n-1)n^{3/2}(1-\frac{25}{24n})^{3/2}}. \quad (3.9)$$

Setting $x = \frac{25}{24n}$, we have $x \leq \frac{1}{48}$ for $n \geq 60$. Applying (3.8) to the right hand side of (3.9), we find that for $n \geq 60$,

$$\frac{\beta}{(n-1)n^{3/2}(1-\frac{25}{24n})^{3/2}} < \frac{\beta}{(n-1)n^{3/2}} \left[1 + \frac{75}{48n} + \frac{3}{8} \left(\frac{25}{24n} \right)^{\frac{3}{2}} \right], \quad (3.10)$$

so that for $n \geq 60$,

$$\begin{aligned} & \frac{72\pi}{(n-1)[24n-25]^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} \\ & < \frac{\beta}{(n-1)n^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} + \frac{\beta}{(n-1)n^{3/2}} \left[\frac{75}{48n} + \frac{3}{8} \left(\frac{25}{24n} \right)^{\frac{3}{2}} \right]. \end{aligned} \quad (3.11)$$

To prove (3.7), we proceed to show that the right hand side of (3.11) is bounded by $\frac{1}{(n-1)^3}$. Noted that for $n \geq 2$,

$$\frac{\beta}{(n-1)n^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} = \frac{\beta}{(n^{5/2} + \beta)(n-1)} + \frac{\beta^2}{(n^{5/2} + \beta)(n-1)n^{3/2}},$$

and $n^{5/2} + \beta > (n-1)^{5/2}$, together with $n^{3/2} > (n-1)^{3/2}$, we have that

$$\frac{\beta}{(n-1)n^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} < \frac{\beta}{(n-1)^{7/2}} + \frac{\beta}{(n-1)^5}. \quad (3.12)$$

Applying (3.12) to (3.11), we obtain that for $n \geq 60$,

$$\begin{aligned} & \frac{72\pi}{(n-1)[24n-25]^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} \\ & < \frac{\beta}{(n-1)^{7/2}} + \frac{\beta^2}{(n-1)^5} + \frac{\beta}{(n-1)n^{3/2}} \left[\frac{75}{48n} + \frac{3}{8} \left(\frac{25}{24n} \right)^{\frac{3}{2}} \right]. \end{aligned} \quad (3.13)$$

Since $\frac{75}{48n} < \frac{2}{n-1}$ and $\frac{3}{8} \left(\frac{25}{24n} \right)^{\frac{3}{2}} < \frac{1}{(n-1)^{3/2}}$ for $n \geq 2$, it follows from (3.13) that for $n \geq 60$,

$$\begin{aligned} & \frac{72\pi}{(n-1)[24n-25]^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} \\ & < \frac{\beta}{(n-1)^{7/2}} + \frac{\beta^2}{(n-1)^5} + \frac{2\beta}{(n-1)^{7/2}} + \frac{\beta}{(n-1)^4}. \end{aligned}$$

Using the fact that $\beta < 2$, we see that

$$\frac{3\beta}{(n-1)^{7/2}} + \frac{\beta^2}{(n-1)^5} + \frac{\beta}{(n-1)^4} < \frac{6}{(n-1)^{7/2}} + \frac{4}{(n-1)^5} + \frac{2}{(n-1)^4}. \quad (3.14)$$

For $n \geq 60$, it is easily checked that the right hand side of (3.14) is bounded by $\frac{1}{(n-1)^3}$. This confirms (3.7).

To prove the claim (3.6), it is enough to show that for $n \geq 2095$,

$$\frac{1}{(n-1)^3} < \frac{4 \log[\mu(n+1)]}{(n+1)^3} - \frac{5}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \quad (3.15)$$

From (2.61) it can be seen that for $n \geq 2095$,

$$\frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} < \frac{5}{(n-1)^3}. \quad (3.16)$$

Since $4 \log[\mu(n+1)] > 18$ for $n \geq 2095$, it follows from (3.16) that for $n \geq 2095$,

$$\begin{aligned} & \frac{4 \log[\mu(n+1)]}{(n+1)^3} - \frac{5}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} \\ & > \frac{18}{(n+1)^3} - \frac{10}{(n-1)^3} > \frac{1}{(n-1)^3}. \end{aligned}$$

So we obtain (3.15), which yields (3.6). For $2 \leq n \leq 2094$, the inequality (3.5) can be easily checked. This completes the proof. \blacksquare

We are now in a position to finish the proof of Theorem 1.4.

Proof of Theorem 1.4. It is known that for $x > 0$,

$$\frac{x}{1+x} < \log(1+x),$$

so that for $n \geq 1$,

$$\frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} < \log\left(1 + \frac{3\pi}{\sqrt{24n^{5/2}}}\right).$$

In light of the above relation, Theorem 3.2 implies that for $n \geq 2$,

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} < \log\left(1 + \frac{3\pi}{\sqrt{24n^{5/2}}}\right),$$

that is,

$$\sqrt[n+1]{p(n+1)} \sqrt[n-1]{p(n-1)} < \left(1 + \frac{3\pi}{\sqrt{24n^{5/2}}}\right) (\sqrt[n]{p(n)})^2,$$

as required. \blacksquare

We remark that $\beta = 3\pi/\sqrt{24}$ is the smallest possible number for the inequality in Theorem 1.4. Suppose that $0 < \gamma < \beta$. By Theorem 3.1, there exists an integer N so as to for $n > N$,

$$n^{5/2} \Delta^2 \log \sqrt[n-1]{p(n-1)} > \gamma.$$

It follows that

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} > \frac{\gamma}{n^{5/2}} > \log \left(1 + \frac{\gamma}{n^{5/2}} \right),$$

which implies that for $n > N$,

$$\frac{\sqrt[n]{p(n)}}{\sqrt[n+1]{p(n+1)}} \left(1 + \frac{\gamma}{n^{5/2}} \right) < \frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}.$$

Acknowledgments. We thank the referee for helpful comments. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, and the National Science Foundation of China.

References

- [1] G.E. Andrews, Combinatorial proof of a partition function limit, *Amer. Math. Monthly.*, 78 (1971), 276–278.
- [2] G.E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
- [3] W.Y.C. Chen, J.J.F. Guo and L.X.W. Wang, Infinitely log-monotonic combinatorial sequences. *Adv. Appl. Math.*, 52 (2014), 99–120.
- [4] W.Y.C. Chen, L.X.W. Wang and G.Y.B. Xie, Finite differences of the logarithm of the partition function. *Math. Comp.*, 85 (2016), 825–847.
- [5] S. Desalvo and I. Pak, Log-concavity of the partition function, *Ramanujan J.*, 38 (2016), 61–73.
- [6] G.H. Hardy, *Twelve Lectures on Subjects Suggested by His Life and Work*, Cambridge University Press, Cambridge, 1940.
- [7] G.H. Hardy and S. Ramanujan, Asymptotic formulae in combinatory analysis, *Proc. London Math. Soc.*, 17 (1918), 75–175.
- [8] D.H. Lehmer, On the series for the partition function, *Trans. Amer. Math. Soc.*, 43 (1938), 271–292.
- [9] D.H. Lehmer, On the remainders and convergence of the series for the partition function, *Trans. Amer. Math. Soc.*, 46 (1939), 362–373.
- [10] H. Rademacher, A convergent series for the partition function $p(n)$, *Proc. Nat. Acad. Sci. USA*, 23 (1937), 78–84.
- [11] Z.W. Sun, On a sequence involving sums of primes, *Bull. Aust. Math. Soc.* 88 (2013), 197–205.
- [12] Y. Wang and B.X. Zhu, Proofs of some conjectures on monotonicity of number-theoretic and combinatorial sequences, *Sci. China Math.* 57 (2014), 2429–2435.