

Turán Problems for Vertex-disjoint Cliques in Multi-partite Hypergraphs

Erica L.L. Liu¹, Jian Wang²

¹Center for Applied Mathematics
Tianjin University
Tianjin 300072, P. R. China

²Department of Mathematics
Taiyuan University of Technology
Taiyuan 030024, P. R. China

E-mail: ¹liulingling@tju.edu.cn, ²wangjian01@tyut.edu.cn

Abstract

For two s -uniform hypergraphs H and F , the Turán number $ex_s(H, F)$ is the maximum number of edges in an F -free subgraph of H . Let s, r, k, n_1, \dots, n_r be integers satisfying $2 \leq s \leq r$ and $n_1 \leq n_2 \leq \dots \leq n_r$. De Silva, Heysse and Young determined $ex_2(K_{n_1, \dots, n_r}, kK_2)$ and De Silva, Heysse, Kapilow, Schenfisch and Young determined $ex_2(K_{n_1, \dots, n_r}, kK_r)$. In this paper, as a generalization of these results, we consider three Turán-type problems for k disjoint cliques in r -partite s -uniform hypergraphs. First, we consider a multi-partite version of the Erdős matching conjecture and determine $ex_s(K_{n_1, \dots, n_r}^{(s)}, kK_s^{(s)})$ for $n_1 \geq s^3k^2 + sr$. Then, using a probabilistic argument, we determine $ex_s(K_{n_1, \dots, n_r}^{(s)}, kK_r^{(s)})$ for all $n_1 \geq k$. Recently, Alon and Shikhelman determined asymptotically, for all F , the generalized Turán number $ex_2(K_n, K_s, F)$, which is the maximum number of copies of K_s in an F -free graph on n vertices. Here we determine $ex_2(K_{n_1, \dots, n_r}, K_s, kK_r)$ with $n_1 \geq k$ and $n_3 = \dots = n_r$. Utilizing a result on rainbow matchings due to Glebov, Sudakov and Szabó, we determine $ex_2(K_{n_1, \dots, n_r}, K_s, kK_r)$ for all n_1, \dots, n_r with $n_4 \geq r^r(k-1)k^{2r-2}$.

Keywords: Turán number; multi-partite hypergraphs; probabilistic argument.

Mathematics Subject Classification (2010): 05C35, 05C65

1 Introduction

An s -uniform hypergraph, or simply an s -graph, is a hypergraph whose edges have exactly s vertices. For an s -graph H , let $V(H)$ be the vertex set of H and $E(H)$ the edge set of H . An s -graph H is called F -free if H does not contain any copy of F as a subgraph. For two s -graphs H and F , the Turán number $ex_s(H, F)$ is the maximum number of edges of an F -free subgraph of H . Denote by $K_t^{(s)}$ the complete s -graph on t vertices. A copy of $K_t^{(s)}$ in an s -graph H is also called a t -clique of H . Let $kK_t^{(s)}$ denote the s -graph consisting of k vertex-disjoint copies of $K_t^{(s)}$. If $t = s$, then $kK_s^{(s)}$ represents a matching of size k . Let n_1, \dots, n_r be integers and V_1, V_2, \dots, V_r be disjoint vertex sets with $|V_i| = n_i$ for each $i = 1, \dots, r$. A complete r -partite s -graph on vertex classes V_1, V_2, \dots, V_r , denoted by

$K^{(s)}(V_1, V_2, \dots, V_r)$ or $K_{n_1, n_2, \dots, n_r}^{(s)}$, is defined to be the s -graph whose edge set consists of all the s -element subsets S of $V_1 \cup V_2 \cup \dots \cup V_r$ such that $|S \cap V_i| \leq 1$ for all $i = 1, \dots, r$. An s -graph H is called an r -partite s -graph on vertex classes V_1, V_2, \dots, V_r if H is a subgraph of $K^{(s)}(V_1, V_2, \dots, V_r)$. For $s = 2$, we often write $K_t, kK_t, K(V_1, V_2, \dots, V_r), K_{n_1, n_2, \dots, n_r}$ and $ex(H, F)$ instead of $K_t^{(2)}, kK_t^{(2)}, K^{(2)}(V_1, V_2, \dots, V_r), K_{n_1, n_2, \dots, n_r}^{(2)}$ and $ex_2(H, F)$. Let $[n]$ denote the set $\{1, 2, \dots, n\}$ and $[m, n]$ denote the set $\{m, m+1, \dots, n\}$ for $m \leq n$.

Turán-type problems were first considered by Mantel [17] in 1907, who determined $ex(K_n, K_3)$. In 1941, Turán [19] showed that the balanced complete t -partite graph on n vertices, called the Turán graph and denoted by $T_{n,t}$, is the unique graph that maximises the number of edges among all K_{t+1} -free graphs on n vertices. Since then, Turán numbers of graphs and hypergraphs have been extensively studied. However, even though lots of progress has been made, most of the Turán problems for bipartite graphs and for hypergraphs are still open. Specifically, none of the Turán numbers $ex_s(K_n^{(s)}, K_t^{(s)})$ with $t > s > 2$ has yet been determined, even asymptotically. We recommend the reader to consult [15, 18] for surveys on Turán numbers of graphs and hypergraphs.

Many problems in additive combinatorics are closely related to Turán-type problems in multi-partite graphs and hypergraphs. Recently, Turán problems in multi-partite graphs have received a lot of attention, see [3, 6, 14]. The following result, which is attributed to De Silva, Heysse and Young, determines $ex(K_{n_1, \dots, n_r}, kK_2)$.

Theorem 1.1. *For $n_1 \leq n_2 \leq \dots \leq n_r$ and $k \leq n_1$,*

$$ex(K_{n_1, n_2, \dots, n_r}, kK_2) = (k-1)(n_2 + \dots + n_r).$$

Since it seems that their preprint has not been published online, we present a proof of Theorem 1.1 in the Appendix for the completeness of the paper. In [6], De Silva, Heysse, Kapilow, Schenfisch and Young determined $ex(K_{n_1, \dots, n_r}, kK_r)$.

Theorem 1.2. [6] *For $n_1 \leq n_2 \leq \dots \leq n_r$ and $k \leq n_1$,*

$$ex(K_{n_1, \dots, n_r}, kK_r) = \sum_{1 \leq i < j \leq r} n_i n_j - n_1 n_2 + (k-1)n_2.$$

In this paper, we consider three Turán-type problems for k disjoint cliques in r -partite s -graphs. Let n_1, n_2, \dots, n_r be integers. For any $A \subset [r]$, denote $\prod_{i \in A} n_i$ by n_A . Define

$$f_k^{(s)}(n_2, \dots, n_r) = (k-1) \sum_{\substack{A: A \subset [2, r] \\ |A|=s-1}} n_A,$$

$$g_k^{(s)}(n_1, n_2, \dots, n_r) = \sum_{\substack{A: A \subset [r] \\ |A|=s}} n_A - n_{[s]} + (k-1)n_{[2, s]},$$

and

$$h_k^{(s)}(n_1, n_2, \dots, n_r) = \sum_{\substack{A: A \subset [r] \\ |A|=s, \{1, 2\} \not\subset A}} n_A + \sum_{\substack{A: A \subset [3, r] \\ |A|=s-2}} (k-1)n_2 n_A.$$

Theorem 1.3. *For $2 \leq s \leq r$, $k \geq 1$ and $n_1 \leq n_2 \leq \dots \leq n_r$, if $n_1 \geq s^3 k + sr$ for $s \leq r-2$; $n_1 \geq s^3 k^2 + sr$ for $s = r-1$ and $n_1 \geq k$ for $s = r$, then*

$$ex_s(K_{n_1, n_2, \dots, n_r}^{(s)}, kK_s^{(s)}) = f_k^{(s)}(n_2, \dots, n_r).$$

It should be mentioned that the problem in Theorem 1.3 can be viewed as a multi-partite version of the Erdős matching conjecture, which states that

$$ex_s(K_n^{(s)}, kK_s^{(s)}) = \max \left\{ \binom{ks-1}{s}, \binom{n}{s} - \binom{n-k+1}{s} \right\}$$

and is still open when n is close to $s(k-1)$, see [5, 7, 10, 11, 12] for recent progress. The lower bound in Theorem 1.3 follows from the following construction. Let H_1 be an r -partite s -graph on vertex classes V_1, V_2, \dots, V_r with sizes n_1, n_2, \dots, n_r , respectively. Let V'_1 be a $(k-1)$ -element subset of V_1 . An edge S of $K^{(s)}(V_1, V_2, \dots, V_r)$ forms an edge of H_1 if and only if $S \cap V'_1 \neq \emptyset$. It is easy to see that H_1 is $kK_s^{(s)}$ -free. Otherwise, if H_1 has a matching of size k , then we have $|V'_1| \geq k$ since each edge of H_1 contains a vertex in V'_1 .

As our second main result, we use a probabilistic argument to determine $ex_s(K_{n_1, \dots, n_r}^{(s)}, kK_r^{(s)})$.

Theorem 1.4. *For $2 \leq s \leq r$, $n_1 \leq n_2 \leq \dots \leq n_r$ and $k \leq n_1$,*

$$ex_s(K_{n_1, \dots, n_r}^{(s)}, kK_r^{(s)}) = g_k^{(s)}(n_1, n_2, \dots, n_r).$$

The lower bound in Theorem 1.4 follows from the following construction. Let H_2 be an r -partite s -graph on vertex classes V_1, V_2, \dots, V_r with sizes n_1, n_2, \dots, n_r , respectively. Let V'_1 be an $(n_1 - k + 1)$ -element subset of V_1 and let H_2 be obtained by deleting all the edges of $K^{(s)}(V'_1, V_2, \dots, V_r)$ from $K^{(s)}(V_1, V_2, \dots, V_r)$. It is easy to see that H_2 is $kK_r^{(s)}$ -free. Otherwise, if there are k vertex-disjoint copies of $K_r^{(s)}$ in H_2 , then we have $|V_1 \setminus V'_1| \geq k$ since each copy of $K_r^{(s)}$ in H_2 contains a vertex in $V_1 \setminus V'_1$.

We also consider the generalized Turán problem in multi-partite graphs. Let $ex(G, T, F)$ denote the maximum number of copies of T in an F -free subgraph of G . The first result of this type is due to Zykov [20], who showed that the Turán graph also maximises the number of s -cliques in an n -vertex K_{t+1} -free graph for $s \leq t$. Recently, Alon and Shikhelman [2] determined $ex(K_n, K_s, F)$ asymptotically for any F with chromatic number $\chi(F) = t+1 > s$. Precisely, they proved that

$$ex(K_n, K_s, F) = k_s(T_{n,t}) + o(n^s),$$

where $k_s(T_{n,t})$ denotes the number of s -cliques in the Turán graph $T_{n,t}$. Later, the error term of this result was further improved by Ma and Qiu [16].

In this paper, we also study the maximum number of s -cliques in a kK_r -free subgraph of K_{n_1, \dots, n_r} . By the same probabilistic argument as in the proof of Theorem 1.4, we obtain the following result.

Theorem 1.5. *For $2 \leq s \leq r$, $n_1 \leq n_2 \leq n_3$ and $k \leq n_1$,*

$$ex(K_{n_1, n_2, \underbrace{n_3, \dots, n_3}_{r-2}}, K_s, kK_r) = h_k^{(s)}(n_1, n_2, \underbrace{n_3, \dots, n_3}_{r-2}).$$

Note that for $r = 3, s \leq 3$ and arbitrary n_1, n_2, n_3 , the Turán number $ex(K_{n_1, n_2, n_3}, K_s, kK_3)$ is determined by Theorem 1.5. Utilizing a result on rainbow matchings due to Glebov, Sudakov and Szabó [13], we also determine $ex(K_{n_1, \dots, n_r}, K_s, kK_r)$ for $r \geq 4$ and n_4 sufficiently larger than k .

Theorem 1.6. *For $r \geq 4, 2 \leq s \leq r, n_1 \leq n_2 \leq \dots \leq n_r$ and $k \leq n_1$, if $n_4 \geq r^r(k-1)k^{2r-2}$, then*

$$ex(K_{n_1, \dots, n_r}, K_s, kK_r) = h_k^{(s)}(n_1, n_2, \dots, n_r).$$

The lower bounds in Theorems 1.5 and 1.6 follow from the same construction as follows. Let G be an r -partite graph on V_1, V_2, \dots, V_r , which are of sizes n_1, n_2, \dots, n_r , respectively. Let V'_1 be an $(n_1 - k + 1)$ -element subset of V_1 . Then G is obtained by deleting all the edges of $K(V'_1, V_2)$ from $K(V_1, V_2, \dots, V_r)$. It is easy to see that G is kK_r -free. Otherwise, if there are k vertex-disjoint copies of K_r in G , then we have $|V_1 \setminus V'_1| \geq k$ since each copy of K_r in G contains a vertex in $V_1 \setminus V'_1$.

The rest of the paper is organized as follows. We will prove Theorem 1.3 in Section 2. In Section 3, we prove Theorem 1.4. In Section 4, we prove Theorems 1.5 and 1.6.

2 Turán number of $kK_s^{(s)}$ in r -partite s -graphs

In this section, we prove Theorem 1.3. First, we consider the case $s = r$, which is the base case for other results in this paper. Aharoni and Howard [1] determined the maximum number of edges in a balanced r -partite r -graph that is $kK_r^{(r)}$ -free. By the same argument, we prove the following result:

Lemma 2.1. *For any integers $1 \leq k \leq n_1 \leq n_2 \leq \dots \leq n_r$,*

$$ex_r(K_{n_1, \dots, n_r}^{(r)}, kK_r^{(r)}) = (k-1)n_2 \cdots n_r.$$

Proof. We shall partition the edge set of $K^{(r)}(V_1, \dots, V_r)$ into $n_2 n_3 \cdots n_r$ matchings of size n_1 each. Let $V_i = \{v_{i,0}, v_{i,1}, \dots, v_{i,n_i-1}\}$ for $i = 1, 2, \dots, r$ and

$$\Lambda = [0, n_2 - 1] \times [0, n_3 - 1] \times \cdots \times [0, n_r - 1].$$

For any $(r-1)$ -tuple $(x_2, x_3, \dots, x_r) \in \Lambda$, define

$$E(x_2, x_3, \dots, x_r) = \left\{ \{v_{1,x}, v_{2,(x+x_2) \bmod n_2}, \dots, v_{r,(x+x_r) \bmod n_r}\} : x \in [0, n_1 - 1] \right\}.$$

It is easy to see that $E(x_2, x_3, \dots, x_r)$ is a matching of size n_1 . Moreover, let

$$\Omega = \{E(x_2, x_3, \dots, x_r) : (x_2, x_3, \dots, x_r) \in \Lambda\}.$$

We shall show that Ω forms a partition of the edge set of $K^{(r)}(V_1, \dots, V_r)$. On one hand, let $e = \{v_{1,x_1}, v_{2,x_2}, \dots, v_{r,x_r}\}$ be an edge in $K^{(r)}(V_1, \dots, V_r)$ with $v_{i,x_i} \in V_i$ for each $i = 1, 2, \dots, r$. Define

$$y_i := (x_i - x_1) \bmod n_i$$

for each $i = 2, \dots, r$. It is easy to check that $e \in E(y_2, y_3, \dots, y_r)$. Moreover, for each $(x_2, x_3, \dots, x_r) \in \Lambda$, $E(x_2, x_3, \dots, x_r) \subset E(K^{(r)}(V_1, \dots, V_r))$ holds. Thus, we have

$$E(K^{(r)}(V_1, \dots, V_r)) = \bigcup_{(x_2, x_3, \dots, x_r) \in \Lambda} E(x_2, x_3, \dots, x_r).$$

On the other hand, for any two different tuples $(y_2, y_3, \dots, y_r), (z_2, z_3, \dots, z_r) \in \Lambda$, we claim that $E(y_2, y_3, \dots, y_r) \cap E(z_2, z_3, \dots, z_r) = \emptyset$. Otherwise if there exists $\{v_{1,x_1}, v_{2,x_2}, \dots, v_{r,x_r}\} \in E(y_2, y_3, \dots, y_r) \cap E(z_2, z_3, \dots, z_r)$, then we have

$$x_i \equiv (x_1 + y_i) \bmod n_i \equiv (x_1 + z_i) \bmod n_i$$

for all $i = 2, \dots, r$. It follows that $y_i \equiv z_i \bmod n_i$. Since $y_i, z_i \in \{0, 1, \dots, n_i - 1\}$, we obtain $y_i = z_i$ for all $i = 2, \dots, r$, a contradiction. Therefore, Ω forms a partition of the edge set of $K^{(r)}(V_1, \dots, V_r)$.

Assume that $H \subseteq K^{(r)}(V_1, \dots, V_r)$ and $e(H) \geq (k-1)n_2 \cdots n_r + 1$. Then the partition

$$\{E(H) \cap E(x_2, x_3, \dots, x_r) : (x_2, x_3, \dots, x_r) \in \Lambda\}$$

of $E(H)$ shows that at least one of the matchings $E(H) \cap E(x_2, x_3, \dots, x_r)$ has size k or more, a contradiction.

For the lower bound, $K_{k-1, n_2, \dots, n_r}^{(r)}$ is a $kK_r^{(r)}$ -free r -graph with $(k-1)n_2 \cdots n_r$ edges. Thus, we conclude that $ex_r(K_{n_1, \dots, n_r}^{(r)}, kK_r^{(r)}) = (k-1)n_2 \cdots n_r$. \square

Let H be an s -graph. For $u, v \in V(H)$ and $e \in E(H)$, we define a *shifting* operator S_{uv} on e as follows:

$$S_{uv}(e) = \begin{cases} (e \setminus \{v\}) \cup \{u\}, & \text{if } v \in e, u \notin e \text{ and } (e \setminus \{v\}) \cup \{u\} \notin E(H), \\ e, & \text{otherwise.} \end{cases}$$

Define $S_{uv}(H)$ be the s -graph with vertex set $V(H)$ and edge set $\{S_{uv}(e) : e \in E(H)\}$.

It is easy to see that $e(S_{uv}(H)) = e(H)$. Let $\nu(H)$ denote the size of a largest matching in H . Frankl [9] showed that applying the shifting operator to H does not increase $\nu(H)$. For the completeness we also include a short proof of this.

Lemma 2.2. [9] *Let H be an s -graph. For any $u, v \in V(H)$,*

$$\nu(S_{uv}(H)) \leq \nu(H).$$

Proof. Suppose for contradiction that $\nu(H) = k$ but $\nu(S_{uv}(H)) = k+1$. Let $M = \{e_1, e_2, \dots, e_{k+1}\}$ be a matching of size $k+1$ in $S_{uv}(H)$. Since each edge in $E(S_{uv}(H)) \setminus E(H)$ contains u , it follows that exactly one of e_1, e_2, \dots, e_{k+1} is not in H . Without loss of generality, we assume that $e_{k+1} \notin E(H)$. Then, $u \in e_{k+1}$, $v \notin e_{k+1}$ and $e'_{k+1} = e_{k+1} \setminus \{u\} \cup \{v\} \in E(H)$. Since $\nu(H) = k$, it is easy to see that $e'_{k+1} \cap e_i = \{v\}$ for some $i \in [k]$. Since $e_i \in E(H) \cap E(S_{uv}(H))$ and $u \notin e_i$, by the definition of S_{uv} we have $e'_i = e_i \setminus \{v\} \cup \{u\} \in E(H)$. Then, $M \setminus \{e_i, e_{k+1}\} \cup \{e'_i, e'_{k+1}\}$ forms a matching of size $k+1$ in H , a contradiction. \square

Let H be an r -partite s -graph on vertex classes V_1, V_2, \dots, V_r , and

$$V_i = \{a_{i,1}, a_{i,2}, \dots, a_{i,n_i}\}$$

for $i = 1, 2, \dots, r$. Define a partial order \prec on $V = \cup_{i=1}^r V_i$ such that

$$a_{i,1} \prec a_{i,2} \prec \dots \prec a_{i,n_i}$$

for each i and vertices from different parts are incomparable. For two different edges $S_1 = \{a_1, a_2, \dots, a_s\}$ and $S_2 = \{b_1, b_2, \dots, b_s\}$ in $K^{(r)}(V_1, \dots, V_r)$, we define $S_1 \prec S_2$ if and only if there exists a permutation $\sigma_1 \sigma_2 \cdots \sigma_s$ of $[s]$ such that $a_j \prec b_{\sigma_j}$ or $a_j = b_{\sigma_j}$ holds for all $j = 1, \dots, s$.

An r -partite s -graph H is called a *stable r -partite s -graph* if $S_{ab}(H) = H$ holds for all $a, b \in V(H)$ with $a \prec b$. If H is stable and $e \in E(H)$, it is easy to see that for any s -element vertex subset S with $S \prec e$, we have $S \in E(H)$. Indeed, let $S = \{a_1, a_2, \dots, a_s\}$ and $e = \{b_1, b_2, \dots, b_s\}$. Without loss of generality, we may assume that $a_i \prec b_i$ for each $i = 1, \dots, s_0$ and $a_i = b_i$ for each $i = s_0 + 1, \dots, s$. Since $S_{a_1 b_1}(H) = H$ and $e \in E(H)$, it is easy to see that $e_1 = e \setminus \{b_1\} \cup \{a_1\} \in E(H)$. Since $S_{a_2 b_2}(H) = H$ and $e_1 \in E(H)$,

it follows that $e_2 = e_1 \setminus \{b_2\} \cup \{a_2\} \in E(H)$. Repeat the same argument for $i = 3, \dots, s_0$ and we shall obtain that $S \in E(H)$.

To obtain a stable r -partite s -graph, we can apply the shifting operator to H iteratively. For an intermediate step, let H^* be the current r -partite s -graph. If H^* is stable, we are done. If H^* is not stable, there exists a pair (a, b) such that $a \prec b$ and $S_{ab}(H^*) \neq H^*$. Then, apply S_{ab} to H^* and we obtain a new r -partite s -graph. Define

$$g(H^*) := \sum_{e \in E(H^*)} \sum_{i=1}^r \sum_{j: a_{i,j} \in e} j.$$

Since after each step $g(H^*)$ decreases strictly and $g(H) > 0$ holds for all the non-empty r -partite s -graphs H , the process will end in finite steps. It should be mentioned that if we apply the shifting operator in different orders, at the end we may arrive at different stable r -partite s -graphs. For more properties of the shifting operator, we refer the reader to [10].

For $u, v \in V(H)$, let $L_H(u)$ denote the set of edges in H containing u and $L_H(u, v)$ denote the set of edges in H containing u and v . Let $d_H(u)$ and $d_H(u, v)$ denote the cardinality of $L_H(u)$ and $L_H(u, v)$, respectively. For $X \subset V(H)$, let $\Gamma_H(X)$ denote the set of edges in H that intersect X . It should be noticed that $\Gamma_H(\{u\})$ is the same as $L_H(u)$. The subscripts will be dropped if there is no confusion. For $S \subset V(H)$, let $H[S]$ denote the s -graph induced by S and $H \setminus S$ the s -graph induced by $V(H) \setminus S$.

Lemma 2.3. *For $3 \leq s \leq r-1$, if $n \geq s^3k + sr$ for $s \leq r-2$ and $n \geq s^3k^2 + sr$ for $s = r-1$, then*

$$ex_s(K_{\underbrace{n, \dots, n}_r}, kK_s^{(s)}) = (k-1) \binom{r-1}{s-1} n^{s-1}.$$

Proof. We prove the lemma by induction on k . For $k = 1$, the lemma holds trivially. Suppose that the lemma holds for all $k' < k$ and H is a $kK_s^{(s)}$ -free subgraph of $K_{\underbrace{n, \dots, n}_r}^{(s)}$

with the maximum number of edges. By Lemma 2.2, we may further assume that H is stable. Let $T_0 = \{a_{1,1}, a_{2,1}, \dots, a_{r,1}\}$, $\nu(H \setminus T_0) = t$ and $M' = \{e_1, \dots, e_t\}$ be a largest matching in $H \setminus T_0$. Since H is stable, $H[T_0]$ is not empty. Then, it is easy to see that $t \leq k-2$. Otherwise, for any edge $e \in H[T_0]$, $\{e\} \cup M'$ forms a matching of size k in H . Since $H \setminus T_0$ is $(t+1)K_s^{(s)}$ -free and $n-1 \geq s^3(t+1) + rs$ for $s \leq r-2$ and $n-1 \geq s^3(t+1)^2 + rs$ for $s = r-1$, by the induction hypothesis, it follows that

$$e(H \setminus T_0) \leq t \binom{r-1}{s-1} (n-1)^{s-1}.$$

If

$$|\Gamma(T_0)| \leq (k-1) \binom{r-1}{s-1} n^{s-1} - t \binom{r-1}{s-1} (n-1)^{s-1},$$

then we conclude that

$$e(H) = e(H \setminus T_0) + |\Gamma(T_0)| \leq (k-1) \binom{r-1}{s-1} n^{s-1}.$$

Thus, we are left with the case

$$|\Gamma(T_0)| > (k-1) \binom{r-1}{s-1} n^{s-1} - t \binom{r-1}{s-1} (n-1)^{s-1}. \quad (2.1)$$

We will show that inequality (2.1) either implies the lemma or leads to a contradiction. The proof splits into two cases according to the value of t .

Case 1. $t = k - 2$. Without loss of generality, assume that $a_{1,1}$ is the vertex in T_0 with the maximum degree within H . Since

$$\sum_{i=1}^r d(a_{i,1}) \geq |\Gamma(T_0)|,$$

by the inequality (2.1) it follows that

$$\begin{aligned} d(a_{1,1}) &\geq \frac{1}{r} |\Gamma(T_0)| \\ &> \frac{1}{r} \binom{r-1}{s-1} ((k-1)n^{s-1} - (k-2)(n-1)^{s-1}) \\ &\geq \frac{1}{r} \binom{r-1}{s-1} n^{s-1}. \end{aligned}$$

Then, the structure of H can be partly described by the following claim.

Claim 1. Every edge in H intersects V_1 .

Proof. Suppose to the contrary that there exists an edge in H that does not intersect V_1 . Since H is stable, there exists an edge in T_0 that does not contain $a_{1,1}$. Let e_0 be such an edge. Let S be the set of vertices covered by the edges in $M' \cup \{e_0\}$, where, as before, M' is a matching of size $k - 2$ in $H \setminus T_0$. Clearly, $|S| = (k - 1)s$. For each $u \in S$, the number of edges containing u and $a_{1,1}$ is at most $\binom{r-2}{s-2} n^{s-2}$. Then, there are at most $(k - 1)s \binom{r-2}{s-2} n^{s-2}$ edges in $L(a_{1,1})$ that intersect S . It follows that the number of edges in $L(a_{1,1})$ that are disjoint from the edges in $M' \cup \{e_0\}$ is at least

$$\begin{aligned} &d(a_{1,1}) - (k - 1)s \binom{r-2}{s-2} n^{s-2} \\ &> \frac{1}{r} \binom{r-1}{s-1} n^{s-1} - (k - 1)s \binom{r-2}{s-2} n^{s-2} \\ &= \binom{r-2}{s-2} n^{s-2} \left(\frac{r-1}{r(s-1)} n - (k - 1)s \right) \\ &> 0, \end{aligned}$$

where the last inequality follows from the assumption that $n \geq 2s^2k$. Thus, let e'_0 be an edge in $L(a_{1,1})$ that is disjoint from the edges in $M' \cup \{e_0\}$. Then $M' \cup \{e_0, e'_0\}$ forms a matching of size k in H , which contradicts the fact that H is $kK_s^{(s)}$ -free. Therefore, the claim holds. \square

Define an r -partite r -graph H^* on vertex classes V_1, \dots, V_r . An r -element subset T of $V(H)$ forms an edge of H^* if $H[T]$ is non-empty and $|T \cap V_i| = 1$ for $i = 1, \dots, r$. Since H is $kK_s^{(s)}$ -free, it follows that H^* is $kK_r^{(r)}$ -free. By Lemma 2.1, we have $e(H^*) \leq (k - 1)n^{r-1}$. Now we prove the result by double counting. Let

$$\Phi = \{(e, T) : e \in E(H), T \in E(K^{(r)}(V_1, V_2, \dots, V_r)), \text{ and } e \subset T\}.$$

For every $T = \{x_1, x_2, \dots, x_r\} \in E(H^*)$ with $x_i \in V_i$ for each i , since by Claim 1 each edge in $H[T]$ contains x_1 , it follows that

$$e(H[T]) \leq \binom{r-1}{s-1}.$$

Moreover, $H[T]$ is non-empty if and only if T forms an edge in H^* . Thus,

$$|\Phi| \leq (k-1)n^{r-1} \binom{r-1}{s-1}.$$

On the other hand, each edge in H appears in n^{r-s} pairs in Φ . Therefore, we have

$$e(H) = |\Phi|/n^{r-s} \leq (k-1) \binom{r-1}{s-1} n^{s-1}.$$

Case 2. $t \leq k-3$. Let X be the set of vertices in T_0 with degree greater than $\frac{1}{2r} \binom{r-1}{s-1} n^{s-1}$ and $Y = T_0 \setminus X$.

First, we prove the following claim, which will be used several times.

Claim 2. $\nu(H \setminus X) \leq k-1-|X|$.

Proof. Suppose to the contrary that $\nu(H \setminus X) \geq k-|X|$. Let M^* be a largest matching in $H \setminus X$. We shall show that M^* can be greedily enlarged to a matching of size k in H , which contradicts the fact that $\nu(H) \leq k-1$. Since $\nu(H \setminus X) \geq k-|X|$, it follows that $|X| \geq k - \nu(H \setminus X) = k - |M^*|$. Let $l = k - |M^*|$ and x_1, x_2, \dots, x_l be l vertices in X . Set $X_i^+ = \{x_{i+1}, x_{i+2}, \dots, x_l\}$ and $M_0 = M^*$. Note that

$$\begin{aligned} d(x_1) &\geq \frac{1}{2r} \binom{r-1}{s-1} n^{s-1} \\ &= \frac{n}{2(s-1)} \cdot \frac{r-1}{r} \cdot \binom{r-2}{s-2} n^{s-2} \\ &\geq \frac{n}{2(s-1)} \cdot \frac{2}{3} \cdot \binom{r-2}{s-2} n^{s-2} \\ &> sk \cdot \binom{r-2}{s-2} n^{s-2} \\ &> (|M_0|s + |X_1^+|) \binom{r-2}{s-2} n^{s-2}, \end{aligned}$$

where the second inequality follows from the fact that $r \geq 3$, the third inequality follows from the assumption that $n \geq 3s^2k$ and the last inequality follows from the fact that $k = |M^*| + l > |M_0| + |X_1^+|$. Since there are at most $(|M_0|s + |X_1^+|) \binom{r-2}{s-2} n^{s-2}$ edges in $L(x_1)$ that intersect $(\cup_{e \in M_0} e) \cup X_1^+$, we can choose e'_1 from $L(x_1)$ such that $M_1 = M_0 \cup \{e'_1\}$ is a matching of size $|M_0| + 1$ and x_2, x_3, \dots, x_l are not used. Now we continue to choose an edge from each of $L(x_2), \dots, L(x_l)$ to enlarge the matching. When dealing with $L(x_i)$, note that $|X_i^+| = l - i$. Since there are at most $(|M_{i-1}|s + |X_i^+|) \binom{r-2}{s-2} n^{s-2}$ edges in $L(x_i)$ that intersect $(\cup_{e \in M_{i-1}} e) \cup X_i^+$ and

$$\begin{aligned} d(x_i) &\geq \frac{1}{2r} \binom{r-1}{s-1} n^{s-1} \\ &> sk \binom{r-2}{s-2} n^{s-2} \\ &> (|M_{i-1}|s + |X_i^+|) \binom{r-2}{s-2} n^{s-2}, \end{aligned}$$

where the last inequality follows from $k = |M^*| + l > |M_{i-1}| + |X_i^+|$, therefore we can choose e'_i from $L(x_i)$ such that $M_i = M_{i-1} \cup \{e'_i\}$ is a matching of size $|M_{i-1}| + 1$ and

$x_{i+1}, x_{i+2}, \dots, x_l$ are not used. Finally, we end up with M_l , which is a matching of size $|M^*| + l = k$. It contradicts the fact that H is $kK_s^{(s)}$ -free. Thus, we conclude that $\nu(H \setminus X) \leq k - 1 - |X|$. \square

Then, we show that the sizes of both X and the matching number of $H \setminus X$ can be determined by the matching number of $H \setminus T_0$.

Claim 3. $|X| = k - 1 - t$.

Proof. By Claim 2 we have

$$t = \nu(H \setminus T_0) \leq \nu(H \setminus X) \leq k - 1 - |X|.$$

Thus, $|X| \leq k - 1 - t$. If $|X| \leq k - 2 - t$, then

$$\begin{aligned} |\Gamma(T_0)| &\leq |X| \binom{r-1}{s-1} n^{s-1} + (r - |X|) \cdot \frac{1}{2r} \binom{r-1}{s-1} n^{s-1} \\ &= |X| \binom{r-1}{s-1} n^{s-1} \left(1 - \frac{1}{2r}\right) + \frac{1}{2} \binom{r-1}{s-1} n^{s-1} \\ &\leq \left((k - 2 - t) \left(1 - \frac{1}{2r}\right) + \frac{1}{2}\right) \binom{r-1}{s-1} n^{s-1} \\ &< (k - 1 - t) \binom{r-1}{s-1} n^{s-1}, \end{aligned}$$

which contradicts the inequality (2.1). Thus, the claim holds. \square

Claim 4. $\nu(H \setminus X) = \nu(H \setminus T_0) = t$.

Proof. By Claims 2 and 3, we have

$$\nu(H \setminus X) \leq k - 1 - |X| = t = \nu(H \setminus T_0).$$

Moreover, since $H \setminus T_0$ is a subgraph of $H \setminus X$, it follows that $\nu(H \setminus X) \geq \nu(H \setminus T_0)$. Thus, the claim holds. \square

We also claim that Y cannot be an empty set. Otherwise, by Claim 3 we have

$$|\Gamma(T_0)| = |\Gamma(X)| \leq |X| \binom{r-1}{s-1} n^{s-1} = (k - 1 - t) \binom{r-1}{s-1} n^{s-1},$$

which contradicts the inequality (2.1).

By Claim 4, we have that all edges in $L_{H \setminus X}(y)$ intersect $\cup_{e \in M'} e$ for each $y \in Y$. Otherwise, if there exists an edge e_0 in $L_{H \setminus X}(y)$ that is disjoint from $\cup_{e \in M'} e$ for some $y \in Y$, then $M' \cup \{e_0\}$ forms a matching of size $t + 1$ in $H \setminus X$, a contradiction. Then, we can obtain an upper bound on $|\Gamma_{H \setminus X}(Y)|$ by the following argument. For $e_i \in M'$, define a bipartite graph G_i on vertex classes Y and e_i , where e_i is viewed as one of the sides of G_i . For $u \in e_i$ and $v \in Y$, $\{u, v\}$ is an edge of G_i if $d_{H \setminus X}(u, v) > (t + 1)s \binom{r-3}{s-3} n^{s-3}$. If there is an i such that $\nu(G_i) \geq 2$, let $\{u_p, v_p\}$ and $\{u_q, v_q\}$ be two disjoint edges of G_i with $u_p, u_q \in e_i$ and $v_p, v_q \in Y$. Since there are at most $ts \binom{r-3}{s-3} n^{s-3}$ edges in $L_{H \setminus X}(u_p, v_p)$ that intersect $(\cup_{e \in M'} e) \cup \{v_q\} \setminus \{u_p\}$, we can find an edge f_p in $L_{H \setminus X}(u_p, v_p)$ that is disjoint from $(\cup_{e \in M'} e) \cup \{v_q\} \setminus \{u_p\}$. Similarly, there are at most $(t + 1)s \binom{r-3}{s-3} n^{s-3}$ edges in $L_{H \setminus X}(u_q, v_q)$ that intersect $(\cup_{e \in M'} e) \cup \{f_p\} \setminus \{u_q\}$. Thus, we can find an edge f_q in $L_{H \setminus X}(u_q, v_q)$ that

is disjoint from $(\cup_{e \in M'} e) \cup \{f_p\} \setminus \{u_q\}$. Now $(M' \setminus \{e_i\}) \cup \{f_p, f_q\}$ forms a matching of size $t + 1$ in $H \setminus X$, which contradicts with Claim 4. Thus, we conclude that each G_i has matching number at most one.

Let $e_i \in M'$ and

$$\Gamma_{H \setminus X}(e_i, Y) = \{e \in E(H \setminus X) : e \cap e_i \neq \emptyset \text{ and } e \cap Y \neq \emptyset\}.$$

The rest of the proof is divided into two subcases according to the size of $|Y|$.

Case 2.1. $|Y| \geq s$. Since $\nu(G_i) \leq 1$, by Lemma 2.1, there are at most $|Y|$ edges in G_i . Then,

$$\begin{aligned} |\Gamma_{H \setminus X}(e_i, Y)| &\leq e(G_i) \binom{r-2}{s-2} n^{s-2} + (|Y||e_i| - e(G_i))(t+1)s \binom{r-3}{s-3} n^{s-3} \\ &= e(G_i) \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n - (t+1)s \right) + |Y||e_i|(t+1)s \binom{r-3}{s-3} n^{s-3} \\ &\leq |Y| \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n - (t+1)s \right) + |Y|s^2(t+1) \binom{r-3}{s-3} n^{s-3}, \\ &= (r - |X|) \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n + (t+1)s(s-1) \right), \end{aligned}$$

where the second inequality follows from the assumption that $n > sk$ and $e(G_i) \leq |Y|$.

By Claim 4, for any $y \in Y$ all edges in $L_{H \setminus X}(y)$ intersect $\cup_{e \in M'} e$. It follows that

$$|\Gamma_{H \setminus X}(Y)| \leq \sum_{i=1}^t |\Gamma_{H \setminus X}(e_i, Y)| \leq t(r - |X|) \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n + (t+1)s(s-1) \right).$$

Since

$$|\Gamma(X)| \leq \sum_{i=1}^{|X|} d(x_i) \leq |X| \binom{r-1}{s-1} n^{s-1},$$

therefore

$$\begin{aligned} |\Gamma(T_0)| &= |\Gamma(X)| + |\Gamma_{H \setminus X}(Y)| \\ &\leq |X| \binom{r-1}{s-1} n^{s-1} + t(r - |X|) \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n + (t+1)s(s-1) \right). \end{aligned} \quad (2.2)$$

By combining the inequalities (2.1) and (2.2) and using the fact that $|X| = k - 1 - t$, we arrive at

$$t \binom{r-1}{s-1} (n^{s-1} - (n-1)^{s-1}) \leq t(r - |X|) \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n + (t+1)s(s-1) \right).$$

Since $|X| \geq 2$ (because $|X| = k - 1 - t$ and in Case 2 we assume that $t \leq k - 3$), we have

$$\frac{(r-1)(r-2)}{(s-1)(s-2)} (n^{s-1} - (n-1)^{s-1}) \leq (r-2) n^{s-3} \left(\frac{r-2}{s-2} n + (t+1)s(s-1) \right). \quad (2.3)$$

By Taylor's Theorem with Lagrange remainder, it can be deduced that

$$n^{s-1} - (n-1)^{s-1} \geq (s-1)n^{s-2} - \frac{(s-1)(s-2)}{2} n^{s-3}. \quad (2.4)$$

By combining the inequalities (2.3) and (2.4), we obtain that

$$\frac{r-1}{s-2}n^{s-2} - \frac{r-1}{2}n^{s-3} \leq n^{s-3} \left(\frac{r-2}{s-2}n + (t+1)s(s-1) \right). \quad (2.5)$$

Since $t \leq k-3$, by simplifying the inequality (2.5) we arrive at

$$n \leq s(s-1)(s-2)(k-2) + \frac{(r-1)(s-2)}{2} < s^3k + sr,$$

which contradicts the fact that $n \geq s^3k + sr$.

Case 2.2. $|Y| \leq s-1$.

For each $i = 1, 2, \dots, t$, since $\nu(G_i) \leq 1$, by Lemma 2.1 we have $e(G_i) \leq s$. Then

$$\begin{aligned} |\Gamma_{H \setminus X}(e_i, Y)| &\leq e(G_i) \binom{r-2}{s-2} n^{s-2} + (|Y||e_i| - e(G_i))(t+1)s \binom{r-3}{s-3} n^{s-3} \\ &= e(G_i) \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2}n - (t+1)s \right) + |Y||e_i|(t+1)s \binom{r-3}{s-3} n^{s-3} \\ &\leq s \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2}n - (t+1)s \right) + |Y|s^2(t+1) \binom{r-3}{s-3} n^{s-3}, \\ &= s \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2}n + (t+1)s(|Y|-1) \right), \end{aligned}$$

where the second inequality follows from the assumption that $n > sk$ and $e(G_i) \leq s$. Thus,

$$|\Gamma_{H \setminus X}(Y)| \leq \sum_{i=1}^t |\Gamma_{H \setminus X}(e_i, Y)| \leq st \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2}n + (t+1)s(|Y|-1) \right). \quad (2.6)$$

Since

$$|\Gamma(X)| \leq \sum_{i=1}^{|X|} d(x_i) \leq |X| \binom{r-1}{s-1} n^{s-1},$$

therefore

$$\begin{aligned} |\Gamma(T_0)| &= |\Gamma(X)| + |\Gamma_{H \setminus X}(Y)| \\ &\leq |X| \binom{r-1}{s-1} n^{s-1} + st \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2}n + (t+1)s(|Y|-1) \right). \end{aligned} \quad (2.7)$$

By combining the inequalities (2.1) and (2.7) and using Claim 3, we arrive at

$$t \binom{r-1}{s-1} (n^{s-1} - (n-1)^{s-1}) \leq st \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2}n + (t+1)s(|Y|-1) \right). \quad (2.8)$$

Then by combining the inequalities (2.4) and (2.8) we obtain that

$$\frac{(r-1)(r-2)}{s-2}n^{s-2} - \frac{(r-1)(r-2)}{2}n^{s-3} \leq sn^{s-3} \left(\frac{r-2}{s-2}n + (t+1)s(|Y|-1) \right). \quad (2.9)$$

By simplifying, we arrive at

$$(r-1-s)n \leq \frac{(r-1)(s-2)}{2} + \frac{s^2(s-2)(t+1)(|Y|-1)}{r-2}.$$

Since $|Y| = r - |X| \leq r - 2$ and $t + 1 \leq k$, it follows that

$$(r - 1 - s)n \leq \frac{(r - 1)(s - 2)}{2} + s^2(s - 2)k.$$

Since $n \geq s^3k + sr$ when $s \leq r - 2$, it leads to a contradiction for $s \leq r - 2$.

For $r = s + 1$, we shall give a slightly better upper bound on $|L(X)|$ as follows. Let $X = \{x_1, \dots, x_{k-1-t}\}$, $X_0 = \emptyset$ and $X_i = \{x_1, \dots, x_i\}$ for $i = 1, 2, \dots, k - 1 - t$. Note that

$$|\Gamma(X)| = \sum_{i=1}^{|X|} |L_{H \setminus X_{i-1}}(x_i)|.$$

Now, it is easy to see that

$$L_{H \setminus X_1}(x_2) \leq \binom{s-1}{s-1} n^{s-1} + (n-1) \binom{s-1}{s-2} n^{s-2} = sn^{s-1} - (s-1)n^{s-2}.$$

For $i \neq 2$, we use the trivial inequality $|L_{H \setminus X_{i-1}}(x_i)| \leq sn^{s-1}$. Since $|X| \geq 2$ (because $|X| = k - 1 - t$ and in Case 2 we assume that $t \leq k - 3$), it follows that

$$|\Gamma(X)| = \sum_{i=1}^{|X|} |L_{H \setminus X_{i-1}}(x_i)| \leq |X|sn^{s-1} - (s-1)n^{s-2}.$$

Then, by the inequality (2.6), we obtain an upper bound on $\Gamma(T_0)$ as follows:

$$\begin{aligned} |\Gamma(T_0)| &= |\Gamma(X)| + |\Gamma_{H \setminus X}(Y)| \\ &\leq |X|sn^{s-1} - (s-1)n^{s-2} + st \binom{r-3}{s-3} n^{s-3} \left(\frac{r-2}{s-2} n + (t+1)s(|Y|-1) \right) \\ &= |X|sn^{s-1} - (s-1)n^{s-2} + st(s-2)n^{s-3} \left(\frac{s-1}{s-2} n + (t+1)s(|Y|-1) \right). \end{aligned} \quad (2.10)$$

By combining the inequalities (2.1) and (2.10) and using Claim 3, we have

$$ts(n^{s-1} - (n-1)^{s-1}) \leq st(s-2)n^{s-3} \left(\frac{s-1}{s-2} n + (t+1)s(|Y|-1) \right) - (s-1)n^{s-2}. \quad (2.11)$$

By simplifying, we obtain that

$$ts(n^{s-1} - (n-1)^{s-1}) \leq (st(s-1) - s + 1)n^{s-2} + s^2(s-2)t(t+1)(|Y|-1)n^{s-3}. \quad (2.12)$$

By combining the inequalities (2.4) and (2.12), we arrive at

$$(s-1)n \leq \frac{s(s-1)(s-2)t}{2} + s^2(s-2)t(t+1)(|Y|-1).$$

Since $|Y| = s + 1 - |X| \leq s - 1$ and $t + 1 \leq k$, it follows that

$$n \leq \frac{s(s-2)}{2}t + s^2t(t+1)(|Y|-1) \cdot \frac{s-2}{s-1} \leq \frac{s^2}{2}k + s^2(s-2)k^2 \leq s^3k^2,$$

which contradicts the fact that $n \geq s^3k^2 + sr$ for $s = r - 1$.

Thus, we complete the proof of Lemma 2.3. \square

In the following proof of Theorem 1.3, we shall use Theorem 1.1 and Lemma 2.3 as base cases.

Proof of Theorem 1.3. Notice that Lemma 2.1 implies the theorem for $s = r$. So we are left with the case $s \leq r - 1$. We prove by induction on $(s, \sum_{i=2}^r (n_i - n_1))$. The base case of $s = 2$ is verified for all r and $n_1 \leq n_2 \leq \dots \leq n_r$ by Theorem 1.1. For every $s \geq 3$, the base case of $\sum_{i=2}^r (n_i - n_1) = 0$ is verified for all r by Lemma 2.3. Suppose now that $\sum_{i=2}^r (n_i - n_1) > 0$. Assume that for all r , the theorem holds for all pairs $(s', \sum_{i=2}^r (n'_i - n'_1))$ such that $s' < s$ or $s' = s$ together with $\sum_{i=2}^r (n'_i - n'_1) < \sum_{i=2}^r (n_i - n_1)$. There exists an $i \in [2, r]$ such that $n_i > n_{i-1}$. Without loss of generality, assume that $i = r$. Let H be a $kK_s^{(s)}$ -free subgraph of $K_{n_1, \dots, n_r}^{(s)}$. By Lemma 2.2 we may assume that H is stable. Let V_r be the vertex set with cardinality n_r and

$$V_r = \{a_{r,1}, a_{r,2}, \dots, a_{r,n_r}\}.$$

Let $H' = H \setminus \{a_{r,n_r}\}$ and

$$H(a_{r,n_r}) = \{S \subset V : S \cup \{a_{r,n_r}\} \in E(H)\}.$$

Clearly, $H(a_{r,n_r})$ is an $(r-1)$ -partite $(s-1)$ -graph with parts of sizes n_1, n_2, \dots, n_{r-1} . We claim that $\nu(H(a_{r,n_r})) \leq k-1$. Otherwise, suppose $M = \{e_1, e_2, \dots, e_k\}$ is a matching of size k in $H(a_{r,n_r})$. Since H is stable and $n_r > k$, $\{e_1 \cup \{a_{r,1}\}, e_2 \cup \{a_{r,2}\}, \dots, e_k \cup \{a_{r,k}\}\}$ forms a matching of size k , which contradicts the fact that H is $kK_s^{(s)}$ -free. Since H' is $kK_s^{(s)}$ -free, by the induction hypothesis on $\sum_{i=2}^r (n_i - n_1)$, we have

$$e(H') \leq f_k^{(s)}(n_2, \dots, n_{r-1}, n_r - 1).$$

Since $H(a_{r,n_r})$ is a $kK_{s-1}^{(s-1)}$ -free $(r-1)$ -partite $(s-1)$ -graph, $n_1 \geq s^3k + sr \geq (s-1)^3k + (s-1)(r-1)$ for $(s-1) \leq (r-1) - 2$ and $n_1 \geq s^3k^2 + sr \geq (s-1)^3k^2 + (s-1)(r-1)$ for $(s-1) = (r-1) - 1$, by the induction hypothesis on s , we have

$$e(H(a_{r,n_r})) \leq f_k^{(s-1)}(n_2, \dots, n_{r-1}).$$

Thus,

$$\begin{aligned} e(H) &= e(H') + e(H(a_{r,n_r})) \\ &\leq f_k^{(s)}(n_2, \dots, n_{r-1}, n_r - 1) + f_k^{(s-1)}(n_2, \dots, n_{r-1}) \\ &= f_k^{(s)}(n_2, \dots, n_{r-1}, n_r), \end{aligned}$$

which completes the proof. \square

3 Turán number of $kK_r^{(s)}$ in r -partite s -graphs

In this section, we generalize the result of [6] to s -graphs by using a probabilistic argument. The following lemma will be useful for us.

Lemma 3.1. Assume that $b > 0$, $w_1 \geq w_2 \geq \dots \geq w_N > 0$ and let (P) be a linear programming model as follows:

$$\begin{aligned} \max \quad & z = \sum_{i=1}^N x_i \\ \text{s.t.} \quad & \sum_{i=1}^N w_i^{-1} x_i \leq b, \\ & 0 \leq x_i \leq w_i, \quad i = 1, 2, \dots, N. \end{aligned}$$

Let M be the integral part of b and $a = w_{M+1}(b - M)$. Then $\sum_{i=1}^M w_i + a$ is the optimal value of (P) .

Proof. Suppose to the contrary that there exists a feasible solution $y = (y_1, y_2, \dots, y_N)$ to (P) such that

$$\sum_{i=1}^N y_i > \sum_{i=1}^M w_i + a.$$

Since y is a feasible solution, it follows that

$$\sum_{i=1}^N w_i^{-1} y_i \leq b = M + w_{M+1}^{-1} a = \sum_{i=1}^M w_i^{-1} w_i + w_{M+1}^{-1} a.$$

Then, since $w_i \geq w_j$ for any $i < j$, we have

$$\sum_{i=1}^M w_i^{-1} (w_i - y_i) \geq \sum_{i=M+1}^N w_i^{-1} y_i - w_{M+1}^{-1} a \geq w_{M+1}^{-1} \left(\sum_{i=M+1}^N y_i - a \right) > w_{M+1}^{-1} \sum_{i=1}^M (w_i - y_i).$$

On the other hand, since

$$\sum_{i=1}^M w_i^{-1} (w_i - y_i) \leq w_M^{-1} \sum_{i=1}^M (w_i - y_i),$$

we arrived at $w_M^{-1} > w_{M+1}^{-1}$, a contradiction. Thus, the lemma holds. \square

Let H be an r -partite s -graph on vertex classes V_1, V_2, \dots, V_r . For any $A \subset [r]$, we shall write $\cup_{i \in A} V_i$ as V_A for short. Denote by $E(V_A)$ the edge set of the induced subgraph $H[V_A]$ and $e(V_A)$ the cardinality of $E(V_A)$.

Proof of Theorem 1.4. Suppose $H \subseteq K_{n_1, \dots, n_r}^{(s)}$ does not contain any copy of $kK_r^{(s)}$. Choose an r -tuple (x_1, x_2, \dots, x_r) from $V_1 \times V_2 \times \dots \times V_r$ uniformly at random. Let $T = \{x_1, x_2, \dots, x_r\}$ and $X(T)$ be the number of edges in $H[T]$. Then

$$\mathbb{E}(X(T)) = \sum_{S \in E(H)} \Pr(S \subset T) = \sum_{\substack{A: A \subset [r] \\ |A|=s}} \sum_{S \in E(V_A)} \frac{1}{n_A} = \sum_{\substack{A: A \subset [r] \\ |A|=s}} \frac{e(V_A)}{n_A}. \quad (3.1)$$

On the other hand, let m be the number of copies of $K_r^{(s)}$ in H . Define an r -partite r -graph H^* on the same vertex classes V_1, V_2, \dots, V_r . An r -element set S forms an edge in H^* if and only if $H[S]$ is a copy of $K_r^{(s)}$. Since H is $kK_r^{(s)}$ -free, it follows that the matching number of H^* is at most $k - 1$. Moreover, the number of edges in H^* is exactly

m . Then, by Lemma 2.1, we have $m \leq (k-1)n_2 \cdots n_r$. Let A_T be the event that $H[T]$ is a copy of $K_r^{(s)}$. Clearly, we have

$$Pr(A_T) = \frac{m}{n_1 n_1 \cdots n_r} \leq \frac{k-1}{n_1}.$$

Thus,

$$\begin{aligned} \mathbb{E}(X(T)) &= \mathbb{E}(X(T)|A_T)Pr(A_T) + \mathbb{E}(X(T)|\overline{A_T})Pr(\overline{A_T}) \\ &\leq \binom{r}{s} Pr(A_T) + \left(\binom{r}{s} - 1 \right) (1 - Pr(A_T)) \\ &= \binom{r}{s} - 1 + Pr(A_T) \\ &\leq \binom{r}{s} - 1 + \frac{k-1}{n_1}. \end{aligned} \tag{3.2}$$

Putting (3.1) and (3.2) together, we obtain that

$$\sum_{\substack{A: A \subset [r] \\ |A|=s}} e(V_A) \frac{1}{n_A} \leq \binom{r}{s} - 1 + \frac{k-1}{n_1}. \tag{3.3}$$

We consider the linear programming model (P1) as follows:

$$\begin{aligned} \max \quad & z = \sum_{\substack{A: A \subset [r] \\ |A|=s}} x_A \\ \text{s.t.} \quad & \sum_{\substack{A: A \subset [r] \\ |A|=s}} n_A^{-1} x_A \leq \binom{r}{s} - 1 + \frac{k-1}{n_1}, \\ & 0 \leq x_A \leq n_A, \quad A \in \binom{[r]}{s}. \end{aligned}$$

Applying Lemma 3.1 by setting $N = \binom{r}{s}$, $b = \binom{r}{s} - 1 + \frac{k-1}{n_1}$ and w_i be the i -th largest value in $\{n_A : A \in \binom{[r]}{s}\}$ for each $i \in 1, 2, \dots, \binom{r}{s}$ in (P), we have $M = \lfloor b \rfloor = \binom{r}{s} - 1$. Since $n_{[s]} \leq n_A$ for all $A \in \binom{[r]}{s}$, it follows that

$$a = w_{M+1}(b - M) = n_{[s]} \cdot \frac{k-1}{n_1} = (k-1)n_{[2,s]}.$$

Thus, the optimal value of (P1) is

$$\sum_{i=1}^M w_i + a = \sum_{\substack{A: A \subset [r] \\ |A|=s, A \neq [s]}} n_A + (k-1)n_{[2,s]} = g_k^{(s)}(n_1, n_2, \dots, n_r).$$

Let y be a vector indexed by the s -element subset A of $[r]$ with $y_A = e(V_A)$. Since $e(V_A) \leq n_A$ and the inequality (3.3) holds, it follows that y is a feasible solution to (P1). Therefore, we have

$$e(H) = \sum_{\substack{A: A \subset [r] \\ |A|=s}} e(V_A) = \sum_{\substack{A: A \subset [r] \\ |A|=s}} y_A \leq g_k^{(s)}(n_1, n_2, \dots, n_r).$$

Thus, the theorem follows. \square

4 The number of s -cliques in r -partite graphs

In this section, we first determine $ex(K_{n_1, \dots, n_r}, K_s, kK_r)$ for the case $n_1 \leq n_2 \leq n_3 = n_4 = \dots = n_r$. Then, by utilizing a result on rainbow matchings, we determine $ex(K_{n_1, \dots, n_r}, K_s, kK_r)$ for all n_1, \dots, n_r with $n_4 \geq r^r(k-1)k^{2r-2}$.

For an r -partite graph G on vertex classes V_1, V_2, \dots, V_r , we use $K_s(G)$ to denote the family of s -element subsets of $V(G)$ that form s -cliques in G and for $u \in V(G)$ we use $K_s(u, G)$ to denote the family of s -element subsets in $K_s(G)$ that contain u . For any $A \subset [r]$, we also use $K_s(V_A)$ to denote $K_s(G[V_A])$. Let $k_s(G)$, $k_s(u, G)$ and $k_s(V_A)$ be the cardinalities of $K_s(G)$, $K_s(u, G)$ and $K_s(V_A)$, respectively.

Proof of Theorem 1.5. Let V_1, V_2, \dots, V_r be the vertex classes such that $|V_i| = n_i$ for each $i = 1, 2, \dots, r$ and $n_4 = \dots = n_r = n_3$. Suppose $G \subseteq K(V_1, V_2, \dots, V_r)$ does not contain any copy of kK_r . Choose an r -tuple (x_1, x_2, \dots, x_r) from $V_1 \times V_2 \times \dots \times V_r$ uniformly at random. Let $T = \{x_1, x_2, \dots, x_r\}$ and $X(T)$ be the number of copies of K_s in $G[T]$. Then,

$$\mathbb{E}(X(T)) = \sum_{S \in K_s(G)} Pr(S \subset T) = \sum_{\substack{A: A \subset [r] \\ |A|=s}} \sum_{S \in K_s(V_A)} \frac{1}{n_A} = \sum_{\substack{A: A \subset [r] \\ |A|=s}} \frac{k_s(V_A)}{n_A}. \quad (4.1)$$

On the other hand, let m be the number of copies of K_r in G . By a similar argument as in the proof of Theorem 1.4, we have $m \leq (k-1)n_2n_3^{r-2}$. If $s = r$, then the theorem holds already (because $h_k^{(r)}(n_1, n_2, \dots, n_r) = (k-1)n_2 \dots n_r$), so we are left with the case $s \leq r-1$. Let A_T be the event that $H[T]$ is a copy of K_r . Clearly, we have

$$Pr(A_T) \leq \frac{k-1}{n_1}.$$

Since there are $\binom{r}{s}$ s -cliques in K_r and at most $\binom{r}{s} - \binom{r-2}{s-2}$ s -cliques in a graph on r vertices that is not a complete graph, it follows that

$$\begin{aligned} \mathbb{E}(X(T)) &= \mathbb{E}(X(T)|A_T)Pr(A_T) + \mathbb{E}(X(T)|\overline{A_T})Pr(\overline{A_T}) \\ &\leq \binom{r}{s}Pr(A_T) + \left(\binom{r}{s} - \binom{r-2}{s-2} \right) (1 - Pr(A_T)) \\ &= \binom{r}{s} - \binom{r-2}{s-2} + \binom{r-2}{s-2}Pr(A_T) \\ &\leq \binom{r}{s} - \binom{r-2}{s-2} + \frac{k-1}{n_1} \binom{r-2}{s-2}. \end{aligned} \quad (4.2)$$

Combining (4.1) and (4.2), we have

$$\sum_{\substack{A: A \subset [r] \\ |A|=s}} k_s(V_A) \frac{1}{n_A} \leq \binom{r}{s} - \binom{r-2}{s-2} + \frac{k-1}{n_1} \binom{r-2}{s-2}. \quad (4.3)$$

We consider the linear programming model (P2) as follows:

$$\begin{aligned} \max \quad & z = \sum_{\substack{A: A \subset [r] \\ |A|=s}} x_A \\ \text{s.t.} \quad & \sum_{\substack{A: A \subset [r] \\ |A|=s}} n_A^{-1} x_A \leq \binom{r}{s} - \binom{r-2}{s-2} + \frac{k-1}{n_1} \binom{r-2}{s-2}, \\ & 0 \leq x_A \leq n_A, \quad A \in \binom{[r]}{s}. \end{aligned}$$

Apply Lemma 3.1 by setting $N = \binom{r}{s}$, $b = \binom{r}{s} - \binom{r-2}{s-2} + \frac{k-1}{n_1} \binom{r-2}{s-2}$ and w_i be the i -th largest value in $\{n_A : A \in \binom{[r]}{s}\}$ for each $i \in 1, 2, \dots, \binom{r}{s}$ in (P). Note that $n_A = n_3^s$ for $A \in \binom{[3, r]}{s}$, $n_A = n_2 n_3^{s-1}$ for $A \in \binom{[2, r]}{s}$ and $2 \in A$, $n_A = n_1 n_3^{s-1}$ for $A \in \binom{[r] \setminus \{2\}}{s}$ and $1 \in A$, $n_A = n_1 n_2 n_3^{s-2}$ for $A \in \binom{[r]}{s}$ and $\{1, 2\} \subset A$. Since $n_3^s \geq n_2 n_3^{s-1} \geq n_1 n_3^{s-1} \geq n_1 n_2 n_3^{s-2}$ and

$$\binom{r-2}{s} + \binom{r-2}{s-1} + \binom{r-2}{s-1} = \binom{r-1}{s} + \binom{r-2}{s-1} = \binom{r}{s} - \binom{r-2}{s-2},$$

it follows that $w_i = n_1 n_2 n_3^{s-2}$ for $i \geq \binom{r}{s} - \binom{r-2}{s-2} + 1$. Since $M = \lfloor b \rfloor \geq \binom{r}{s} - \binom{r-2}{s-2}$, we have $w_{M+1} = n_1 n_2 n_3^{s-2}$. Thus, the optimal value of (P2) is

$$\begin{aligned} \sum_{i=1}^M w_i + a &= \sum_{i=1}^{\binom{r}{s} - \binom{r-2}{s-2}} w_i + \sum_{i=\binom{r}{s} - \binom{r-2}{s-2} + 1}^M w_i + w_{M+1}(b - M) \\ &= \sum_{\substack{A: A \subset [r] \\ |A|=s, \{1,2\} \not\subset A}} n_A + \sum_{i=\binom{r}{s} - \binom{r-2}{s-2} + 1}^M w_{M+1} + (b - M)w_{M+1} \\ &= \sum_{\substack{A: A \subset [r] \\ |A|=s, \{1,2\} \not\subset A}} n_A + \left(M - \binom{r}{s} + \binom{r-2}{s-2} + b - M \right) n_1 n_2 n_3^{s-2} \\ &= \sum_{\substack{A: A \subset [r] \\ |A|=s, \{1,2\} \not\subset A}} n_A + \frac{k-1}{n_1} \binom{r-2}{s-2} n_1 n_2 n_3^{s-2} \\ &= \sum_{\substack{A: A \subset [r] \\ |A|=s, \{1,2\} \not\subset A}} n_A + \sum_{\substack{A: A \subset [3, r] \\ |A|=s-2}} (k-1) n_2 n_A \\ &= h_k^{(s)}(n_1, n_2, \underbrace{n_3, \dots, n_3}_{r-2}). \end{aligned}$$

Let y be a vector indexed by s -element subset A of $[r]$ with $y_A = k_s(V_A)$. Since $k_s(V_A) \leq n_A$ and the inequality (4.3) holds, it follows that y is a feasible solution to (P2). Therefore, we obtain that

$$k_s(G) = \sum_{\substack{A: A \subset [r] \\ |A|=s}} k_s(V_A) \leq h_k^{(s)}(n_1, n_2, \underbrace{n_3, \dots, n_3}_{r-2}).$$

Thus, the theorem holds. \square

Let $f, k \geq 1$ be integers. A k -matching is a matching of size k . Given a coloring $c : E(G) \rightarrow [f]$ of the edges of an r -graph G , we call a matching $M \subset E(G)$ a rainbow matching if all its edges have distinct colors. An (f, k) -colored r -graph $G = (V, E)$ is an r -uniform multi-hypergraph whose edges are colored in f colors such that every color class contains a k -matching. Denote by $f(r, k)$ the largest number f of colors such that there exists an (f, k) -colored r -partite r -graph without a rainbow k -matching. Recently, Glebov, Sudakov and Szabó [13] gave an upper bound on $f(r, k)$.

Theorem 4.1. [13] For arbitrary integers $r, k \geq 2$, $f(r, k) < (r+1)^{r+1} (k-1) k^{2r}$.

Now we consider the maximum number of copies of K_s in a kK_r -free r -partite graph for $n_3 \leq n_4 \leq \dots \leq n_r$.

Proof of Theorem 1.6. Let $r \geq 4$, n_1, n_2 and n_3 be fixed integers. The proof is by induction on $(s, \sum_{i=4}^r (n_i - n_3))$. The base case of $s = 2$ is verified for all r and $n_1 \leq n_2 \leq \dots \leq n_r$ by Theorem 1.2. For every $s \geq 3$, the base case of $n_1 \leq n_2 \leq n_3 = n_4 = \dots = n_r$ is verified for all r by Theorem 1.5. Assume that for all r , the theorem holds for all pairs $(s', \sum_{i=4}^r (n'_i - n'_3))$ such that $s' < s$ or $s' = s$ together with $\sum_{i=4}^r (n'_i - n'_3) < \sum_{i=4}^r (n_i - n_3)$.

Suppose $G \subseteq K_{n_1, \dots, n_r}$ does not contain a copy of kK_r . Since $\sum_{i=4}^r (n_i - n_3) > 0$, there exists an $i \in [4, r]$ such that $n_i > n_{i-1}$. Without loss of generality, assume that $i = r$. For $u \in V_r$, let $G(u)$ denote the $(r-1)$ -partite graph on vertex classes V_1, \dots, V_{r-1} , and a pair $\{v_i, v_j\}$ forms an edge in $G(u)$ if and only if $\{u, v_i\}, \{u, v_j\}$ and $\{v_i, v_j\}$ are all edges in G . If there is a vertex $u \in V_r$ such that $G(u)$ is kK_{r-1} -free, then by induction on s , we have $k_s(u, G) = k_{s-1}(G(u)) \leq h_k^{(s-1)}(n_1, n_2, \dots, n_{r-1})$. Moreover, by induction on $\sum_{i=4}^r (n_i - n_3)$, we obtain that $k_s(G \setminus \{u\}) \leq h_k^{(s)}(n_1, n_2, \dots, n_{r-1}, n_r - 1)$. Therefore,

$$\begin{aligned} k_s(G) &= k_s(G \setminus \{u\}) + k_s(u, G) \\ &\leq h_k^{(s)}(n_1, n_2, \dots, n_{r-1}, n_r - 1) + h_k^{(s-1)}(n_1, n_2, \dots, n_{r-1}) \\ &= h_k^{(s)}(n_1, n_2, \dots, n_{r-1}, n_r). \end{aligned}$$

Otherwise, suppose that for all $u \in V_r$, there are at least k vertex-disjoint copies of K_{r-1} in $G(u)$. Since G is kK_r -free, we have $k \geq 2$. Let H be an $(r-1)$ -partite $(r-1)$ -uniform multi-hypergraph on vertex classes V_1, \dots, V_{r-1} . For any $u \in V_r$, if $\{u_1, \dots, u_{r-1}\}$ forms a copy of K_{r-1} in $G(u)$, let $\{u_1, \dots, u_{r-1}\}$ be an edge in H with color u . Then H is (n_r, k) -colored. Since $n_r \geq n_4 \geq r^r(k-1)k^{2r-2} > f(r-1, k)$, by Theorem 4.1, there is a rainbow k -matching $\{e_{i_1}, \dots, e_{i_k}\}$ in H . Thus, there are k vertices $\{u_{i_1}, \dots, u_{i_k}\} \subset V_r$ such that $\{e_{i_1} \cup \{u_{i_1}\}, \dots, e_{i_k} \cup \{u_{i_k}\}\}$ forms a kK_r in G , a contradiction. Thus, we complete the proof. \square

Acknowledgement. The authors would like to thank two anonymous referees for their helpful suggestions. The second author was supported by the National Natural Science Foundation of China (No. 11701407) and Scientific and Technological Innovation Programs of Higher Education Institutions in Shanxi (No. 183090222-S).

References

- [1] R. Aharoni and D. Howard, A rainbow r -partite version of the Erdős-Ko-Rado theorem, *Combin. Probab. Comput.* 26 (2017) 321–337.
- [2] N. Alon and C. Shikhelman, Many T copies in H -free graphs, *J. Combin. Theory Ser. B* 121 (2016) 146–172.
- [3] P. Bennett, S. English and M. Talanda-Fisher, Weighted Turán problems with applications, *Discrete Math.* 342 (2019) 2165–2172.
- [4] B. Bollobás, On genaralized graphs, *Acta Math. Acad. Sci. Hungar.* 16 (1965) 447–452.

- [5] B. Bollobás, D. E. Daykin and P. Erdős, Sets of independent edges of a hypergraph, Quart. J. Math. Oxford Ser. 21 (1976) 25–32.
- [6] J. De Silva, K. Heyse, A. Kapilow, A. Schenfisch and M. Young, Turán numbers of vertex-disjoint cliques in r -partite graphs, Discrete Math. 341 (2018) 492–496.
- [7] P. Erdős, A problem on independent r -tuples, Ann. Univ. Sci. Budapest. Eötvös Sect. Math. 8 (1965) 93–95.
- [8] P. Frankl, An extremal problem for two families of sets, European J. Combin. 3 (1982) 125–127.
- [9] P. Frankl, The shifting technique in extremal set theory, Surv. Combin. 123 (1987) 81–110.
- [10] P. Frankl, V. Rödl and A. Ruciński, On the maximum number of edges in a triple system not containing a disjoint family of a given size, Combin. Probab. Comput. 21 (2012) 141–148.
- [11] P. Frankl, Improved bounds for Erdős’ matching conjecture, J. Combin. Theory Ser. A 120 (2013) 1068–1072.
- [12] P. Frankl, On the maximum number of edges in a hypergraph with given matching number, Discrete Appl. Math. 216 (2017) 562–581.
- [13] R. Glebov, B. Sudakov and T. Szabó, How many colors guarantee a rainbow matching? Electron. J. Combin. 21 (2014) #P1.27.
- [14] J. Han and Y. Zhao, Turán number of disjoint triangles in 4-partite graphs, arXiv e-prints (Jun. 2019) arXiv:1906.01812.
- [15] P. Keevash, Hypergraph Turán problems, Surv. Combin. 392 (2011) 83–140.
- [16] J. Ma and Y. Qiu, Some sharp results on the generalized Turán numbers, European J. Combin. 84 (2019) <https://doi.org/10.1016/j.ejc.2019.103026>.
- [17] W. Mantel, Problem 28, Wiskundige Opgaven 10 (1907) 60–61.
- [18] A. Sidorenko, What we know and what we do not know about Turán numbers, Graphs Combin. 11 (1995) 179–199.
- [19] P. Turán, On an extremal problem in graph theory, (Hungarian), Mat. Fiz. Lapok 48 (1941) 436–452.
- [20] A.A. Zykov, On some properties of linear complexes, Mat. Sb. 66 (1949) 163–188.

A A proof of Theorem 1.1.

Lemma A.1. *For $n_1 \leq n_2 \leq n_3$ and $k \leq n_1$,*

$$ex(K_{n_1, n_2, n_3}, kK_2) = (k-1)(n_2 + n_3).$$

Proof. First, we prove the lemma for $n_1 = n_2 = n_3 = n$ by induction on k . Clearly, the lemma holds trivially for $k = 1$. We assume that the result holds for all k' with $k' < k \leq n$. Suppose G is a kK_2 -free 3-partite graph with vertex set $V = X \cup Y \cup Z$ and let

$$X = \{x_1, \dots, x_n\}, \quad Y = \{y_1, \dots, y_n\} \quad \text{and} \quad Z = \{z_1, \dots, z_n\}.$$

Define a partial order \prec on $X \cup Y \cup Z$ such that

$$x_1 \prec \dots \prec x_n, \quad y_1 \prec \dots \prec y_n, \quad z_1 \prec \dots \prec z_n,$$

and vertices from different parts are incomparable. Assume that G has maximal number of edges. Thus, $\nu(G) = k - 1$. By Lemma 2.2, we may further assume that G is stable. Let $T_0 = \{x_1, y_1, z_1\}$ and $G' = G \setminus T_0$. Furthermore, let $\nu(G') = t$ and let $M' = \{e_1, \dots, e_t\}$ be a largest matching in G' . If $G[T_0]$ is not a triangle, since G is stable, there exist two vertex sets $V_i, V_j \in \{X, Y, Z\}$ such that $G[V_i \cup V_j]$ is empty. It follows that G is a bipartite graph. Then by Lemma 2.1 with $r = 2$, we conclude that $e(G) \leq 2(k - 1)n$ and the lemma holds. If $G[T_0]$ is a triangle, then we have $k - 4 \leq \nu(G') \leq k - 2$, where $\nu(G') \leq k - 2$ follows from $G[T_0]$ being non-empty, and $\nu(G') \geq k - 4$ follows from there being only three vertices in T_0 and from $\nu(G) = k - 1$. The proof splits into three cases according to the value of $\nu(G')$.

Case 1. $\nu(G') = k - 2$. For every edge $\{u_i, v_i\} \in M'$, it is easy to see that the number of edges between $\{u_i, v_i\}$ and T_0 is at most 4 since G is a 3-partite graph. Thus, there are at most $4(k - 2)$ edge between $\cup_{e \in M'} e$ and T_0 . If $|\Gamma(T_0)| > 4(k - 2) + 3$, then we will find an edge between T_0 and $V(G') \setminus (\cup_{e \in M'} e)$. Without loss of generality, assume $\{x_1, u\}$ is such an edge. Then $M' \cup \{\{x_1, u\}, \{y_1, z_1\}\}$ forms a matching of size k , which contradicts the fact that G is kK_2 -free. If $|\Gamma(T_0)| \leq 4(k - 2) + 3$, then by the induction hypothesis, we have

$$\begin{aligned} e(G) &= |\Gamma(T_0)| + e(G') \\ &\leq 4(k - 2) + 3 + 2(k - 2)(n - 1) \\ &= 2(k - 1)n - 2n + 2k - 1 \\ &\leq 2(k - 1)n. \end{aligned}$$

Case 2. $\nu(G') = k - 3$. If $|\Gamma(T_0)| \leq 4n + 2(k - 3)$, then by the induction hypothesis, we have

$$e(G) = |\Gamma(T_0)| + e(G') \leq 4n + 2(k - 3) + 2(k - 3)(n - 1) = 2(k - 1)n.$$

Thus, the lemma holds. If $|\Gamma(T_0)| > 4n + 2(k - 3)$, let $G'' = G' \setminus (\cup_{e \in M'} e)$ and consider the edges between T_0 and $V(G'')$. Since there are at most $4(k - 3)$ edges between T_0 and $(\cup_{e \in M'} e)$, the number of edges between T_0 and $V(G'')$ is at least $4n + 2(k - 3) + 1 - 4(k - 3) - 3 = 4n - 2k + 4$. For any $u \in V(G)$ and $S \subset V(G)$, let $d(u, S)$ be the number of neighbors of u in S . Then, it follows that

$$d(x_1, V(G'')) + d(y_1, V(G'')) + d(z_1, V(G'')) \geq 4n - 2k + 4.$$

Since $(Y \cup Z) \setminus (\cup M') \setminus T_0$ has at most $2(n - 1) - (k - 3) = 2n - k + 1$ vertices, we have $d(x_1, V(G'')) \leq 2n - k + 1$. Similarly, $d(y_1, V(G'')) \leq 2n - k + 1$ and $d(z_1, V(G'')) \leq 2n - k + 1$. Therefore, for any $v \in \{x_1, y_1, z_1\}$, $d(v, V(G'')) \geq 4n - 2k + 4 - 2(2n - k + 1) = 2$.

It follows from Hall's theorem that there exist three disjoint edges $\{x_1, u_1\}, \{y_1, u_2\}$ and $\{z_1, u_3\}$ with $u_1, u_2, u_3 \in V(G'')$. These edges together with edges in M' form a matching of size k , which contradicts the fact that G is kK_2 -free.

Case 3. $\nu(G') = k - 4$. Since $|\Gamma(T_0)| < 6n$, by the induction hypothesis, we have

$$e(G) = |\Gamma(T_0)| + e(G') \leq 6n + 2(k - 4)(n - 1) \leq 2(k - 1)n.$$

Thus, the lemma holds for $n_1 = n_2 = n_3 = n$.

At last, we prove the lemma for the general case $n_1 \leq n_2 \leq n_3$ by induction on $n_2 + n_3 - 2n_1$. Since we've already proven the base case $n_2 + n_3 - 2n_1 = 0$, now assume that $n_2 + n_3 - 2n_1 > 0$. There exists $i = 2$ or 3 such that $n_i > n_{i-1}$. Without loss of generality, assume that $i = 3$. If there exists $v \in Z$ such that $d(v) \leq k - 1$, we have

$$\begin{aligned} e(G) &= d(v) + e(G \setminus v) \\ &\leq k - 1 + (k - 1)(n_2 + n_3 - 1) \\ &= (k - 1)(n_2 + n_3). \end{aligned}$$

If $d(v) \geq k$ for every $v \in Z$, since $|Z| \geq k$, it is easy greedily to find a matching of size k , a contradiction. Thus, we complete the proof. \square

Proof of Theorem 1.1. The cases $r = 2$ and $r = 3$ follow from Lemmas 2.1 and A.1, respectively. Thus, we are left with the case $r \geq 4$ which we prove by induction on k . Clearly, the result holds for $k = 1$. Assume that the result holds for all $k' < k$. Let $G \subseteq K_{n_1, \dots, n_r}$ be a kK_2 -free graph with the maximum number of edges. Thus, $\nu(G) = k - 1$. Denote by X_i the set of vertices in V_i with degree at least $2k - 1$ and put $x_i = |X_i|$ for $i = 1, \dots, r$. Let $n = n_1 + \dots + n_r$ and $x = x_1 + \dots + x_r$. Now we divide the proof into two cases according to the value of x .

Case 1. $x \geq 1$. Let $X = \bigcup_{i=1}^r X_i$ and $G' = G \setminus X$. Since $d(u) \geq 2k - 1$ for each $u \in X$, it is easy to see that $x \leq k - 1$ and $\nu(G') \leq k - 1 - x$ because otherwise one could greedily find a matching of size k . Let $\bar{x}_i = x - x_i$ and $n_{i_0} - x_{i_0} = \min_{i \in [r]} \{n_i - x_i\}$. By the induction hypothesis, we have

$$\begin{aligned} e(G) &= |\Gamma(X)| + e(G') \\ &\leq \sum_{i < j} x_i x_j + \sum_{i=1}^r x_i \left(\sum_{j \neq i} (n_j - x_j) \right) + (k - 1 - x) \left(\sum_{i=1}^r (n_i - x_i) - \min_{i \in [r]} \{n_i - x_i\} \right) \\ &= (k - 1)n - (k - 1)(x + n_{i_0} - x_{i_0}) + \sum_{i < j} x_i x_j + \sum_{i=1}^r x_i (n_{i_0} - x_{i_0} - (n_i - x_i)) \\ &\leq (k - 1)n - (k - 1)(n_{i_0} + \bar{x}_{i_0}) + \sum_{i < j} x_i x_j \\ &= (k - 1)(n - n_{i_0}) - (k - 1)\bar{x}_{i_0} + x_{i_0}\bar{x}_{i_0} + \sum_{\substack{i < j \\ i, j \neq i_0}} x_i x_j \\ &\leq (k - 1)(n - n_1) - \bar{x}_{i_0}^2 + \sum_{\substack{i < j \\ i, j \neq i_0}} x_i x_j \\ &= (k - 1)(n - n_1) - \sum_{i \neq i_0} x_i^2 - \sum_{\substack{i < j \\ i, j \neq i_0}} x_i x_j \\ &\leq (k - 1)(n_2 + \dots + n_r), \end{aligned}$$

where the second inequality follows from $n_{i_0} - x_{i_0} - (n_i - x_i) \leq 0$ and the third inequality follows from $n_{i_0} \geq n_1$ and $x_{i_0} + \bar{x}_{i_0} = x \leq k - 1$. Thus, the theorem holds.

Case 2. $x = 0$. Then all the vertices in G have degree at most $2k - 2$. Let $M = \{\{u_1, v_1\}, \dots, \{u_{k-1}, v_{k-1}\}\}$ be a largest matching of G , $A = \{u_1, \dots, u_{k-1}, v_1, \dots, v_{k-1}\}$ and $B = V(G) \setminus A$. Since M is a largest matching, B is an independent set of G . Let t_i be the number of edges between $\{u_i, v_i\}$ and B . We claim that $t_i \leq 2k - 2$. Otherwise, there exist $u, v \in B$ such that both $\{u_i, u\}$ and $\{v_i, v\}$ are edges of G , and then $M' \setminus \{\{u_i, v_i\}\} \cup \{\{u_i, u\}, \{v_i, v\}\}$ forms a matching of size k , a contradiction. Since $d(v) \leq 2k - 2$ for every $v \in V(G)$ and $d_B(u_i) + d_B(v_i) = t_i$, we have $d_A(u_i) + d_A(v_i) \leq 4k - 4 - t_i$. Thus, we have

$$\begin{aligned}
e(G) &= e(A, B) + e(A) \\
&= \sum_{v \in A} d_B(v) + \frac{1}{2} \sum_{v \in A} d_A(v) \\
&= \sum_{i=1}^{k-1} (d_B(u_i) + d_B(v_i)) + \frac{1}{2} \sum_{i=1}^{k-1} (d_A(u_i) + d_A(v_i)) \\
&\leq \sum_{i=1}^{k-1} t_i + \frac{1}{2} \sum_{i=1}^{k-1} (4k - 4 - t_i) \\
&= \frac{1}{2} \sum_{i=1}^{k-1} t_i + \frac{1}{2} (k-1)(4k-4) \\
&\leq (k-1)(3k-3) \\
&< (k-1)(n_2 + \dots + n_r),
\end{aligned}$$

where the last inequality follows from $r \geq 4$ and $n_1 \geq k$. Thus, we complete the proof. \square