

LARGE GLOBAL SOLUTIONS FOR NONLINEAR SCHRÖDINGER EQUATIONS II, MASS-SUPERCritical, ENERGY-SUBCRITICAL CASES

MARIUS BECEANU, QINGQUAN DENG, AVY SOFFER, AND YIFEI WU

ABSTRACT. In this paper, we consider the defocusing mass-supercritical, energy-subcritical nonlinear Schrödinger equation,

$$i\partial_t u + \Delta u = |u|^p u, \quad (t, x) \in \mathbb{R}^{d+1},$$

with $p \in (\frac{4}{d}, \frac{4}{d-2})$. We prove that under some restrictions on d, p , any radial function in the rough space $H^{s_0}(\mathbb{R}^d)$, for some $s_0 < s_c$ with the support away from the origin, there exists an incoming/outgoing decomposition, such that the initial data in the outgoing part leads to the global well-posedness and scattering forward in time; while the initial data in the incoming part leads to the global well-posedness and scattering backward in time. The proof is based on Phase-Space analysis of the nonlinear dynamics.

CONTENTS

1. Introduction	2
2. Preliminary	7
2.1. Some notations	7
2.2. Some basic lemmas	8
2.3. Linear Schrödinger operator	9
2.4. Some lemmas about Fourier integral operators	10
3. The incoming/outgoing waves	11
3.1. Definitions of the incoming/outgoing waves	11
3.2. Basic properties of the incoming/outgoing functions	13
3.3. Boundedness of the incoming/outgoing projection operators	26
4. Estimates on the incoming/outgoing linear flow	39
4.1. “Incoming/outgoing” decomposition of the linear flow	39
4.2. Improved Strichartz’s estimates	46
5. Proof of the Theorem 1.3	48
5.1. Definitions of the modified incoming and outgoing components	48
5.2. Space-time estimates	50
5.3. Morawetz estimates	52

2010 *Mathematics Subject Classification*. Primary 35Q55.

Key words and phrases. Nonlinear Schrödinger equation, global well-posedness, scattering.

5.4. Energy estimate	54
5.5. The proofs	56
Acknowledgements	57
References	57

1. INTRODUCTION

The aim of this work is the study of global existence and scattering theory for the inter-critical nonlinear Schrödinger equation (NLS) in 3 – 5 dimensions:

$$\begin{cases} i\partial_t u + \Delta u = \mu|u|^p u, \\ u(0, x) = u_0(x), \end{cases} \quad (1.1)$$

with $\mu = \pm 1, p > 0$. Here $u(t, x) : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$ is a complex-valued function. The case $\mu = 1$ is referred to the defocusing case, and $\mu = -1$ is referred to the focusing case. The class of solutions to equation (1.1) are invariant under the scaling

$$u(t, x) \rightarrow u_\lambda(t, x) = \lambda^{\frac{2}{p}} u(\lambda^2 t, \lambda x) \quad \text{for } \lambda > 0, \quad (1.2)$$

which maps the initial data as

$$u(0) \rightarrow u_\lambda(0) := \lambda^{\frac{2}{p}} u_0(\lambda x) \quad \text{for } \lambda > 0.$$

Denote

$$s_c = \frac{d}{2} - \frac{2}{p}.$$

Then the scaling leaves \dot{H}^{s_c} norm invariant, that is,

$$\|u_\lambda(t)\|_{\dot{H}^{s_c}} = \|u(\lambda^2 t)\|_{\dot{H}^{s_c}}, \quad \|u(0)\|_{\dot{H}^{s_c}} = \|u_\lambda(0)\|_{\dot{H}^{s_c}},$$

which is called *critical regularity* s_c . It is also considered as the lowest regularity that problem (1.1) is well-posed for general $H^s(\mathbb{R}^d)$ -data. Indeed, one can find some special initial datum belonging to $H^s(\mathbb{R}^d)$, $s < s_c$ such that the problem (1.1) is ill-posed.

The H^1 -solution of equation (1.1) also enjoys the mass, momentum and energy conservation laws, which read

$$\begin{aligned} M(u(t)) &:= \int |u(t, x)|^2 dx = M(u_0), \\ P(u(t)) &:= \operatorname{Im} \int \overline{u(t, x)} \nabla u(t, x) dx = P(u_0), \\ E(u(t)) &:= \frac{1}{2} \int |\nabla u(t, x)|^2 dx + \frac{\mu}{p+2} \int |u(t, x)|^{p+2} dx = E(u_0). \end{aligned} \quad (1.3)$$

The well-posedness and scattering theory for Cauchy problem (1.1) with initial data in $H^s(\mathbb{R}^d)$ were extensively studied, which we here briefly review. The local well-posedness theory follows from a standard fixed point argument, implying that for all $u_0 \in H^s(\mathbb{R}^d)$, $s \geq s_c$, there exists $T_0 > 0$ such that its corresponding solution $u \in C([0, T_0], H^s(\mathbb{R}^d))$. In fact, the above T_0 depends on $\|u_0\|_{H^s(\mathbb{R}^d)}$ when $s > s_c$ and also the profile of u_0 when $s = s_c$. Some of the results can be found in Cazenave and Weissler [11].

The fixed point argument used in local theory can be applied directly to prove the global well-posedness for solutions to equation (1.1) with small initial data in $H^s(\mathbb{R}^d)$ with $s \geq s_c$. In the mass-supercritical, energy-subcritical cases, that is, $\frac{4}{d} < p < \frac{4}{d-2}$, if we consider the solution in energy space $H^1(\mathbb{R}^d)$, the local theory above together with conservation laws (1.3), yields the global well-posedness for all initial data $u_0 \in H^1(\mathbb{R}^d)$ in the defocusing case $\mu = 1$, and for any initial data $u_0 \in H^1(\mathbb{R}^d)$ with some restrictions in the focusing case. Furthermore, the scattering under the same conditions were also obtained by Ginibre, Velo [22] in the defocusing case and Duyckaerts, Holmer and Roudenko [18] in the focusing case.

Recently, conditional global and scattering results for $s_c > 1$ with the assumption of $u \in L_t^\infty(I, \dot{H}_x^{s_c}(\mathbb{R}^d))$ were considered by many authors, beginning with the major work [25, 26], and then developed by [9, 17, 19, 20, 36, 37] and cited references. That is, if the initial data $u_0 \in \dot{H}^{s_c}(\mathbb{R}^d)$ and the solution has a priori estimate

$$\sup_{0 < t < T_{out}(u_0)} \|u(t)\|_{\dot{H}_x^{s_c}(\mathbb{R}^d)} < +\infty, \quad (1.4)$$

then $T_{out}(u_0) = +\infty$ and the solution scatters in $\dot{H}^{s_c}(\mathbb{R}^d)$, here $[0, T_{out}(u_0))$ is the maximal interval for existence of the solution. Consequently, these results give blowup criterion in which the lifetime only depends on the critical norm $\|u\|_{L_t^\infty \dot{H}_x^{s_c}(\mathbb{R}^d)}$. However, no such large global results are known for general initial data $u_0 \in \dot{H}^{s_c}(\mathbb{R}^d)$.

In the case when $s_c < 1$ the use of a priori estimates on the solution as a condition was also developed. The work of Bourgain [6] on the NLS, made assumptions on the space-time norm of the solution in space-time subsets, by deriving necessary conditions for blow-up. For example, Kenig and Merle [25] proved that for the NLS in the intercritical case, in dimension 3 and cubic nonlinearity case, global existence and scattering hold under the condition

$$\sup_{0 < t < T_{out}(u_0)} \|u(t)\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^d)} < \infty.$$

See also for examples some developments in [29, 49].

These conditional results, and other works, used the elaborate method introduced by Kenig-Merle, involving profile decomposition, concentration compactness and the localized Strichartz estimates. In contrast, our results use explicit conditions on the *initial data alone*. A further new consequence of our analysis is that supercritical/rough solutions exist, which are large and that in fact the standard assumption of initial data being in the space H^{s_c} is not required.

We follow here a new paradigm, based on phase space analysis and propagation estimates, to give explicit conditions on the initial data, which implies global existence and scattering. This will allow us to give explicit conditions on large and rough initial data, for which global existence and scattering hold. Our method does not use the above techniques based on profile decomposition and concentration compactness.

When the initial data is rough, $s < s_c$, we do not have general global well-posedness. Yet, as we will show, generic conditions for global existence and scattering allow such initial data. So, we conclude that rough initial data does not guarantee blowup.

Rough data may appear naturally in some applications: initial data corrupted with noise (as in nonlinear optics applications), and in the construction of invariant measures for the NLS dynamics, see [4, 5] and further developments in [15, 16, 30, 40, 41, 46] and the cited

references. In these last cases it is known that the relevant measure is supported on rough spaces.

The phase-space analysis we use seeks to identify the initial data that can not move into the origin and blow-up. So, naturally, this analysis requires the distinction between outgoing waves under the free flow, which move away from the origin (for positive times), and the incoming waves which move towards the origin.

In particular we show, that under some extra conditions, initial data which is outgoing, will lead to global solution, even if it is rough, that is, the data belongs to the space H^s , with $s < s_c$, where s_c is the critical Sobolev norm. While it looks unsurprising, it is in fact a subtle result!

Under the free flow this initial data (radial) will move away from the origin, and therefore will get smaller in L^∞ , by the assumption of radial symmetry, and some H^s regularity. Moreover, since the speed depends on the frequency, it gets smaller and faster for higher frequencies. Indeed, as an estimate essentially proved in Proposition 4.1 below, we show that

$$\|\chi_{\leq 1+Nt} e^{it\Delta} P_{\geq N} f_{out}\|_{H^1} \lesssim \|f\|_{L^2}. \quad (1.5)$$

However, the property of being outgoing is not necessarily preserved by the nonlinearity. Furthermore, in a major difference from the wave equation case (see [2, 3]), part of the solution moves *backward* towards the origin, even under the free flow. Since the data is rough, we do not have local existence (see the ill-posedness results in [13] and [27]), so, we can not move a short time forward, in the usual perturbative way e.g., as in [38].

To counter these effects, we need optimal propagation estimates in space, frequency and time. In particular we need to use the extra smoothing effect for waves which move in the “wrong” direction, the classical forbidden regions, see (1.5). We also need gain of regularity in the Strichartz estimate for radial functions, which allows us to show that in fact the Duhamel part, contributed by the nonlinearity is essential in H^1 , which is based on the following result. A more general result will be stated in Section 4.

Proposition 1.1. *Let $d = 3, 4, 5$. For any radial function $f \in L^2(\mathbb{R}^d)$ satisfying*

$$\text{supp } f \subset \{x : |x| \geq 1\},$$

there exists some suitable decomposition, says incoming and outgoing decomposition,

$$f = f_+ + f_-,$$

such that the following supercritical space-time estimate holds for any $N \geq 1$,

$$\|e^{\pm it\Delta} P_N f_{\pm}\|_{L_t^\infty H_x^1(\mathbb{R}^+ \times \mathbb{R}^d) + L_t^2 L_x^\infty(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim \|P_N f\|_{L^2}.$$

Remark 1.2. In general, the standard Strichartz estimates (see [24]) imply

$$\|e^{\pm it\Delta} P_N f\|_{L_t^2 L_x^\infty(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim N^{\frac{d}{2}-1} \|P_N f\|_{L^2},$$

which is scaling invariant and thus the index $\frac{d}{2}-1$ is optimal. Hence, the decomposition above presents a way in which we are able to obtain a kind of supercritical Strichartz estimates.

The proposition is based on the decomposition

$$L^2 \cap \{f : \text{supp } f \subset \{|x| \geq 1\}\} = L_+^2 + L_-^2,$$

for which the spaces L_{\pm}^2 can be roughly described that if $f \in L_+^2$, then for any $t > 0$,

$$e^{it\Delta}P_N f = \chi_{\geq \delta \langle Nt \rangle} F(t) + O(N^{-a} \langle t \rangle^{-b}), \quad \text{for some } F(t) \in L^2;$$

and if $f \in L_-^2$, then the equality above holds for any $t < 0$. Here $\delta > 0$ is a small constant, $a > 0$ and $b > 0$.

More precisely, if f is supported away from zero, then we can write $f = f_+ + f_- + f_0$ with $f_0 \in H^1$, $f_+ = P_+ f \in L_+^2$, $f_- = P_- f \in L_-^2$, where P_{\pm} are incoming/outgoing projection operators defined in Definition 3.4. The crucial step in the construction above is the proof we give that the range of P_+ is almost invariant (in the sense above) under the free flow for $t > 0$, and similarly for P_- , $t < 0$. Since the problem under study is energy subcritical, the solution is stable under H^1 perturbations. We emphasize that P^{\pm} are not completely invariant under the free flow. There is always an H^1 correction. Moreover if it is true (invariance of the range) at one time, it will not be true at later times. These present obstacles in the study of the long time behavior of the solution.

These estimates together with the approximate energy identity are applied to the part w of the solution. w is defined by the following decomposition for the nonlinear solution:

$$w = u - e^{it\Delta}(P_{\geq N} f)_+,$$

(with a slight modification). Then, w satisfies the following equation:

$$i\partial_t w + \Delta w = |u|^p u.$$

Therefore, by showing that w is in H^1 , we conclude that all the singular part, is carried away as a free wave to infinity.

Hence, since the problem is energy subcritical, we can control the effect of the nonlinearity, even though it is large. One should note however, that better, supercritical smoothing estimates only hold for outgoing waves.

Our construction of the projections on incoming/outgoing waves follows a similar approach of T. Tao [44]. We make a different decomposition near the origin.

A comparison to such in/out decompositions in scattering theory may be illuminating: the problem of global existence is the exact dual in the phase space, of the scattering problem. For global existence we need to control short time behavior, near the origin in space, at high frequency; the complete opposite of scattering. In scattering theory the decomposition into in and out waves, which was inspired by Enss method, leads to various definitions of such projections, beginning with the works of Mourre [39]. Mourre constructed the projection by taking the projection on the positive spectral part of the dilation generator A , defined as $x \cdot p + p \cdot x \equiv 2A$. Here x is multiplication by x in space and p , the momentum operator in Quantum Mechanics, is defined by $p \equiv -i\nabla_x$. He defined the projections on the incoming/outgoing parts by $P^{\pm}(A)$, where $P^{\pm}(y)$ stand for the characteristic functions, in y , of the positive/negative real axis.

Clearly A is a nice PDO, but not function of A .

Then, propagation estimates and other constructions, including estimates based on replacing A by $\gamma \equiv f(x)x \cdot p + p \cdot f(x)x$ were developed by Sigal and Soffer [47]. Here $f(x) \sim 1/|x|$, $|x| > 1$. In this construction, the “out” component of the solution still has some small portion moving inward, but this portion decays fast in time. See however [48]. Unlike the constructions we use, which are adapted to the radial nonlinear problem, the

scattering theory constructions above apply to non-radial functions, and so are the known estimates.

However, the propagation estimates obtained are not suitable, as they stand, for the dual space problem. They apply for general Hamiltonians, not just the Laplacian. It may be used for short times as well, when the frequency is large; that is still not explicit in the literature. The problem comes from the fact that near zero in space, the frequency can be very large, and yet A and γ remain bounded.

In the nonlinear context, this problem is far more severe, since the high frequency part is not stable under nonlinear perturbations.

New constructions for in/out decomposition were first introduced by T. Tao [44]. They are based on a decomposition in terms of spherical waves, of the form e^{ikr}/r , in three dimensions.

Still the problem at zero remains, much work was done to deal with this part of the phase-space. Further works in this direction, and others, sharp propagation estimates were done in e.g. Killip, Tao, Visan [31], and Li and Zhang [28, 33, 34].

We follow, up to some modifications near the origin, similar constructions and estimates.

We will use them in a different way, to localize and propagate rough initial data in particular, so as to get explicit conditions for global existence and scattering. Our strategy is to show that in some sense the phase-space localization of the initial data is stable, up to smooth corrections. So, we show that the solution is a sum of the linear, rough/supercritical part plus a correction coming from the Duhamel term (the nonlinear part).

By using radial symmetry, we show there is a gain of regularity, and the Duhamel term contributes essentially in H^1 space, the correction w . It will also follow, that if the initial w is small, then it can be controlled in the focusing case. For this we need only to have the initial data small in H^s , for some $s < s_c$.

Therefore, since the nonlinearity is inter-critical, and we have the Morawetz estimates and energy estimates in hand, we can then get global existence theory as in the H^1 case, after using also frequency cut-off and a continuity argument.

Similar improved smoothing for the Duhamel term was obtained before by Bourgain [7], in cases where the data is *subcritical* and below H^1 .

In the defocusing case, we can then cover outgoing initial data of arbitrary size, and in the focusing case small (but in the rough/supercritical norm!) data. Since the initial data is supercritical, in general, there is no well-posedness; to achieve that, we need the initial data to be supported at some positive distance away from the origin in space, together with the outgoing condition.

Our main result is following, which we focus on the defocusing case.

Theorem 1.3. *Let $\mu = 1$, $d = 3, 4, 5$. Then there exist $s_0 < 1$ and $p_1(d) < \frac{4}{d-2}$, such that for any $p \in [p_1(d), \frac{4}{d-2})$, the following is true. Suppose that f is a radial function for which there exists $\varepsilon_0 > 0$ such that*

$$\chi_{\leq \varepsilon_0} f \in H^1(\mathbb{R}^d), \quad (1 - \chi_{\leq \varepsilon_0}) f \in H^{s_0}(\mathbb{R}^d). \quad (1.6)$$

Then the solution u to the equation (1.1) with the initial data

$$u_0 = f_+ \quad (\text{or } u_0 = f_-)$$

exists globally forward (or backward) in time, and u is unique in $C(\mathbb{R}^+; H^{s_0}(\mathbb{R}^d)) \cap X_+$ (or $u \in C(\mathbb{R}^-; H^{s_0}(\mathbb{R}^d)) \cap X_-$), in which X_\pm are some auxiliary space-time space. Moreover, the solution depends continuously on the initial data in $H^{s_0}(\mathbb{R}^d)$. Here f_+ and f_- are the modified outgoing and incoming components of f respectively, which are given in Proposition 1.1. Furthermore, there exists $u^+ \in H^{s_0}(\mathbb{R}^d)$ (or $u^- \in H^{s_0}(\mathbb{R}^d)$), such that when $t \rightarrow +\infty$ (or $t \rightarrow -\infty$),

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} u^+\|_{H^1(\mathbb{R}^d)} = 0 \quad (\text{or } \lim_{t \rightarrow -\infty} \|u(t) - e^{it\Delta} u^-\|_{H^1(\mathbb{R}^d)} = 0). \quad (1.7)$$

Remark 1.4. We make several remarks regarding the above statements.

- (1). Note that s_0 is independent of p , hence $s_0 < s_c$ when p is close to $\frac{4}{d-2}$. Moreover, since $f = f_+ + f_-$, if f is not in $H^{s_c}(\mathbb{R}^d)$, at least one of f_+ and f_- is not in $H^{s_c}(\mathbb{R}^d)$. Therefore, we obtain a class of the global solutions for the defocusing energy-subcritical nonlinear Schrödinger equation in the supercritical space $H^{s_0}(\mathbb{R}^d)$.
- (2). It is worth noting that there is no smallness restriction in Theorem 1.3. Here we are not going to pursue the sharp conditions on s_0 and $p_1(d)$ in this paper.

Moreover, the restriction on the dimensions $d = 3, 4, 5$ is not essential, and the analogous results are valid in more general dimensions. However, for the sake of readability, we will not go into details.

- (3). The theorem implies that the incoming/outgoing solution has the “smoothing effect”. Indeed, we can show that for initial data belonging to $L^2(\mathbb{R}^d)$, there exist some $s_* > 0, r_0 > 2$, such that the solution $u(t)$ of (1.1) corresponding to such initial data, is in $W^{s,r}(\mathbb{R}^d)$ for any $t > 0$, any $r \in (2, r_0]$ and any $s \in [0, s_*]$, and moreover, the L^r -norm decays at infinite time in the sense that

$$\|u(t)\|_{L^r(\mathbb{R}^d)} \rightarrow 0, \quad \text{as } t \rightarrow +\infty.$$

- (4). By rescaling, we only need to prove the theorem when $\varepsilon_0 = 1$.

Organization of the paper. In Section 2, we give some preliminaries, which include some basic lemmas, some estimates on the linear Schrödinger operator and the Fourier integral operators. In Section 3, we give the definitions of the incoming/outgoing waves, their basic properties, the boundness in $\dot{H}^{s_c}(\mathbb{R}^d)$. In Section 4, we give some supercritical spacetime estimates on the incoming/outgoing linear flow. In Section 5, we give the proof of the main theorem.

2. PRELIMINARY

2.1. Some notations. We write $X \lesssim Y$ or $Y \gtrsim X$ to indicate $X \leq CY$ for some constant $C > 0$. If C depends upon some additional parameters, we will indicate this with subscripts; for example, $X \lesssim_a Y$ denotes the assertion that $X \leq C(a)Y$ for some $C(a)$ depending on a . We use $O(Y)$ to denote any quantity X such that $|X| \lesssim Y$. We use the notation $X \sim Y$ whenever $X \lesssim Y \lesssim X$.

The notation $a+$ denotes $a + \epsilon$ for any small ϵ , and $a-$ for $a - \epsilon$. $|\nabla|^\alpha = (-\Delta)^{\alpha/2}$. $\langle \cdot \rangle = (1 + |\cdot|^2)^{\frac{1}{2}}$. We denote $\mathcal{S}(\mathbb{R}^d)$ to be the Schwartz Space in \mathbb{R}^d , and $\mathcal{S}'(\mathbb{R}^d)$ to be the topological dual of $\mathcal{S}(\mathbb{R}^d)$. Let $h \in \mathcal{S}'(\mathbb{R}^{d+1})$, we use $\|h\|_{L_t^q L_x^p}$ to denote the mixed norm $\left(\int \|h(\cdot, t)\|_{L^p}^q dt \right)^{\frac{1}{q}}$, and $\|h\|_{L_{xt}^q} := \|h\|_{L_x^q L_t^q}$. Sometimes, we use the notation $q' = \frac{q}{q-1}$.

Throughout this paper, we use $\chi_{\leq a}$ for $a \in \mathbb{R}^+$ to be the smooth function

$$\chi_{\leq a}(x) = \begin{cases} 1, & |x| \leq a, \\ 0, & |x| \geq \frac{101}{100}a. \end{cases}$$

Moreover, we denote $\chi_{\geq a} = 1 - \chi_{\leq a}$, $\chi_{a \leq \cdot \leq b} = \chi_{\leq b} - \chi_{\leq a}$ and $\chi_a = \chi_{\leq 2a} - \chi_{\leq a}$ for short.

Also, we need some Fourier operators. For each number $N > 0$, we define the Fourier multipliers $P_{\leq N}, P_{\geq N}, P_N$ as

$$\begin{aligned} \widehat{P_{\leq N}f}(\xi) &:= \chi_{\leq N}(\xi) \hat{f}(\xi), \\ \widehat{P_{\geq N}f}(\xi) &:= \chi_{\geq N}(\xi) \hat{f}(\xi), \\ \widehat{P_N f}(\xi) &:= \chi_N(\xi) \hat{f}(\xi). \end{aligned}$$

We will usually use these multipliers when N are *dyadic numbers* (that is, of the form 2^k for some integer k).

2.2. Some basic lemmas. First, we need the following radial Sobolev embedding, see, e.g., [45].

Lemma 2.1. *Let α, q, p, s be the parameters which satisfy*

$$\alpha > -\frac{d}{q}; \quad \frac{1}{q} \leq \frac{1}{p} \leq \frac{1}{q} + s; \quad 1 \leq p, q \leq \infty; \quad 0 < s < d$$

with

$$\alpha + s = d\left(\frac{1}{p} - \frac{1}{q}\right).$$

Moreover, at most one of the equalities holds:

$$p = 1, \quad p = \infty, \quad q = 1, \quad q = \infty, \quad \frac{1}{p} = \frac{1}{q} + s.$$

Then for any radial function u such that $|\nabla|^s u \in L^p(\mathbb{R}^d)$,

$$\| |x|^\alpha u \|_{L^q(\mathbb{R}^d)} \lesssim \| |\nabla|^s u \|_{L^p(\mathbb{R}^d)}.$$

The second is the following fractional Leibniz rule, see [8, 32] and the references therein.

Lemma 2.2. *Let $0 < s < 1$, $1 < p \leq \infty$, and $1 < p_1, p_2, p_3, p_4 \leq \infty$ with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, $\frac{1}{p} = \frac{1}{p_3} + \frac{1}{p_4}$, and let $f, g \in \mathcal{S}(\mathbb{R}^d)$, then*

$$\| |\nabla|^s(fg) \|_{L^p} \lesssim \| |\nabla|^s f \|_{L^{p_1}} \|g\|_{L^{p_2}} + \| |\nabla|^s g \|_{L^{p_3}} \|f\|_{L^{p_4}}.$$

A simple consequence is the following elementary inequality, see [1] for the proof.

Lemma 2.3. *For any $a > 0$, $1 < p \leq \infty$, $0 \leq \gamma < \frac{d}{p}$, and $|\nabla|^\gamma g \in L^p(\mathbb{R}^d)$,*

$$\| |\nabla|^\gamma (\chi_{\leq a} g) \|_{L^p(\mathbb{R}^d)} \lesssim \| |\nabla|^\gamma g \|_{L^p(\mathbb{R}^d)}. \quad (2.1)$$

Here the implicit constant is independent of a . The same estimate holds for $\chi_{\geq a} g$.

Moreover, we need the following mismatch result, which is helpful in commuting the spatial and the frequency cutoffs.

Lemma 2.4 (Mismatch estimates, see [33]). *Let ϕ_1 and ϕ_2 be smooth functions obeying*

$$|\phi_j| \leq 1 \quad \text{and} \quad \text{dist}(\text{supp}\phi_1, \text{supp}\phi_2) \geq A,$$

for some large constant A . Then for $\sigma \geq 0$, $M \geq 1$, $1 \leq r \leq q \leq \infty$ and for any $m \geq 0$,

$$\|\phi_1 |\nabla|^\sigma P_{\geq M}(\phi_2 f)\|_{L_x^q(\mathbb{R}^d)} \lesssim M^{\sigma-m} A^{-m+\frac{d}{q}-\frac{d}{r}} \|\phi_2 f\|_{L_x^r(\mathbb{R}^d)}. \quad (2.2)$$

Remark 2.5. In this paper, we will frequently use the following estimate from Lemma 2.4,

$$\|\chi_{\leq \frac{1}{4}} P_M(\chi_{\geq 1} f)\|_{L^\infty(\mathbb{R}^d)} \lesssim M^{-m} \|\tilde{P}_M(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}, \quad \text{for any } m \geq 0,$$

where \tilde{P}_N is defined by

$$\widehat{\tilde{P}_N f}(\xi) := \chi_{2^{10}N \leq \cdot \leq 2^{10}N}(\xi) \hat{f}(\xi).$$

In the following, we shall slightly abuse notation and write \tilde{P}_N by P_N .

Furthermore, we need the following elementary formulas, see [1].

Lemma 2.6. *Let $f \in \mathcal{S}(\mathbb{R}^d)^d$ and $g \in \mathcal{S}(\mathbb{R}^d)$, then for any integer N ,*

$$\nabla_\xi \cdot (f \nabla_\xi)^{N-1} \cdot (fg) = \sum_{\substack{l_1, \dots, l_N \in \mathbb{N}^d, l' \in \mathbb{N}^d; \\ |l_j| \leq j; |l_1| + \dots + |l_N| + |l'| = N}} C_{l_1, \dots, l_N, l'} \partial_\xi^{l_1} f \cdots \partial_\xi^{l_N} f \partial_\xi^{l'} g,$$

where we have used the notations

$$\nabla_\xi = \{\partial_{\xi_1}, \dots, \partial_{\xi_d}\}; \quad \partial_\xi^l f = \sum_{j=1}^d C_j \partial_{\xi_1}^{l^1} \cdots \partial_{\xi_d}^{l^d} f_j, \quad \text{for any } l = \{l^1, \dots, l^d\} \in \mathbb{N}^d.$$

Here $C_{l_1, \dots, l_N, l'}$, C_j are some absolute constants only depending on N and d .

2.3. Linear Schrödinger operator. Let the operator $S(t) = e^{it\Delta}$ be the linear Schrödinger flow, that is,

$$(i\partial_t + \Delta)S(t) \equiv 0.$$

The following are some fundamental properties of the operator $e^{it\Delta}$. The first is the explicit formula, see, e.g., Cazenave [10].

Lemma 2.7. *For all $\phi \in \mathcal{S}(\mathbb{R}^d)$, $t \neq 0$,*

$$S(t)\phi(x) = \frac{1}{(4\pi it)^{\frac{d}{2}}} \int_{\mathbb{R}^d} e^{\frac{ix-y|^2}{4t}} \phi(y) dy.$$

Moreover, for any $r \geq 2$,

$$\|S(t)\phi\|_{L_x^r(\mathbb{R}^d)} \lesssim |t|^{-d(\frac{1}{2}-\frac{1}{r})} \|\phi\|_{L^{r'}(\mathbb{R}^d)}.$$

The following is the standard Strichartz estimate, see for example [24].

Lemma 2.8. *Let I be a time interval and let $u : I \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a solution to the inhomogeneous Schrödinger equation*

$$iu_t - \Delta u + F = 0.$$

Then for any $t_0 \in I$, any pairs (q_j, r_j) , $j = 1, 2$ satisfying

$$q_j \geq 2, \quad r_j \geq 2, \quad \frac{2}{q_j} + \frac{d}{r_j} = \frac{d}{2}, \quad \text{and} \quad (q_j, r_j, d) \neq (2, +\infty, 2),$$

the following estimate holds,

$$\|u\|_{C(I; L^2(\mathbb{R}^d))} + \|u\|_{L_t^{q_1} L_x^{r_1}(I \times \mathbb{R}^d)} \lesssim \|u(t_0)\|_{L_x^2(\mathbb{R}^d)} + \|F\|_{L_t^{q'_2} L_x^{r'_2}(I \times \mathbb{R}^d)}.$$

We also need the special Strichartz estimates for radial data, which were firstly proved by Shao [42], and then developed in [12, 23].

Lemma 2.9 (Radial Strichartz estimates). *Let $g \in L^2(\mathbb{R}^d)$ be a radial function, k be an integer, then for any triple (q, r, γ) satisfying*

$$\gamma \in \mathbb{R}, \quad q \geq 2, \quad r > 2, \quad \frac{2}{q} + \frac{2d-1}{r} < \frac{2d-1}{2}, \quad \text{and} \quad \frac{2}{q} + \frac{d}{r} = \frac{d}{2} + \gamma, \quad (2.3)$$

we have that

$$\| |\nabla|^\gamma e^{it\Delta} g \|_{L_t^q L_x^r(\mathbb{R} \times \mathbb{R}^d)} \lesssim \|g\|_{L^2(\mathbb{R}^d)}.$$

Furthermore, let $F \in L_t^{\tilde{q}'} L_x^{\tilde{r}'}(\mathbb{R}^{d+1})$ be a radial function in x , then

$$\left\| \int_0^t e^{i(t-s)\Delta} F(s) ds \right\|_{L_t^q L_x^r(\mathbb{R}^{d+1})} + \left\| |\nabla|^{-\gamma} \int_0^t e^{i(t-s)\Delta} F(s) ds \right\|_{L_t^\infty L_x^2(\mathbb{R}^{d+1})} \lesssim \|F\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}'}(\mathbb{R}^{d+1})},$$

where the triples (q, r, γ) , $(\tilde{q}, \tilde{r}, -\gamma)$ satisfy (2.3).

2.4. Some lemmas about Fourier integral operators. The following are some lemmas related to the estimate of the Fourier integral operators, we refer to Stein [43] for the proofs. The first one is from the application of the “stationary phase” theory.

Lemma 2.10. *Let ϕ, ψ be smooth functions defined in \mathbb{R} and $\lambda \in \mathbb{R}^+$, and ϕ satisfies*

$$\phi(x_0) = \phi'(x_0) = 0, \quad \text{and} \quad \phi''(x_0) \neq 0.$$

If ψ is supported in a sufficiently small neighborhood of x_0 , then

$$\int_{\mathbb{R}} e^{i\lambda\phi(x)} \psi(x) dx = a_0 \lambda^{-\frac{1}{2}} + O(\lambda^{-\frac{3}{2}}), \quad \text{when } \lambda \rightarrow +\infty,$$

with

$$a_0 = \psi(x_0) \left(\frac{2\pi}{i\phi''(x_0)} \right)^{\frac{1}{2}}.$$

The second result is the estimate of the Fourier integral operator, which can be regarded as an extension of the Plancherel identity. The following result can be found in [21].

Proposition 2.11. *Let T be the Fourier integral operator given by*

$$Tf(x) = \int_{\mathbb{R}^d} e^{ix \cdot \xi} a(x, \xi) \chi_{\leq 1}(\xi) \hat{f}(\xi) d\xi. \quad (2.4)$$

Suppose that $a(x, \xi) \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ and satisfies that for all multi-indices $\alpha \in \mathbb{R}^d$,

$$\sup_{\xi \in \mathbb{R}^d} \langle \xi \rangle^{|\alpha|} \left\| \partial_\xi^\alpha a(\cdot, \xi) \right\|_{L_x^\infty(\mathbb{R}^d)} < +\infty.$$

Then the operator T defined in (2.4) satisfies that

$$T : L^p(\mathbb{R}^d) \mapsto L^p(\mathbb{R}^d), \quad \text{is bounded, for all } 1 < p < \infty.$$

As an immediate consequence of the proposition above, we have the estimates on the following Fourier integral operators which are used in this paper.

Corollary 2.12. *Let $\beta \in \mathbb{R}, N > 0$, and let T be the Fourier integral operator given by*

$$Tf(x) = \int_{\mathbb{R}^d} e^{i\beta x \cdot \xi} \chi_{\leq 1}(x \cdot \xi) \chi_{\leq N}(\xi) f(\xi) d\xi. \quad (2.5)$$

Then there exists some $C > 0$ which is independent of N, β , such that

$$\|Tf\|_{L^2} \leq C|\beta|^{-\frac{d}{2}} \|\chi_{\leq N} f\|_{L^2}.$$

The same estimate holds if $\chi_{\leq 1}$ is replaced by $\chi_{\geq 1}, \chi'_{\leq 1}, \chi'_{\geq 1}$ or $\chi''_{\geq 1}$ in (2.5).

Proof. Since the operator T is scaling invariant in L^2 , we can use the rescaling argument. Indeed, the result is obtained by changing to the new variables $\eta = N^{-1}\xi, y = \beta Nx$, and then applying Proposition 2.11 and the Plancherel identity. The cases of $\chi_{\leq 1}, \chi'_{\leq 1}, \chi'_{\geq 1}$ and $\chi''_{\geq 1}$ can be treated in the same way. Since $\chi_{\geq 1}(x \cdot \xi) = 1 - \chi_{\leq 1}(x \cdot \xi)$, combining with the Plancherel identity, the analogous estimate when $\chi_{\leq 1}(x \cdot \xi)$ is replaced by $\chi_{\geq 1}(x \cdot \xi)$ in (2.5) is also proved. \square

3. THE INCOMING/OUTGOING WAVES

3.1. Definitions of the incoming/outgoing waves. First of all, we give the definitions of the incoming/outgoing waves for the Schrödinger flow.

3.1.1. The deformed Fourier transform. We denote the standard Fourier transform by \hat{f} or $\mathcal{F}f$ as

$$(\mathcal{F}f(\xi) \text{ or })\hat{f}(\xi) = \int_{\mathbb{R}^d} e^{-2\pi i x \cdot \xi} f(x) dx,$$

and its inverse transform

$$(\mathcal{F}^{-1}f(x) \text{ or })\check{f}(x) = \int_{\mathbb{R}^d} e^{2\pi i x \cdot \xi} f(\xi) d\xi.$$

Now we give a deformed Fourier transform, and its basic properties.

Definition 3.1. *Let $\alpha \in \mathbb{R}, \beta \in \mathbb{R}$, and let $f \in \mathcal{S}(\mathbb{R}^d)$ with $|x|^\beta f \in L^1_{loc}(\mathbb{R}^d)$. We define*

$$\mathcal{F}f(\xi) = |\xi|^\alpha \int_{\mathbb{R}^d} e^{-2\pi i x \cdot \xi} |x|^\beta f(x) dx. \quad (3.1)$$

Then it is easy to see the following inverse transform, that is,

Lemma 3.2. *Let $f \in \mathcal{S}(\mathbb{R}^d)$, $|x|^\beta f \in L^1_{loc}(\mathbb{R}^d)$ and $|\xi|^{-\alpha} \mathcal{F}f \in L^1_{loc}(\mathbb{R}^d)$, then for any $x \in \mathbb{R}^d \setminus \{0\}$,*

$$f(x) = |x|^{-\beta} \int_{\mathbb{R}^d} e^{2\pi i x \cdot \xi} |\xi|^{-\alpha} \mathcal{F}f(\xi) d\xi. \quad (3.2)$$

Proof. From the definition,

$$|\xi|^{-\alpha} \mathcal{F}f(\xi) = \mathcal{F}(|x|^\beta f)(\xi). \quad (3.3)$$

Hence, by the inverse Fourier transform, we have

$$|x|^\beta f(x) = \mathcal{F}^{-1}(|\xi|^{-\alpha} \mathcal{F}f)(x).$$

This gives the formula in (3.2). \square

We give the following remark on the conditions in Lemma 3.2.

Remark 3.3. If $f \in \mathcal{S}(\mathbb{R}^d)$ is radial, and $\beta > -d$, then $|x|^\beta f \in L_{loc}^1(\mathbb{R}^d)$. Indeed,

$$\int_{|x| \leq 1} |x|^\beta f(x) dx = c(d) \int_0^1 r^{\beta+d-1} f(r) dr \lesssim \|f\|_{L^\infty(\mathbb{R}^d)}.$$

Similarly, if $\alpha < d$, we can prove that $|\xi|^{-\alpha} \mathcal{F}f \in L_{loc}^1(\mathbb{R}^d)$.

We now give the radial version of the deformed Fourier transform and its inverse transform. We note that if f is radial, so is $\mathcal{F}f$. Moreover,

$$\begin{aligned} \mathcal{F}f(\rho) &= \rho^\alpha \int_0^{+\infty} \int_{|\theta|=1} e^{-2\pi i \rho r \omega \cdot \theta} d\theta f(r) r^{\beta+d-1} dr \\ &= \rho^\alpha \int_0^{+\infty} \widehat{d\omega}(\rho r \omega) r^{\beta+d-1} f(r) dr, \end{aligned}$$

where $\xi = \rho\omega$. Note that $\widehat{d\omega}$ has radial symmetry, and

$$\widehat{d\omega}(\rho r \omega) = \widehat{d\omega}(\rho r) = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho r \sin \theta} \cos^{d-2} \theta d\theta.$$

(It is equal to $2\pi(\rho r)^{-\frac{d-2}{2}} J_{\frac{d-2}{2}}(2\pi \rho r)$, where $J_{\frac{d-2}{2}}$ is a bessel function.) Therefore,

$$\mathcal{F}f(\rho) = \rho^\alpha \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho r \sin \theta} \cos^{d-2} \theta r^{\beta+d-1} f(r) d\theta dr. \quad (3.4)$$

Similarly, we have

$$f(r) = r^{-\beta} \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \cos^{d-2} \theta \rho^{-\alpha+d-1} \mathcal{F}f(\rho) d\theta d\rho. \quad (3.5)$$

3.1.2. Definitions. In this subsection, we define the incoming and outgoing components of functions and present their properties. Our definition here is inspired by T. Tao [44]. For convenience, we denote

$$J(r) = \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \cos^{d-2} \theta d\theta. \quad (3.6)$$

Then we have

$$J(-r) = \int_{-\frac{\pi}{2}}^0 e^{2\pi i r \sin \theta} \cos^{d-2} \theta d\theta.$$

Let

$$K(r) = \chi_{\geq 2}(r) \left[-\frac{1}{2\pi i r} + \frac{d-3}{(2\pi i r)^3} \right], \quad \text{for } d = 3, 4, 5.$$

Definition 3.4. Let $\alpha < d, \beta > -d$, and the function $f \in L_{loc}^1(\mathbb{R}^d)$ be radial. We define the incoming component of f as

$$f_{in}(r) = r^{-\beta} \int_0^{+\infty} \left(J(-\rho r) + K(\rho r) \right) \rho^{-\alpha+d-1} \mathcal{F}f(\rho) d\rho;$$

the outgoing component of f as

$$f_{out}(r) = r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{-\alpha+d-1} \mathcal{F}f(\rho) d\rho.$$

Moreover, for any fixed integer k , we define the frequency restricted incoming component of f as

$$f_{in,k}(r) = r^{-\beta} \int_0^{+\infty} \left(J(-\rho r) + K(\rho r) \right) \chi_{2^k}(\rho) \rho^{-\alpha+d-1} \mathcal{F}f(\rho) d\rho;$$

correspondingly, the frequency restricted outgoing component of f as

$$f_{out,k}(r) = r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{-\alpha+d-1} \mathcal{F}f(\rho) d\rho.$$

Remark 3.5. If we consider the higher dimensional version, then $K(r)$ should be modified. Here we use an appropriately modified Fourier transform in order to cancel the singularity from r^{-1} at origin and meanwhile to guarantee the boundedness of the incoming/outgoing projection operators on L^2 .

In the whole of the present paper, we set the numbers in Definition 3.4 that

$$\beta = \frac{d-1}{2} - 2, \quad \text{and} \quad \alpha = 0.$$

(These numbers should be changed if one considers the cases when $d \neq 3, 4, 5$.)

From the definitions and (3.5), we have

$$f(r) = f_{out}(r) + f_{in}(r).$$

Moreover,

$$f_{out/in}(r) = \sum_{k=-\infty}^{+\infty} f_{out/in,k}(r).$$

Correspondingly, for $k_0 \in \mathbb{Z}$, we denote

$$f_{out/in,\geq k_0}(r) = \sum_{k=k_0}^{+\infty} f_{out/in,k}(r).$$

3.2. Basic properties of the incoming/outgoing functions. First, we give the estimates on the following oscillatory integrals. The first two can be regarded as some asymptotic behaviors of the restricted forms of $J(r)$.

Lemma 3.6. *Let $d = 3, 4, 5$. There exists constant $c \in \mathbb{C}$ only depending on d , such that when $r \rightarrow +\infty$,*

$$\int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta = cr^{-\frac{d-1}{2}} e^{2\pi i r} + O(r^{-\frac{d+1}{2}}); \quad (3.7)$$

and

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta = r^{-\frac{d-1}{2}} \left(\bar{c} e^{-2\pi i r} + c e^{2\pi i r} \right) + O(r^{-\frac{d+1}{2}}). \quad (3.8)$$

Remark 3.7. The estimates presented in this lemma are not sharp. For example, the first estimate can be improved as the form of $r^{-\frac{d-1}{2}} (p_K(1/r) e^{2\pi i r} + O(r^{-K-1}))$ for any $K \in \mathbb{Z}^+$, where p_K is some polynomial function of order K . However, the version presented in this lemma is simpler and enough in this paper.

Proof of Lemma 3.6. We only prove the first estimate, since the second one is the sum of

$$\int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \quad \text{and} \quad \int_0^{\frac{\pi}{2}} e^{-2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta.$$

When $d = 3$,

$$\begin{aligned} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos \theta d\theta &= \int_0^1 e^{2\pi i r s} \chi_{\geq \frac{1}{2}}(s) ds \\ &= \frac{1}{2\pi i r} \int_0^1 \partial_s (e^{2\pi i r s}) \chi_{\geq \frac{1}{2}}(s) ds = \frac{1}{2\pi i r} e^{2\pi i r} - \frac{1}{2\pi i r} \int_0^1 e^{2\pi i r s} \chi'_{\geq \frac{1}{2}}(s) ds. \end{aligned}$$

By integration-by-parts K times, we have

$$\int_0^1 e^{2\pi i r s} \chi'_{\geq \frac{1}{2}}(s) ds = O(r^{-K}).$$

Hence we obtain that

$$\int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos \theta d\theta = \frac{1}{2\pi i r} e^{2\pi i r} + O(r^{-K}).$$

When $d = 4$, using the formula,

$$2\pi i r \cos \theta \cdot e^{2\pi i r \sin \theta} = \partial_\theta e^{2\pi i r \sin \theta}, \quad (3.9)$$

we have

$$\begin{aligned} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^2 \theta d\theta &= \frac{1}{2\pi i r} \int_0^{\frac{\pi}{2}} \partial_\theta e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos \theta d\theta \\ &= \frac{1}{2\pi i r} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \left(\chi_{\geq \frac{\pi}{6}}(\theta) \sin \theta - \chi'_{\geq \frac{\pi}{6}}(\theta) \cos \theta \right) d\theta. \end{aligned}$$

Note that

$$\int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi'_{\geq \frac{\pi}{6}}(\theta) \cos \theta d\theta = \int_0^1 e^{2\pi i r s} \chi'_{\geq \frac{1}{2}}(s) ds = O(r^{-K}).$$

Moreover, using Lemma 2.10,

$$\begin{aligned} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \sin \theta d\theta &= \frac{1}{2} \int_0^\pi e^{2\pi i r \sin \theta} \chi_{\frac{\pi}{6} \leq \cdot \leq \frac{5\pi}{6}}(\theta) \sin \theta d\theta \\ &= \left(\frac{1}{2\pi i r} \right)^{\frac{1}{2}} e^{2\pi i r} + O(r^{-\frac{3}{2}}). \end{aligned}$$

This gives the required estimate in dimension four.

When $d = 5$, we have

$$\begin{aligned} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^3 \theta d\theta &= \int_0^1 e^{2\pi i r s} \chi_{\geq \frac{1}{2}}(s) (1 - s^2) ds \\ &= \frac{1}{2\pi i r} \int_0^1 \partial_s (e^{2\pi i r s}) \chi_{\geq \frac{1}{2}}(s) (1 - s^2) ds. \end{aligned}$$

By integration-by-parts once and noting the zero boundary values, it is equal to

$$\frac{1}{2\pi i r} \int_0^1 e^{2\pi i r s} \left(-\chi'_{\geq \frac{1}{2}}(s) (1 - s^2) + 2\chi_{\geq \frac{1}{2}}(s) s \right) ds.$$

Now arguing similarly as above and integration-by-parts many times, we have

$$\frac{1}{2\pi ir} \int_0^1 e^{2\pi ir s} \chi'_{\geq \frac{1}{2}}(s) (1-s^2) ds = O(r^{-K}),$$

and

$$\frac{1}{2\pi ir} \int_0^1 e^{2\pi ir s} \chi_{\geq \frac{1}{2}}(s) s ds = \frac{1}{(2\pi ir)^2} \left(1 - \frac{1}{2\pi ir}\right) e^{2\pi ir} + O(r^{-K}).$$

This obeys the form as claimed. Hence we finish the proof of the lemma. \square

Second, we have the following estimates.

Lemma 3.8. *When $r \rightarrow +\infty$,*

$$\int_0^{\frac{\pi}{2}} e^{2\pi ir \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta = -\frac{1}{2\pi ir} + \frac{d-3}{(2\pi ir)^3} + O(r^{-5}).$$

Moreover,

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{2\pi ir \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta = O(r^{-10}).$$

Proof. Using the formula (3.9) and integration-by-parts,

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} e^{2\pi ir \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &= \frac{1}{2\pi ir} \int_0^{\frac{\pi}{2}} \partial_\theta e^{2\pi ir \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-3} \theta d\theta \\ &= -\frac{1}{2\pi ir} - \frac{1}{2\pi ir} \int_0^{\frac{\pi}{2}} e^{2\pi ir \sin \theta} \eta_1(\theta) d\theta, \end{aligned}$$

where

$$\eta_1(\theta) = \partial_\theta \left(\chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-3} \theta \right).$$

Note that $\eta_1(0) = \eta_1(\frac{\pi}{2}) = 0$, then using the formula (3.9) and integration-by-parts again, we have

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} e^{2\pi ir \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &= -\frac{1}{2\pi ir} + \frac{1}{(2\pi ir)^2} \int_0^{\frac{\pi}{2}} e^{2\pi ir \sin \theta} \eta_2(\theta) d\theta, \end{aligned}$$

where

$$\eta_2(\theta) = \partial_\theta \left(\frac{\eta_1(\theta)}{\cos \theta} \right).$$

Note that $|\eta_2(\theta)| \lesssim 1$. Moreover, by a direct computation, we have

$$\eta_2(0) = -(d-3), \quad \eta_2\left(\frac{\pi}{2}\right) = 0.$$

Then repeating the process above again, we obtain

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &= -\frac{1}{2\pi i r} + \frac{d-3}{(2\pi i r)^3} + \frac{1}{(2\pi i r)^3} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \eta_3(\theta) d\theta. \end{aligned}$$

Here

$$\eta_3(\theta) = \partial_\theta \left(\frac{\eta_2(\theta)}{\cos \theta} \right).$$

Moreover, we note that $\eta_3(0) = \eta_3(\frac{\pi}{2}) = 0$. Then the same process gives

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &= -\frac{1}{2\pi i r} + \frac{d-3}{(2\pi i r)^3} + \frac{1}{(2\pi i r)^5} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \eta_4(\theta) d\theta, \end{aligned} \quad (3.10)$$

where the function η_4 satisfies $|\eta_4(\theta)| \lesssim 1$. Then we obtain the desired result in the first estimate.

The second one, because of the zero boundary values in every step, can be obtained by integration-by-parts 10 times. \square

As consequences of the lemmas above, we have the following variant forms of $J(r) - K(r)$. The first one is

Corollary 3.9. *Let $d = 3, 4, 5$, then*

$$\chi_{\leq 1}(r) \left(J(r) - K(r) \right) = \chi_{\leq 1}(r) \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \cos^{d-2} \theta d\theta; \quad (3.11)$$

and

$$\begin{aligned} \chi_{\geq 1}(r) \left(J(r) - K(r) \right) &= \chi_{\geq 1}(r) \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &+ \chi_{1 \leq \cdot \leq 2}(r) \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta + \chi_{\geq 2}(r) \frac{1}{(2\pi i r)^5} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \tilde{\eta}_d(\theta) d\theta, \end{aligned} \quad (3.12)$$

where $\tilde{\eta}_d, d = 3, 4, 5$ are uniformly bounded functions.

Proof. From the definition of $K(r)$, we have $K(r) = 0$ in the part of $\chi_{\leq 1}(r)$. Hence, by the definition of $J(r)$ directly, we have (3.11).

For (3.12), we split $J(r)$ into the following two parts,

$$\int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \quad \text{and} \quad \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta.$$

For the latter part, we use the identity (3.10) to obtain that

$$\chi_{\geq 2}(r) \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta = K(r) + \chi_{\geq 2}(r) \frac{1}{(2\pi i r)^5} \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \eta_4(\theta) d\theta.$$

Hence, setting $\tilde{\eta}_d = \eta_4$, we get (3.12). Note that from the definitions in the proof of Lemma 3.8, η_4 are uniformly bounded functions whenever $d = 3, 4, 5$. This proves the lemma. \square

The following result shows the asymptotic behaviors of $J(r) - K(r)$ and $J(r) + J(-r)$.

Corollary 3.10. *Let $d = 3, 4, 5$, then*

$$J(r) - K(r) = \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta + O(\langle r \rangle^{-5}) \quad (3.13)$$

$$= O(\langle r \rangle^{-\frac{d-1}{2}}); \quad (3.14)$$

and

$$J(r) + J(-r) = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta + O(\langle r \rangle^{-10}). \quad (3.15)$$

Proof. Note that when $r \lesssim 1$, the following four functions: $J(r) - K(r)$, $J(r) + J(-r)$,

$$\int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \quad \text{and} \quad \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta$$

are uniformly bounded. Hence, the estimates of (3.13) and (3.15) hold when $r \lesssim 1$.

When $r \gtrsim 1$, from (3.12),

$$J(r) - K(r) = \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta + O(r^{-5});$$

and from Lemma 3.8,

$$J(r) + J(-r) = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta + O(r^{-10}).$$

Hence, we obtain (3.13) and (3.15). Combining with (3.13) and (3.7), we obtain (3.14). This finishes the proof of the corollary. \square

3.2.1. Matching estimates in frequency and physical spaces. In this subsection, we will show that the incoming/outgoing projection would almost preserve the localization in both space and frequency. The first result below is related to the preservation of frequency. Specifically, if a function f has high frequency $f = P_{2^k} f$, then its incoming/outgoing component will have almost the same frequency plus a smooth perturbation.

Proposition 3.11. *Let $k_0 \geq 0$. Suppose that $f \in L^2(\mathbb{R}^d)$ with $\text{supp } f \subset \{x : |x| \geq 1\}$, then*

$$(P_{\geq 2^{k_0}} f)_{\text{out/in}} = (P_{\geq 2^{k_0}} f)_{\text{out/in}, \geq k_0-1} + h$$

with

$$\|h\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-k_0} \|f\|_{H^{-10}(\mathbb{R}^d)}. \quad (3.16)$$

Here $\mu(d) = 2$ if $d = 3, 4$, and $\mu(5) = 3$. Similarly,

$$(P_{\leq 2^{k_0}} f)_{\text{out/in}} = (P_{\leq 2^{k_0}} f)_{\text{out/in}, \leq k_0+1} + \tilde{h}$$

with \tilde{h} verifying the same estimate to (3.16).

Proof. We only give the estimate of the first part, since the second one can be treated in the same manner. Moreover, considering the support of f , for simplifying the notations, we write

$$f = \chi_{\geq 1} f.$$

To give the desired estimate, we use the Littlewood-Paley decomposition and write

$$P_{\geq 2^{k_0}} f = \sum_{k=k_0}^{+\infty} P_{2^k} f.$$

From (3.3), we note that

$$\mathcal{F}(P_{2^k} f)(\rho) = \mathcal{F}\left(|x|^\beta P_{2^k} f\right)(\rho).$$

Accordingly, we decompose $\mathcal{F}(P_{2^k} f)(\rho)$ into the following two parts,

$$\chi_{\geq 2^{k_0-1}}(\rho) \mathcal{F}\left(|x|^\beta P_{2^k} f\right)(\rho) + \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F}\left(|x|^\beta P_{2^k} f\right)(\rho).$$

Now due to the support of f , we further decompose the expression above as

$$\begin{aligned} & \chi_{\geq 2^{k_0-1}}(\rho) \mathcal{F}\left(|x|^\beta P_{2^k} f\right)(\rho) + \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F}\left(\chi_{\geq \frac{1}{2}} |x|^\beta \cdot P_{2^k} \chi_{\geq 1} f\right)(\rho) \\ & + \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F}\left(\chi_{\leq \frac{1}{2}}(x) |x|^\beta \cdot P_{2^k} \chi_{\geq 1} f\right)(\rho). \end{aligned}$$

We denote

$$\begin{aligned} g_1(\rho) &= \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F}\left(\chi_{\geq \frac{1}{2}}(x) |x|^\beta \cdot P_{2^k} \chi_{\geq 1} f\right)(\rho); \\ g_2(\rho) &= \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F}\left(\chi_{\leq \frac{1}{2}}(x) |x|^\beta \cdot P_{2^k} \chi_{\geq 1} f\right)(\rho). \end{aligned}$$

Then we have

$$\mathcal{F}(P_{2^k} f)(\rho) = \chi_{\geq 2^{k_0-1}}(\rho) \mathcal{F}(P_{2^k} f)(\rho) + g_1 + g_2. \quad (3.17)$$

From Definition 3.4, we have

$$(P_{2^k} f)_{out} = r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} \mathcal{F}(P_{2^k} f)(\rho) d\rho,$$

thus using (3.17), we write

$$\begin{aligned} (P_{2^k} f)_{out} &= r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} \chi_{\geq 2^{k_0-1}}(\rho) \mathcal{F}(P_{2^k} f)(\rho) d\rho \\ &+ r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_1(\rho) d\rho \\ &+ r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_2(\rho) d\rho. \end{aligned}$$

Denote

$$\begin{aligned} h_k(r) &= r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_1(\rho) d\rho \\ &+ r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_2(\rho) d\rho, \end{aligned} \quad (3.18)$$

then

$$(P_{2^k} f)_{out} = (P_{2^k} f)_{out, \geq k_0-1} + h_k.$$

Moreover, let

$$h(r) = \sum_{k=k_0}^{+\infty} h_k(r),$$

then

$$(P_{\geq 2^{k_0}} f)_{out/in} = (P_{\geq 2^{k_0}} f)_{out/in, \geq k_0-1} + h.$$

Now we give the control of h_k . To this end, we first claim that for any $M \in \mathbb{Z}^+$,

$$\|g_1\|_{L_\rho^\infty} + \|g_2\|_{L_\rho^\infty} \lesssim_M 2^{-Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \quad (3.19)$$

To prove this claim, we note that

$$g_1(\rho) = \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F} \left(P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta) \cdot P_{2^k} f \right)(\rho).$$

Here we have used the fact $k \geq k_0$. Then by Hausdorff-Young's inequality,

$$\begin{aligned} \|g_1\|_{L_\rho^\infty} &\lesssim \left\| \mathcal{F} \left(P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta) \cdot P_{2^k} f \right) \right\|_{L_\rho^\infty} \\ &\lesssim \|P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta) \cdot P_{2^k} f\|_{L_x^1(\mathbb{R}^d)} \\ &\lesssim \|P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta)\|_{L_x^2(\mathbb{R}^d)} \|P_{2^k} f\|_{L_x^2(\mathbb{R}^d)}. \end{aligned}$$

Note that

$$\|P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta)\|_{L_x^2(\mathbb{R}^d)} \lesssim 2^{-Mk} \|(-\Delta)^{\frac{M}{2}} P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta)\|_{L_x^2(\mathbb{R}^d)} \lesssim 2^{-Mk},$$

thus we obtain that

$$\|g_1\|_{L_\rho^\infty} \lesssim 2^{-Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

For g_2 , we rewrite it as

$$g_2(\rho) = \chi_{\leq 2^{k_0-1}}(\rho) \mathcal{F} \left(\chi_{\leq \frac{1}{2}}(x) |x|^\beta \cdot \chi_{\leq \frac{3}{4}}(x) P_{2^k} \chi_{\geq 1} f \right)(\rho).$$

Then

$$\begin{aligned} \|g_2\|_{L_\rho^\infty} &\lesssim \left\| \mathcal{F} \left(\chi_{\leq \frac{1}{2}}(x) |x|^\beta \cdot \chi_{\leq \frac{3}{4}}(x) P_{2^k} \chi_{\geq 1} f \right) \right\|_{L_\rho^\infty} \\ &\lesssim \|\chi_{\leq \frac{1}{2}}(x) |x|^\beta \cdot \chi_{\leq \frac{3}{4}}(x) P_{2^k} \chi_{\geq 1} f\|_{L_x^1(\mathbb{R}^d)} \\ &\lesssim \|\chi_{\leq \frac{1}{2}}(x) |x|^\beta\|_{L_x^2(\mathbb{R}^d)} \|\chi_{\leq \frac{3}{4}} P_{2^k} \chi_{\geq 1} f\|_{L_x^2(\mathbb{R}^d)}. \end{aligned}$$

Since $\beta = \frac{d-5}{2} > -\frac{d}{2}$, we have

$$\|\chi_{\leq \frac{1}{2}} |x|^\beta\|_{L_x^2(\mathbb{R}^d)} \lesssim 1.$$

Moreover, by the mismatch estimate given in Lemma 2.4, we have

$$\|\chi_{\leq \frac{3}{4}} P_{2^k} \chi_{\geq 1} f\|_{L_x^2(\mathbb{R}^d)} \lesssim 2^{-Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Hence, we obtain that

$$\|g_2\|_{L_\rho^\infty} \lesssim 2^{-Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

This together with the estimate on g_1 , gives the claim (3.19).

Based on (3.19), we consider $\|h_k\|_{L^2(\mathbb{R}^d)}$. Note that

$$\|h_k\|_{L^2(\mathbb{R}^d)} = c \left\| r^{\frac{d-1}{2}} h_k(r) \right\|_{L_r^2}.$$

Further,

$$\begin{aligned} r^{\frac{d-1}{2}} h_k(r) = & r^2 \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_1(\rho) d\rho \\ & + r^2 \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_2(\rho) d\rho. \end{aligned}$$

We only consider the first term. Since the estimates for g_1, g_2 are the same, see (3.19), they can be treated in the same way. To do this, we split it into the following two parts again,

$$r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \rho^{d-1} g_1(\rho) d\rho, \quad (3.20a)$$

and

$$r^2 \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{\geq 1}(\rho r) \rho^{d-1} g_1(\rho) d\rho. \quad (3.20b)$$

Since $|J(\rho r)| \lesssim 1$, (3.20a) can be controlled by

$$\int_0^{+\infty} \chi_{\leq 1}(\rho r) \rho^{d-3} |g_1(\rho)| d\rho.$$

Since $d \geq 3$,

$$\begin{aligned} \|(3.20a)\|_{L_r^2} & \lesssim \left\| \int_0^{+\infty} \chi_{\leq 1}(\rho r) \rho^{d-3} \chi_{\leq 2^{k_0-1}}(\rho) |g_1(\rho)| d\rho \right\|_{L_r^2} \\ & \lesssim \int_0^{+\infty} \rho^{d-\frac{7}{2}} \chi_{\leq 2^{k_0-1}}(\rho) |g_1(\rho)| d\rho \\ & \lesssim 2^{(d-\frac{5}{2})k_0-Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

Then, choosing M large enough, we have

$$\|(3.20a)\|_{L_r^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \quad (3.21)$$

For (3.20b), according to (3.13), we split it into the following two parts,

$$r^2 \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \rho^{d-1} g_1(\rho) d\rho, \quad (3.22-1)$$

and

$$r^2 \int_0^{+\infty} O(\langle \rho r \rangle^{-5}) \chi_{\geq 1}(\rho r) \rho^{d-1} g_1(\rho) d\rho. \quad (3.22-2)$$

To estimate (3.22-1), we use the formula

$$e^{2\pi i \rho r \sin \theta} = \frac{1}{(2\pi i r \sin \theta)^2} \partial_\rho^2 e^{2\pi i \rho r \sin \theta}$$

to reduce it as

$$\int_0^{+\infty} \int_0^{\frac{\pi}{2}} \frac{1}{(2\pi i \sin \theta)^2} \partial_\rho^2 e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \rho^{d-1} g_1(\rho) d\rho.$$

By integration-by-parts, it is equal to

$$\int_0^{+\infty} \int_0^{\frac{\pi}{2}} \frac{1}{(2\pi i \sin \theta)^2} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \partial_\rho^2 \left[\chi_{\geq 1}(\rho r) \rho^{d-1} g_1(\rho) \right] d\rho.$$

Then by Corollary 2.12, we have

$$\begin{aligned} \|(3.22-1)\|_{L_r^2} &\lesssim \int_0^{\frac{\pi}{2}} \frac{1}{(2\pi \sin \theta)^{\frac{5}{2}}} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &\quad \cdot \left(\|\rho^{d-3} g_1(\rho)\|_{L_\rho^2} + \|\rho^{d-2} \partial_\rho g_1(\rho)\|_{L_\rho^2} + \|\rho^{d-1} \partial_\rho^2 g_1(\rho)\|_{L_\rho^2} \right). \end{aligned}$$

Note that

$$\begin{aligned} \|\rho^{d-3} g_1(\rho)\|_{L_\rho^2} &\lesssim \|\chi_{\leq 2^{k_0-1}}(\rho) \rho^{d-3}\|_{L_\rho^2} \|g_1\|_{L_\rho^\infty} \\ &\lesssim 2^{(d-\frac{5}{2})k_0-Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}; \end{aligned}$$

and furthermore, by the Plancherel identity,

$$\begin{aligned} \|\rho^{d-2} \partial_\rho g_1(\rho)\|_{L_\rho^2} &\lesssim 2^{\frac{d-3}{2}k_0} \left\| \rho^{\frac{d-1}{2}} \chi_{\leq 2^{k_0-1}}(\rho) \nabla_\xi \mathcal{F} \left((\chi_{\geq \frac{1}{2}} |x|^\beta) \cdot P_{2^k} f \right) (\rho) \right\|_{L_\rho^2} \\ &\lesssim 2^{\frac{d-3}{2}k_0} \left\| \chi_{\leq 2^{k_0-1}}(\xi) \nabla \mathcal{F} \left(P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta) \cdot P_{2^k} f \right) (\xi) \right\|_{L_\xi^2(\mathbb{R}^d)} \\ &\lesssim 2^{\frac{d-3}{2}k_0} \|P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta x) \cdot P_{2^k} f\|_{L^2(\mathbb{R}^d)} \\ &\lesssim 2^{\frac{d-3}{2}k_0} \|P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^\beta x)\|_{L^\infty(\mathbb{R}^d)} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)} \\ &\lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

Moreover, using the following estimate instead,

$$\|P_{\geq 2^{k-2}} (\chi_{\geq \frac{1}{2}} |x|^{\beta+2})\|_{L^\infty(\mathbb{R}^d)} \lesssim 2^{-Mk},$$

we also have

$$\|\rho^{d-1} \Delta g_1\|_{L_\rho^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Since $\partial_\rho^2 = \Delta - \frac{d-1}{\rho} \partial_\rho$, by the estimates above, we get

$$\|\rho^{d-1} \partial_\rho^2 g_1(\rho)\|_{L_\rho^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Therefore,

$$\|(3.22-1)\|_{L_r^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

For (3.22-2), it is bounded by

$$r^{-1} \int_0^{+\infty} \chi_{\geq 1}(\rho r) \rho^{d-4} |g_1(\rho)| d\rho.$$

Similar as (3.20a),

$$\|(3.22-2)\|_{L_r^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Combining the above two estimates for (3.22), we obtain

$$\|(3.20b)\|_{L_r^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Then collecting the above two findings for (3.20), we have

$$\|h_k\|_{L^2(\mathbb{R}^d)} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Now we consider the estimates of h_k with the high-order derivatives. For $\|\Delta h_k\|_{L^2(\mathbb{R}^d)}$, it is equivalent to

$$\left\| r^{\frac{d-1}{2}} \partial_{rr} h_k \right\|_{L_r^2} + \left\| r^{\frac{d-1}{2}-1} \partial_r h_k \right\|_{L_r^2}.$$

Note that from the definition of h_k in (3.18), both of $r^{\frac{d-1}{2}} \partial_{rr} h_k$ and $r^{\frac{d-1}{2}-1} \partial_r h_k$ are the combination of the following three parts,

$$\begin{aligned} & \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho; \\ & r \int_0^{+\infty} \partial_r \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho; \end{aligned} \quad (3.23)$$

and

$$r^2 \int_0^{+\infty} \partial_r^2 \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho,$$

where $j = 1, 2$. When $d = 5$, we also need to estimate $\|\nabla \Delta h_k\|_{L^2(\mathbb{R}^d)}$, which is equivalent to

$$\left\| r^{\frac{d-1}{2}} \partial_{rrr} h_k \right\|_{L_r^2} + \left\| r^{\frac{d-1}{2}-1} \partial_{rr} h_k \right\|_{L_r^2} + \left\| r^{\frac{d-1}{2}-2} \partial_r h_k \right\|_{L_r^2}.$$

Note that $\beta = 0$ in this case, hence all of $r^{\frac{d-1}{2}} \partial_{rrr} h_k$, $r^{\frac{d-1}{2}-1} \partial_{rr} h_k$ and $r^{\frac{d-1}{2}-2} \partial_r h_k$ are the combination of the following three parts,

$$\begin{aligned} & \int_0^{+\infty} \partial_r \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho; \\ & r \int_0^{+\infty} \partial_r^2 \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho; \end{aligned}$$

and

$$r^2 \int_0^{+\infty} \partial_r^3 \left(J(\rho r) - K(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho.$$

(Note that there is no singularity in r , due to $\beta = 0$.)

By the formulas given in Corollary 3.9, we have the explicit form of $\partial_r^l \left(J(\rho r) - K(\rho r) \right)$ for $l = 1, 2, 3$, then similar argument can be used to estimate all the terms which can be dominated by $2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}$. For the sake of completeness, we take the term (3.23) for example and give the estimation. To do this, we split (3.23) into the following two parts,

$$r \int_0^{+\infty} \partial_r \left((J(\rho r) - K(\rho r)) \chi_{\leq 1}(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho; \quad (3.24a)$$

and

$$r \int_0^{+\infty} \partial_r \left((J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \right) \rho^{d-1} g_j(\rho) d\rho. \quad (3.24b)$$

For (3.24a), we use the following estimate which is from (3.11),

$$\left| \partial_r \left((J(\rho r) - K(\rho r)) \chi_{\leq 1}(\rho r) \right) \right| \lesssim \rho \chi_{\leq 1}(\rho r).$$

Then (3.24a) can be dominated by

$$\int_0^{+\infty} \chi_{\leq 1}(\rho r) \rho^{d-1} |g_j(\rho)| d\rho.$$

Hence,

$$\begin{aligned} \|(3.24a)\|_{L_r^2} &\lesssim \int_0^{+\infty} \rho^{d-\frac{3}{2}} |g_j(\rho)| d\rho \\ &\lesssim 2^{(d-\frac{1}{2})k_0} \|g_j\|_{L_\rho^\infty} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

For (3.24b), we use the following estimate which is from (3.12),

$$\begin{aligned} \partial_r \left(\chi_{\geq 1}(\rho r) (J(\rho r) - K(\rho r)) \right) &= 2\pi i \rho \chi_{\geq 1}(\rho r) \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \sin \theta \cos^{d-2} \theta d\theta \\ &\quad + \chi_{\geq 1}(\rho r) \cdot O(\rho^{-4} r^{-5}). \end{aligned}$$

Accordingly, (3.24b) can be split into the following two subparts again,

$$2\pi i r \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \sin \theta \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \rho^d g_j(\rho) d\rho; \quad (3.25-1)$$

and

$$r \int_0^{+\infty} O(\rho^{-4} r^{-5}) \chi_{\geq 1}(\rho r) \rho^{d-1} g_j(\rho) d\rho. \quad (3.25-2)$$

Then the part (3.25-1) can be treated similarly as (3.22-1); the part (3.25-2) can be treated similarly as (3.22-2), and thus we have

$$\|(3.24b)\|_{L_r^2} \leq \|(3.25-1)\|_{L_r^2} + \|(3.25-2)\|_{L_r^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Therefore, from the estimates on (3.24), we get

$$\|(3.23)\|_{L_r^2} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Combining the estimates above, we establish that

$$\|h_k\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-\frac{1}{2}Mk} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Therefore, by summation in k and choosing M suitably large, we obtain that

$$\|h\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-k_0} \|f\|_{H^{-10}(\mathbb{R}^d)}.$$

This finishes the proof of the proposition. \square

The following result shows that if f is supported outside of a ball, then $f_{out/in}$ is also almost supported outside of the ball.

Lemma 3.12. *Let $\mu(d)$ be defined in Proposition 3.11. Suppose that $\text{supp} f \subset \{x : |x| > 1\}$, then*

$$\|\chi_{\leq \frac{1}{4}}(P_{\geq 1} f)_{out/in}\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim \|f\|_{H^{-1}(\mathbb{R}^d)},$$

and for any $k \in \mathbb{Z}^+$,

$$\|\chi_{\leq \frac{1}{4}}(P_{2^k} f)_{out/in, \geq k-1}\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-2k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Proof. We only consider the estimates on the outgoing part f_{out} , because the ones on the incoming part can be proved in a similar way. Using the Littlewood-Paley decomposition, we write

$$P_{\geq 1} f = \sum_{k=0}^{+\infty} P_{2^k} f.$$

Then

$$\|\chi_{\leq \frac{1}{4}}(P_{\geq 1}f)_{out/in}\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim \sum_{k=0}^{+\infty} \|\chi_{\leq \frac{1}{4}}(P_{2^k}f)_{out}\|_{H^{\mu(d)}(\mathbb{R}^d)}.$$

Using Proposition 3.11, we have

$$(P_{2^k}f)_{out} = (P_{2^k}f)_{out, \geq k-1} + h, \quad \text{with} \quad \|h\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-k} \|P_{2^k}f\|_{H^{-10}(\mathbb{R}^d)}.$$

Hence,

$$\|\chi_{\leq \frac{1}{4}}(P_{2^k}f)_{out}\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim \|\chi_{\leq \frac{1}{4}}(P_{2^k}f)_{out, \geq k-1}\|_{H^{\mu(d)}(\mathbb{R}^d)} + \|P_{2^k}f\|_{H^{-10}(\mathbb{R}^d)}.$$

Therefore, we only need to consider $\|\chi_{\leq \frac{1}{4}}(P_{2^k}f)_{out, \geq k-1}\|_{H^{\mu(d)}(\mathbb{R}^d)}$. We first consider the L^2 -norm, which is equal to

$$\|r^{\frac{d-1}{2}} \chi_{\leq \frac{1}{4}}(r)(P_{2^k}f)_{out, \geq k-1}\|_{L_r^2}.$$

From the definition, we have

$$r^{\frac{d-1}{2}}(P_{2^k}f)_{out, \geq k-1} = r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}f)(\rho) d\rho.$$

As in the proof of the previous proposition, we split it into the following two parts,

$$r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}f)(\rho) d\rho; \quad (3.26a)$$

and

$$r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}f)(\rho) d\rho. \quad (3.26b)$$

For (3.26a), using the definition of $\mathcal{F}(P_{2^k}f)$, we rewrite it as

$$r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \int_0^{+\infty} (J(\rho s) + J(-\rho s)) s^{\beta+d-1} P_{2^k}f(s) ds d\rho.$$

Using (3.15) and (3.6), we split (3.26a) into the following two subparts again,

$$\begin{aligned} & r^2 \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \cos^{d-2} \theta d\theta \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' s^{\beta+d-1} P_{2^k}f(s) ds d\rho, \end{aligned} \quad (3.27-1)$$

and

$$r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} P_{2^k}f(s) ds d\rho. \quad (3.27-2)$$

The term (3.27-1) can be rewritten as

$$\begin{aligned} & r^2 \int_0^{\frac{\pi}{2}} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{+\infty} \int_0^{+\infty} e^{2\pi i \rho(r \sin \theta - s \sin \theta')} \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} s^{\beta+d-1} P_{2^k}f(s) ds d\rho \\ & \cdot \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta \cos^{d-2} \theta' d\theta' d\theta. \end{aligned}$$

When $r \leq \frac{1}{3}$, $s \geq \frac{9}{10}$, $|\sin \theta'| \geq \frac{2}{5}$, we have

$$|r \sin \theta - s \sin \theta'| \gtrsim r + s. \quad (3.28)$$

Then using the formula,

$$e^{2\pi i \rho(r \sin \theta - s \sin \theta')} = \frac{1}{2\pi i(r \sin \theta - s \sin \theta')} \partial_\rho \left(e^{2\pi i \rho(r \sin \theta - s \sin \theta')} \right), \quad (3.29)$$

and integration-by-parts 5 times, (3.27-1) turns to

$$\begin{aligned} & r^2 \int_0^{\frac{\pi}{2}} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{+\infty} \int_0^{+\infty} \frac{e^{2\pi i \rho(r \sin \theta - s \sin \theta')}}{[2\pi i(r \sin \theta - s \sin \theta')]^5} \partial_\rho^5 \left[\chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \right] \\ & \cdot s^{\beta+d-1} P_{2^k} f(s) d\rho ds \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta \cos^{d-2} \theta' d\theta' d\theta. \end{aligned} \quad (3.30)$$

Note that $r\chi'_{\leq 1}(\rho r) \lesssim \rho^{-1}$, we have that

$$\left| \partial_\rho^5 \left[\chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \right] \right| \lesssim \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-6}.$$

Here and in the following, for simplicity, we regard $\chi_{\geq 1}$ and its derivatives as the same. Then this last estimate combining with (3.28) and (3.30), gives the bound of (3.27-1) as

$$\int_0^{+\infty} \int_0^{+\infty} \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-8} s^{\beta+d-6} |P_{2^k} f(s)| d\rho ds.$$

Therefore,

$$\|(3.27-1)\|_{L_r^2(\{r \leq \frac{1}{4}\})} \lesssim 2^{(d-\frac{15}{2})k} \|s^{\frac{d-1}{2}} P_{2^k} f\|_{L_s^2} \lesssim 2^{(d-7)k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Note that $|J(\rho r)| \lesssim 1$, thus (3.27-2) is also bounded by

$$\int_0^{+\infty} \int_0^{+\infty} \chi_{\leq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \langle \rho s \rangle^{-10} \rho^{d-3} s^{\beta+d-1} |P_{2^k} f(s)| d\rho ds.$$

Hence, we get

$$\|(3.27-2)\|_{L_r^2(\{r \leq \frac{1}{4}\})} \lesssim 2^{(d-12)k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Combining these two estimates on (3.27), we obtain that

$$\|(3.26a)\|_{L_r^2(\{r \leq \frac{1}{4}\})} \lesssim 2^{-2k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \quad (3.31)$$

Now we consider (3.26b), which is equal to

$$\begin{aligned} & r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} (J(\rho s) + J(-\rho s)) s^{\beta+d-1} P_{2^k} f(s) ds d\rho. \end{aligned}$$

Using (3.15) again, we split (3.26b) into the following two subparts again,

$$\begin{aligned} & r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' s^{\beta+d-1} P_{2^k} f(s) ds d\rho, \end{aligned} \quad (3.32-1)$$

and

$$r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \rho^{d-1} \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} P_{2^k} f(s) ds d\rho. \quad (3.32-2)$$

For (3.32-1), using the formula in (3.12), we can write

$$\chi_{\geq 1}(\rho r) \left(J(\rho r) - K(\rho r) \right) = \chi_{\geq 1}(\rho r) \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \eta(\theta, \rho r) d\theta, \quad (3.33)$$

where the function $\eta(\theta, \rho r)$ is defined as

$$\eta(\theta, \rho r) = \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta + \chi_{1 \leq \cdot \leq 2}(\rho r) \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta + \frac{\chi_{\geq 2}(\rho r)}{(2\pi i \rho r)^5} \tilde{\eta}_d(\theta).$$

Then from (3.33), the term (3.32-1) can be rewritten as

$$\begin{aligned} r^2 \int_0^{\frac{\pi}{2}} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{+\infty} \int_0^{+\infty} e^{2\pi i \rho(r \sin \theta - s \sin \theta')} \chi_{\geq 1}(\rho r) \chi_{\geq 2^{k-1}}(\rho) \eta(\theta, \rho r) \\ \cdot \rho^{d-1} s^{\beta+d-1} P_{2^k} f(s) ds d\rho \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' d\theta. \end{aligned}$$

Now using (3.28) and treating similarly as (3.27-1), we obtain that

$$\|(3.32-1)\|_{L_r^2(\{r \leq \frac{1}{4}\})} \lesssim 2^{(d-12)k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

For (3.32-2), applying (3.14), we have that

$$\chi_{\geq 1}(\rho r) \left(J(\rho r) - K(\rho r) \right) = O((\rho r)^{-\frac{d-1}{2}}). \quad (3.34)$$

Hence,

$$|(3.32-2)| \lesssim r^{2-\frac{d-1}{2}} \int_0^{+\infty} \int_0^{+\infty} \chi_{\geq 2^{k-1}}(\rho) \langle \rho s \rangle^{-10} \rho^{\frac{d-1}{2}} s^{\beta+d-1} |P_{2^k} f(s)| d\rho ds.$$

This gives that

$$\|(3.32-2)\|_{L_r^2(\{r \leq \frac{1}{4}\})} \lesssim 2^{\frac{d-19}{2}k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Therefore, we obtain that

$$\|(3.26b)\|_{L_r^2(\{r \leq \frac{1}{4}\})} \lesssim 2^{\frac{d-19}{2}k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}.$$

Combining the estimates above, we obtain that

$$\|\chi_{\leq \frac{1}{4}}(P_{2^k} f)_{out, \geq k-1}\|_{L^2(\mathbb{R}^d)} \lesssim 2^{-2k} \|P_{2^k} f\|_{L^2(\mathbb{R}^d)}. \quad (3.35)$$

Now similar argument (as in the proof of Proposition 3.11) can be used to treat the functions in the high-order derivatives. Thus we obtain the desired estimate in $H^{\mu(d)}(\mathbb{R}^d)$ space, and prove the lemma. \square

3.3. Boundedness of the incoming/outgoing projection operators. The main results in this section are the boundedness of incoming/outgoing projection operator in $L^2(\mathbb{R}^d)$.

Proposition 3.13. *Suppose that $f \in L^2(\mathbb{R}^d)$, then for any $k \in \mathbb{Z}^+$,*

$$\|f_{out/in, k}\|_{L^2(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

Here the implicit constant is independent of k .

Remark 3.14. We remark here that we have a slightly stronger estimate read as

$$\|f_{out/in}\|_{L^2(\mathbb{R}^d)} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

The proof follows from the similar argument as in the proof of Proposition 3.13 below, with a further spatial cut-off on the function f . However, the log-loss is not essential in this paper, so to simplify the proof we only present a slightly weaker estimate as in Proposition 3.13.

Proof of Proposition 3.13. We only consider the estimates on $f_{out,k}$. Since $f_{out,k}$ is radial, we have

$$\|f_{out,k}\|_{L^2(\mathbb{R}^d)} = c \|r^{\frac{d-1}{2}} f_{out,k}(r)\|_{L_r^2}.$$

By Definition 3.4, we write

$$\begin{aligned} r^{\frac{d-1}{2}} f_{out,k}(r) &= r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}f(\rho) d\rho \\ &= r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}f(\rho) d\rho \end{aligned} \quad (3.36a)$$

$$+ r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}f(\rho) d\rho. \quad (3.36b)$$

Estimates on (3.36a). Note that

$$(3.36a) = r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} (J(\rho s) + J(-\rho s)) s^{\beta+d-1} f(s) ds d\rho.$$

Accordingly, we further split it into the following two subparts again,

$$r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} (J(\rho s) + J(-\rho s)) \chi_{\leq 10}(\rho s) s^{\beta+d-1} f(s) ds d\rho; \quad (3.37-1)$$

and

$$r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} (J(\rho s) + J(-\rho s)) \chi_{\geq 10}(\rho s) s^{\beta+d-1} f(s) ds d\rho. \quad (3.37-2)$$

For (3.37-1), noting that $|J(\rho r)|, |J(\rho s)| \lesssim 1$, we have that

$$|(3.37-1)| \lesssim r^2 \int_0^{+\infty} \int_0^{+\infty} \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \chi_{\leq 10}(\rho s) s^{\beta+d-1} |f(s)| ds d\rho.$$

Thus,

$$\begin{aligned} \|(3.37-1)\|_{L_r^2} &\lesssim \int_0^{+\infty} \int_0^{+\infty} \|r^2 \chi_{\leq 1}(\rho r)\|_{L_r^2} \chi_{2^k}(\rho) \rho^{d-1} \chi_{\leq 10}(\rho s) s^{\beta+d-1} |f(s)| ds d\rho \\ &\lesssim \int_0^{+\infty} \chi_{2^k}(\rho) \rho^{d-\frac{7}{2}} \|\chi_{\leq 10}(\rho s) s^{d-3}\|_{L_s^2} \|s^{\frac{d-1}{2}} f(s)\|_{L_s^2} d\rho \\ &\lesssim \int_0^{+\infty} \rho^{-1} \chi_{2^k}(\rho) d\rho \|s^{\frac{d-1}{2}} f(s)\|_{L_s^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \end{aligned} \quad (3.38)$$

Now we estimate (3.37-2). From (3.15), we write

$$\begin{aligned} (3.37-2) &= r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta} \\ &\quad \cdot \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 10}(\rho s) s^{\beta+d-1} f(s) ds d\rho \end{aligned} \quad (3.39-i)$$

$$+ r^2 \int_0^{+\infty} J(\rho r) \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O((\rho s)^{-10}) \chi_{\geq 10}(\rho s) s^{\beta+d-1} f(s) ds d\rho. \quad (3.39-ii)$$

The terms (3.39-i) is equal to

$$\begin{aligned} & r^2 \int_0^{\frac{\pi}{2}} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{+\infty} \int_0^{+\infty} e^{2\pi i \rho(-s \sin \theta' + r \sin \theta)} \chi_{\leq 1}(\rho r) \chi_{2^k}(\rho) \chi_{\geq 10}(\rho s) \\ & \cdot \rho^{d-1} s^{\beta+d-1} f(s) d\rho ds \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' \cos^{d-2} \theta d\theta' d\theta. \end{aligned} \quad (3.40)$$

Note that the power $\beta + d - 1$ is too large to be integrable in s , but we can use integration-by-parts to decrease the power of s , due to the non-resonance of the phase. Indeed, when $|\theta'| \geq \frac{\pi}{7}$, $s \geq 5r$, we have (3.28). Thus using the formula (3.29) and integration-by-parts 10 times, (3.39-i) can be controlled by

$$r^2 \int_0^{+\infty} \int_0^{+\infty} \chi_{\lesssim 1}(\rho r) \chi_{2^k}(\rho) \chi_{\gtrsim 1}(\rho s) \rho^{d-11} s^{\beta+d-11} |f(s)| d\rho ds.$$

Therefore, we obtain

$$\|(3.39-i)\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

The term (3.39-ii) can be controlled by

$$\int_0^{+\infty} \int_0^{+\infty} \chi_{\leq 1}(\rho r) \chi_{\geq 10}(\rho s) \chi_{2^k}(\rho) \rho^{d-13} s^{d-13} \cdot s^{\frac{d-1}{2}} |f(s)| ds d\rho.$$

Hence, similar as the estimate on (3.37-1), we have

$$\|(3.39-ii)\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

Combining this last estimate with the estimate on (3.39-i), we get

$$\|(3.37-2)\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

This together with (3.38), gives

$$\|(3.36a)\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.41)$$

Estimates on (3.36b). Note that

$$\begin{aligned} (3.36b) &= r^2 \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \\ & \cdot \rho^{d-1} \int_0^{+\infty} (J(\rho s) + J(-\rho s)) s^{\beta+d-1} f(s) ds d\rho. \end{aligned}$$

Note that for $J(\rho r) - K(\rho r)$, we have the equality given in (3.13); and for $J(\rho s) + J(-\rho s)$ we have the equality given in (3.15). Then we divide (3.36b) into the following four parts. The first part is

$$\begin{aligned} & r^2 \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.42-1)$$

the second part is

$$\begin{aligned} & r^2 \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.42-2)$$

the third part is

$$\begin{aligned} & r^2 \int_0^{+\infty} O((\rho r)^{-5}) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.42-3)$$

and the fourth part is

$$r^2 \int_0^{+\infty} O((\rho r)^{-5}) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} f(s) ds d\rho. \quad (3.42-4)$$

Estimate on (3.42-1). We split (3.42-1) into the following two subparts again,

$$\begin{aligned} & r^2 \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' (\chi_{\leq \frac{1}{10}r}(s) + \chi_{\geq 10r}(s)) s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.43-i)$$

and

$$\begin{aligned} & r^2 \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' \chi_{\frac{1}{10}r \leq \cdot \leq 10r}(s) s^{\beta+d-1} f(s) ds d\rho. \end{aligned} \quad (3.43-ii)$$

For (3.43-i), we rewrite it as

$$\begin{aligned} & r^2 \int_0^{\frac{\pi}{2}} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{+\infty} \int_0^{+\infty} e^{2\pi i \rho(r \sin \theta - s \sin \theta')} \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \chi_{\geq \frac{\pi}{6}}(\theta) \\ & \cdot \cos^{d-2} \theta \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' (\chi_{\leq \frac{1}{10}r}(s) + \chi_{\geq 10r}(s)) s^{\beta+d-1} f(s) d\rho ds d\theta' d\theta. \end{aligned}$$

Note that we have (3.28) in this situation. Thus treated similarly as the term (3.39-i), using the formula (3.29) and integration-by-parts 10 times, (3.43-i) has the bound of

$$r^2 \chi_{\gtrsim 2^{-k}}(r) \int_0^{+\infty} \int_0^{+\infty} (r+s)^{-10} \rho^{-10} \chi_{2^k}(\rho) \rho^{d-1} s^{\beta+d-1} |f(s)| d\rho ds.$$

Hence,

$$\|(3.43-i)\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.44)$$

For (3.43-ii), by dyadic decomposition, it is equal to

$$\sum_{j=-k}^{\infty} \chi_{2^j}(r) \cdot (3.43-ii).$$

Note that $\chi_{2j}(r) \cdot (3.43\text{-ii})$ can be written by

$$\begin{aligned} & r^2 \chi_{2j}(r) \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{\geq 1}(\rho r) \chi_{2k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' \chi_{2j-2 \leq \cdot \leq 2j+2}(s) s^{\beta+d-1} f(s) ds d\rho \\ & + \text{other terms.} \end{aligned} \quad (3.45)$$

Here the “*other terms*” can be treated in the same manner as (3.43-i) and thus we ignore them. For (3.45), from Lemma 3.6, we have the following two formulas,

$$\int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta = c(\rho r)^{-\frac{d-1}{2}} e^{2\pi i \rho r} + O((\rho r)^{-\frac{d+1}{2}});$$

and

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' = (\rho s)^{-\frac{d-1}{2}} \left(\bar{c} e^{-2\pi i \rho s} + c e^{2\pi i \rho s} \right) + O((\rho s)^{-\frac{d+1}{2}}).$$

Accordingly, we divide (3.45) into four parts as (3.42-1)–(3.42-4). The first part is

$$\begin{aligned} & r^2 \chi_{2j}(r) \int_0^{+\infty} (\rho r)^{-\frac{d-1}{2}} e^{2\pi i \rho r} \chi_{\geq 1}(\rho r) \chi_{2k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} (\rho s)^{-\frac{d-1}{2}} \left(\bar{c} e^{-2\pi i \rho s} + c e^{2\pi i \rho s} \right) \chi_{2j-2 \leq \cdot \leq 2j+2}(s) s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.46\text{-iia})$$

the second part is

$$\begin{aligned} & r^2 \chi_{2j}(r) \int_0^{+\infty} (\rho r)^{-\frac{d-1}{2}} e^{2\pi i \rho r} \chi_{\geq 1}(\rho r) \chi_{2k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} O((\rho s)^{-\frac{d+1}{2}}) \chi_{2j-2 \leq \cdot \leq 2j+2}(s) s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.46\text{-iib})$$

the third part is

$$\begin{aligned} & r^2 \chi_{2j}(r) \int_0^{+\infty} O((\rho r)^{-\frac{d+1}{2}}) \chi_{\geq 1}(\rho r) \chi_{2k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} (\rho s)^{-\frac{d-1}{2}} \left(\bar{c} e^{-2\pi i \rho s} + c e^{2\pi i \rho s} \right) \chi_{2j-2 \leq \cdot \leq 2j+2}(s) s^{\beta+d-1} f(s) ds d\rho; \end{aligned} \quad (3.46\text{-iic})$$

and the fourth part is

$$\begin{aligned} & r^2 \chi_{2j}(r) \int_0^{+\infty} O((\rho r)^{-\frac{d+1}{2}}) \chi_{\geq 1}(\rho r) \chi_{2k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} O((\rho s)^{-\frac{d+1}{2}}) \chi_{2j-2 \leq \cdot \leq 2j+2}(s) s^{\beta+d-1} f(s) ds d\rho. \end{aligned} \quad (3.46\text{-iid})$$

First, we consider (3.46-iia), which is equivalent to

$$\begin{aligned} & 2^{(2-\frac{d-1}{2})j} \chi_{2j}(r) \int_0^{+\infty} e^{2\pi i \rho r} \chi_{\geq 1}(\rho r) \chi_{2k}(\rho) \\ & \cdot \int_0^{+\infty} \left(\bar{c} e^{-2\pi i \rho s} + c e^{2\pi i \rho s} \right) \chi_{2j-2 \leq \cdot \leq 2j+2}(s) s^{\frac{d-5}{2}} s^{\frac{d-1}{2}} f(s) ds d\rho. \end{aligned}$$

Then using Corollary 2.12 and Plancherel's identity, we have

$$\begin{aligned} \|(3.46\text{-iia})\|_{L_r^2} &\lesssim 2^{\frac{5-d}{2}j} \left\| \int_0^{+\infty} \left(\bar{c}e^{-2\pi i \rho s} + ce^{2\pi i \rho s} \right) \chi_{2^{j-2} \leq \cdot \leq 2^{j+2}}(s) s^{\frac{d-5}{2}} s^{\frac{d-1}{2}} f(s) ds \right\|_{L_\rho^2} \\ &\lesssim 2^{\frac{5-d}{2}j} \left\| \chi_{2^{j-2} \leq \cdot \leq 2^{j+2}}(s) s^{\frac{d-5}{2}} s^{\frac{d-1}{2}} f(s) \right\|_{L_s^2} \lesssim \|f\|_{L^2(\{|x| \sim 2^j\})}. \end{aligned} \quad (3.47)$$

Using the relationship $r \sim s$, the term (3.46-iib)–(3.46-iid) can be treated in the similar way as (3.42-2)–(3.42-4). We just take the term (3.46-iib) for example. Indeed, by Corollary 2.12, we have

$$\begin{aligned} \|(3.46\text{-iib})\|_{L_r^2} &\lesssim 2^{(2-\frac{d-1}{2})j} \left\| \rho^{-\frac{d-1}{2}} \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O((\rho s)^{-\frac{d+1}{2}}) \chi_{2^{j-2} \leq \cdot \leq 2^{j+2}}(s) s^{\beta+d-1} f(s) ds \right\|_{L_\rho^2} \\ &\lesssim \left\| \chi_{2^{j-2} \leq \cdot \leq 2^{j+2}}(s) s^{\frac{d-1}{2}} f(s) \right\|_{L_s^2} \lesssim \|f\|_{L^2(\{|x| \sim 2^j\})}. \end{aligned} \quad (3.48)$$

Similar as (3.42-3), (3.42-4), we have

$$\|(3.46\text{-iic})\|_{L_r^2} + \|(3.46\text{-iid})\|_{L_r^2} \lesssim \|f\|_{L^2(\{|x| \sim 2^j\})}. \quad (3.49)$$

Combining with (3.48), (3.49) and (3.47), we obtain that

$$\|(3.45)\|_{L_r^2} \lesssim \|f\|_{L^2(\{|x| \sim 2^j\})}.$$

Hence, we have

$$\|\chi_{2^j}(r) \cdot (3.43\text{-ii})\|_{L_r^2} \lesssim \|f\|_{L^2(\{|x| \sim 2^j\})}.$$

Therefore,

$$\|(3.43\text{-ii})\|_{L_r^2}^2 \lesssim \sum_{j=-k}^{+\infty} \|\chi_{2^j}(r) \cdot (3.43\text{-ii})\|_{L_r^2}^2 \lesssim \sum_{j=-k}^{+\infty} \|f\|_{L^2(\{|x| \sim 2^j\})}^2 \lesssim \|f\|_{L^2(\mathbb{R}^d)}^2.$$

This last estimate together with (3.44), yields that

$$\|(3.42-1)\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.50)$$

Now we consider the second part (3.42-2). Using the formula,

$$\partial_\rho e^{2\pi i \rho r \sin \theta} = 2\pi i r \sin \theta \cdot e^{2\pi i \rho r \sin \theta},$$

and integration-by-parts, we reduce (3.42-2) to

$$\begin{aligned} &\int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \frac{\chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta}{(2\pi i \sin \theta)^2} d\theta \partial_\rho^2 \left[\chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \right. \\ &\quad \left. \cdot \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} f(s) ds \right] d\rho. \end{aligned}$$

Then by Corollary 2.12, we have

$$\begin{aligned} \|(3.42-2)\|_{L_r^2} &\lesssim \int_0^{\frac{\pi}{2}} \left\| \partial_\rho^2 \left[\chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} f(s) ds \right] \right\|_{L_\rho^2} \chi_{\geq \frac{\pi}{6}}(\theta) \frac{\cos^{d-2} \theta}{\sin^{\frac{5}{2}} \theta} d\theta \\ &\lesssim \left\| \partial_\rho^2 \left[\chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O(\langle \rho s \rangle^{-10}) s^{\beta+d-1} f(s) ds \right] \right\|_{L_\rho^2}. \end{aligned}$$

Here we regard $\chi_1, \chi'_1, \chi''_1$ as the same because they have the same properties we need. From the explicit formula in (3.10), we note that

$$\partial_\rho^j [O(\langle \rho s \rangle^{-10})] = O(s^j \langle \rho s \rangle^{-10}), \quad \text{for } j = 0, 1, 2.$$

Hence, we get that

$$\begin{aligned}
& \left| \partial_\rho^2 \left[\chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O((\rho s)^{-10}) s^{\beta+d-1} f(s) ds \right] \right| \\
& \lesssim \int_0^{+\infty} \chi_{2^k}(\rho) \rho^{d-1} \langle \rho s \rangle^{-10} s^2 s^{\beta+d-1} |f(s)| ds + \int_0^{+\infty} \chi_{2^k}(\rho) \rho^{d-3} \langle \rho s \rangle^{-10} s^{\beta+d-1} |f(s)| ds \\
& \lesssim \int_0^{+\infty} \chi_{2^k}(\rho) [(\rho s)^{d-1} + (\rho s)^{d-3}] \langle \rho s \rangle^{-10} s^{\frac{d-1}{2}} |f(s)| ds.
\end{aligned}$$

Therefore, by Hölder's inequality, we have

$$\|(3.42-2)\|_{L_r^2} \lesssim \|s^{\frac{d-1}{2}} f\|_{L_s^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.51)$$

For the term (3.42-3), we split it into the following two subsubparts again,

$$\begin{aligned}
& r^2 \int_0^{+\infty} O((\rho r)^{-5}) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\
& \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' \chi_{\leq 2^{-k}}(s) s^{\beta+d-1} f(s) ds d\rho; \quad (3.52-i)
\end{aligned}$$

and

$$\begin{aligned}
& r^2 \int_0^{+\infty} O((\rho r)^{-5}) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\
& \cdot \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' \chi_{\geq 2^{-k}}(s) s^{\beta+d-1} f(s) ds d\rho. \quad (3.52-ii)
\end{aligned}$$

By Hölder's inequality, we have

$$\begin{aligned}
| (3.52-i) | & \lesssim r^{-3} \left\| \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-6} \right\|_{L_\rho^2} \\
& \cdot \left\| \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\leq 2^{-k}}(s) s^{\beta+d-1} f(s) ds \right\|_{L_\rho^2} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta',
\end{aligned}$$

then by Plancherel's identity, it is further bounded by

$$r^{-3} \chi_{\geq 2^{-k-1}}(r) 2^{(d-\frac{1}{2}-5)k} \left\| \chi_{\leq 2^{-k}}(s) s^{d-3} s^{\frac{d-1}{2}} f(s) \right\|_{L_s^2}.$$

Hence, since $d \geq 3$, we have

$$\|(3.52-i)\|_{L_r^2} \lesssim \|s^{\frac{d-1}{2}} f\|_{L_s^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.53)$$

Now we consider (3.52-ii). From Lemma 3.6, we have

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta'} \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\theta' = (\rho s)^{-\frac{d-1}{2}} \left(\bar{c} e^{-2\pi i \rho s} + c e^{2\pi i \rho s} \right) + O((\rho s)^{-\frac{d+1}{2}}).$$

Accordingly, (3.52-ii) can be further divided into the following two sub³-parts again,

$$\begin{aligned}
& r^2 \int_0^{+\infty} O((\rho r)^{-5}) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\
& \cdot \int_0^{+\infty} (\rho s)^{-\frac{d-1}{2}} \left(\bar{c} e^{2\pi i \rho s} + c e^{-2\pi i \rho s} \right) \chi_{\geq 2^{-k}}(s) s^{\beta+d-1} f(s) ds d\rho; \quad (3.54-ia)
\end{aligned}$$

and

$$\begin{aligned} & r^2 \int_0^{+\infty} O((\rho r)^{-5}) \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \rho^{d-1} \\ & \cdot \int_0^{+\infty} O((\rho s)^{-\frac{d+1}{2}}) \chi_{\geq 2^{-k}}(s) s^{\beta+d-1} f(s) ds d\rho. \end{aligned} \quad (3.54\text{-iib})$$

Thanks to the power of $(\rho s)^{-\frac{d+1}{2}}$ in the integral, we can deal with (3.54-ia) as in (3.52-i), which implies that

$$|(3.54\text{-ia})| \lesssim r^{-3} \chi_{\geq 2^{-k-1}}(r) 2^{(\frac{d}{2}-5)k} \left\| \chi_{\geq 2^{-k}}(s) s^{\frac{d-5}{2}} s^{\frac{d-1}{2}} f(s) \right\|_{L_s^2}.$$

Since $d \leq 5$, we have

$$\|(3.54\text{-ia})\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.55)$$

The term (3.54-iib) is controlled by

$$r^{-3} \int_0^{+\infty} \int_0^{+\infty} \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \chi_{\geq 2^{-k}}(s) \rho^{\frac{d-13}{2}} s^{\frac{d-7}{2}} s^{\frac{d-1}{2}} |f(s)| ds d\rho.$$

Hence, by Hölder's inequality, we obtain that

$$\|(3.54\text{-iib})\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.56)$$

Therefore, together with (3.55) and (3.56), we get

$$\|(3.52\text{-ii})\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

Hence, combining this last estimate with (3.53), we get

$$\|(3.42\text{-3})\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.57)$$

For the term (3.42-4), it can be controlled by

$$r^{-3} \int_0^{+\infty} \int_0^{+\infty} \chi_{\geq 1}(\rho r) \chi_{2^k}(\rho) \langle \rho s \rangle^{-10} \rho^{d-6} s^{\beta+d-1} f(s) ds d\rho.$$

Similar as (3.54-iib), we have that

$$\|(3.42\text{-4})\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}. \quad (3.58)$$

This together with (3.51), (3.57) and (3.58), gives

$$\|(3.36\text{b})\|_{L_r^2} \lesssim \|f\|_{L^2(\mathbb{R}^d)}.$$

Hence, this last estimate above combined with (3.41), yields the desired estimate. \square

Based on Proposition 3.13, we have the following estimate which can be regarded as the extension of the Bernstein estimate.

Corollary 3.15. *For any $s \in [0, 4]$, and any integer $k \geq 0$, suppose that $f \in \dot{H}^s(\mathbb{R}^d)$, then*

$$\left\| \chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out/in, k-1 \leq \cdot \leq k+1} \right\|_{\dot{H}^s(\mathbb{R}^d)} \lesssim 2^{sk} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

Here the implicit constant is independent of k .

Proof. We only consider the case when $s = 4$, since the other cases can be obtained by the interpolation.

First, we give some reductions. Note that for the radial function g , we have the following formula, for any integer $s \geq 0$,

$$(-\Delta)^{\frac{s}{2}}g(x) = \sum_{j=0}^s c_j r^{j-s} \partial_r^j g(r),$$

where c_j are the constants dependent on d and j , and $r = |x|$. Therefore,

$$\begin{aligned} & \left\| \chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1}f))_{out/in, k-1 \leq \cdot \leq k+1} \right\|_{\dot{H}^4(\mathbb{R}^d)} \\ & \lesssim \sum_{j=0}^4 c_j \left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} \partial_r^j (P_{2^k}(\chi_{\geq 1}f))_{out/in, k-1 \leq \cdot \leq k+1}(r) \right\|_{L_r^2}. \end{aligned}$$

Hence, we only need to show that for any integer $j \leq 4$,

$$\left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} \partial_r^j (P_{2^k}(\chi_{\geq 1}f))_{out/in, k-1 \leq \cdot \leq k+1}(r) \right\|_{L_r^2} \lesssim 2^{jk} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}. \quad (3.59)$$

Furthermore, from Definition 3.4,

$$\begin{aligned} \partial_r^j (P_{2^k}(\chi_{\geq 1}f))_{out/in, k-1 \leq \cdot \leq k+1}(r) &= \sum_{j_1=0}^j c_{j_1} r^{-\beta-j+j_1} \\ &\cdot \int_0^{+\infty} \partial_r^{j_1} (J(\rho r) - K(\rho r)) \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho, \end{aligned} \quad (3.60)$$

where we have simplified the notation and denote

$$\chi_{2^k}(\rho) = \chi_{2^{k-1} \leq \cdot \leq 2^{k+1}}(\rho),$$

and the constants c_j may change line to line in the following. Moreover, we note that

$$\partial_r^{j_1} (J(\rho r) - K(\rho r)) = \left(\frac{\rho}{r}\right)^{j_1} \partial_\rho^{j_1} (J(\rho r) - K(\rho r)).$$

Hence, (3.60) reduces to

$$\sum_{j_1=0}^j c_{j_1} r^{-\beta-j} \int_0^{+\infty} \partial_\rho^{j_1} (J(\rho r) - K(\rho r)) \chi_{2^k}(\rho) \rho^{d-1+j_1} \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho.$$

Further, by integration-by-parts, it turns to

$$\sum_{j_1=0}^j c_{j_1} r^{-\beta-j} \int_0^{+\infty} (J(\rho r) - K(\rho r)) \partial_\rho^{j_1} [\chi_{2^k}(\rho) \rho^{d-1+j_1} \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho)] d\rho.$$

Therefore, we obtain that

$$\begin{aligned} & \partial_r^j (P_{2^k}(\chi_{\geq 1}f))_{out/in, k-1 \leq \cdot \leq k+1}(r) \\ &= \sum_{j_1=0}^j c_{j_1} r^{-\beta-j} \int_0^{+\infty} (J(\rho r) - K(\rho r)) \partial_\rho^{j_1} [\chi_{2^k}(\rho) \rho^{j_1} \rho^{d-1}] \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho \\ & \quad + \sum_{j_1=0}^j c_{j_1} r^{-\beta-j} \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{2^k}(\rho) \rho^{j_1} \rho^{d-1} \partial_\rho^{j_1} [\mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho)] d\rho. \end{aligned}$$

Therefore, to prove (3.59), we only need to show the following two estimates: For any $j_1 \leq j$,

$$\left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} r^{-\beta-j} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \partial_\rho^{j_1} \left[\chi_{2^k}(\rho) \rho^{j_1} \rho^{d-1} \right] \cdot \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho \right\|_{L_r^2} \lesssim 2^{jk} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}; \quad (3.61a)$$

and

$$\left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} r^{-\beta-j} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{j_1} \rho^{d-1} \cdot \partial_\rho^{j_1} \left[\mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) \right] d\rho \right\|_{L_r^2} \lesssim 2^{jk} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}. \quad (3.61b)$$

For (3.61a), note that $r \gtrsim 1, k \geq 0$, it thus reduces to show

$$\left\| r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \partial_\rho^{j_1} \left[\chi_{2^k}(\rho) \rho^{j_1} \rho^{d-1} \right] \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho \right\|_{L^2(\mathbb{R}^d)} \lesssim \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}.$$

Then we can prove it in the same way as Proposition 3.13. Indeed, we can regard

$$\partial_\rho^{j_1} \left[\chi_{2^k}(\rho) \rho^{j_1} \rho^{d-1} \right] \quad \text{as} \quad \chi_{2^k}(\rho) \rho^{d-1},$$

because they have the same properties used in the proof of Proposition 3.13.

For (3.61b), we first consider

$$\partial_\rho^{j_1} \left[\mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) \right],$$

which is equal to

$$\int_0^{+\infty} \partial_\rho^{j_1} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1} P_{2^k}(\chi_{\geq 1}f)(s) ds.$$

Using the relationship

$$\partial_\rho^{j_1} \left(J(\rho s) + J(-\rho s) \right) = \left(\frac{s}{\rho} \right)^{j_1} \partial_s^{j_1} \left(J(\rho s) + J(-\rho s) \right),$$

we have

$$\partial_\rho^{j_1} \left[\mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) \right] = \frac{1}{\rho^{j_1}} \int_0^{+\infty} \partial_s^{j_1} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1+j_1} P_{2^k}(\chi_{\geq 1}f)(s) ds.$$

Then by integration-by-parts, to prove (3.61b), it suffices to show the following two estimates. The first one is

$$\left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} r^{-\beta-j} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \cdot \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1} P_{2^k}(\chi_{\geq 1}f)(s) ds d\rho \right\|_{L_r^2} \lesssim \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}; \quad (3.62-1)$$

and the second one is

$$\begin{aligned} & \left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} r^{-\beta-j} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \right. \\ & \quad \cdot \left. \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1+j_1} \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2} \\ & \lesssim 2^{jk} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \end{aligned} \quad (3.62-2)$$

For (3.62-1), we drop $\chi_{\geq 1}(r) r^{-j}$ and reduce the left-hand side of (3.62-1) to

$$\begin{aligned} & \left\| r^{\frac{d-1}{2}-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \right. \\ & \quad \cdot \left. \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1} P_{2^k}(\chi_{\geq 1} f)(s) ds d\rho \right\|_{L_r^2}. \end{aligned}$$

Note that by Definition 3.4, it is equal to

$$\left\| (P_{2^k}(\chi_{\geq 1} f))_{out,k} \right\|_{L^2(\mathbb{R}^d)}.$$

Thus the estimate (3.62-1) follows from Proposition 3.13.

For (3.62-2), we split its left-hand side term into the following two subparts again:

$$\begin{aligned} & \left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} r^{-\beta-j} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \right. \\ & \quad \cdot \left. \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1+j_1} \chi_{\leq 1}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}; \end{aligned} \quad (3.63-i)$$

and

$$\begin{aligned} & \left\| \chi_{\geq 1}(r) r^{\frac{d-1}{2}} r^{-\beta-j} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \right. \\ & \quad \cdot \left. \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}. \end{aligned} \quad (3.63-ii)$$

By Definition 3.4 again, (3.63-i) is controlled by

$$\left\| \left(\chi_{\leq 1}(r) r^{j_1} \partial_r^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right)_{out,k} \right\|_{L^2(\mathbb{R}^d)}.$$

Hence by Proposition 3.13, it is further bounded by

$$\left\| \chi_{\leq 1}(r) r^{j_1} \partial_r^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right\|_{L^2(\mathbb{R}^d)}. \quad (3.64)$$

Note that for any radial function g ,

$$\partial_r^{j_1} g(r) = \left(\frac{x}{|x|} \cdot \nabla \right)^{j_1} g(x). \quad (3.65)$$

Therefore, using (3.64) and the formula (3.65), we have that

$$(3.63-i) \lesssim 2^{j_1 k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

For (3.63-ii), we note that

$$(3.63\text{-ii})^2 = \sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{\frac{d-1}{2}} r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \right. \\ \left. \cdot \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}^2.$$

Due to this, we may split (3.63-ii)² into the following three subsubparts again. The first one is

$$\sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{\frac{d-1}{2}} r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) \right. \\ \left. \cdot s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \chi_{2^{h-2} \leq \cdot \leq 2^{h+2}}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}^2; \quad (3.66\text{-ii-1})$$

the second one is

$$\sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{\frac{d-1}{2}} r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) \right. \\ \left. \cdot s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \chi_{\geq 2^{h+2}}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}^2; \quad (3.66\text{-ii-2})$$

and the third one is

$$\sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{\frac{d-1}{2}} r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} \left(J(\rho s) + J(-\rho s) \right) \right. \\ \left. \cdot s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \chi_{\leq 2^{h-2}}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}^2. \quad (3.66\text{-ii-3})$$

Similar as (3.63-i), from Definition 3.4 and Proposition 3.13, we have

$$(3.66\text{-ii-1}) \lesssim \sum_{h=0}^{+\infty} 2^{-jh} \left\| r^{j_1} \chi_{2^{h-2} \leq \cdot \leq 2^{h+2}}(r) \chi_{\geq 1}(r) \partial_r^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right\|_{L^2(\mathbb{R}^d)}^2.$$

Since $j_1 \leq j$, it is further dominated by

$$\sum_{h=0}^{+\infty} \left\| \chi_{2^{h-2} \leq \cdot \leq 2^{h+2}}(r) \chi_{\geq 1}(r) \partial_r^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right\|_{L^2(\mathbb{R}^d)}^2 \lesssim \left\| \chi_{\geq 1}(r) \partial_r^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right\|_{L^2(\mathbb{R}^d)}^2.$$

Using (3.65) again, it is bounded by $2^{2j_1 k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}^2$.

For (3.66-ii-2) and (3.66-ii-3), we only consider the former, since they can be treated in the same manner. From Corollary 3.10, we split (3.66-ii-2) into the following two terms again,

$$\sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{\frac{d-1}{2}} r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-2\pi i \rho s \sin \theta} \right. \\ \left. \cdot \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \chi_{\geq 2^{h+2}}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \right\|_{L_r^2}^2; \quad (3.67\text{-ii-2a})$$

and

$$\begin{aligned} & \sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{-\beta} \int_0^{+\infty} \left(J(\rho r) - K(\rho r) \right) \chi_{2^k}(\rho) \rho^{d-1} \int_0^{+\infty} O((\rho s)^{-10}) \right. \\ & \quad \cdot \chi_{\geq 1}(s) \chi_{\geq 2^{h+2}}(s) s^{\beta+d-1+j_1} \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] ds d\rho \left. \right\|_{L^2(\mathbb{R}^d)}^2. \end{aligned} \quad (3.67\text{-ii-2b})$$

For (3.67-ii-2a), due to the non-resonance of the phase, it can be treated similarly as (3.43-i). Indeed, from Corollary 3.9, we write

$$J(\rho r) - K(\rho r) = \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \eta(\theta, \rho r) d\theta,$$

where $\eta(\theta, \rho r)$ is defined as

$$\begin{aligned} \eta(\theta, \rho r) &= \chi_{\leq 1}(\rho r) \cos^{d-2} \theta + \chi_{\geq 1}(\rho r) \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta \\ &\quad + \chi_{1 \leq \cdot \leq 2}(\rho r) \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta + \chi_{\geq 2}(\rho r) \frac{1}{(2\pi i \rho r)^5} \tilde{\eta}_d(\theta). \end{aligned}$$

Therefore, (3.67-ii-2a) can be rewritten as

$$\begin{aligned} & \sum_{h=0}^{+\infty} 2^{-jh} \left\| \chi_{2^h}(r) r^{\frac{d-1}{2}} r^{-\beta} \int_0^{\frac{\pi}{2}} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{+\infty} \int_0^{+\infty} e^{2\pi i \rho(r \sin \theta - s \sin \theta')} \rho^{d-1} \eta(\theta, \rho r) \chi_{2^k}(\rho) \right. \\ & \quad \cdot s^{\beta+d-1+j_1} \chi_{\geq 1}(s) \chi_{\geq 2^{h+2}}(s) \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \chi_{\geq \frac{\pi}{6}}(\theta') \cos^{d-2} \theta' d\rho ds d\theta' d\theta \left. \right\|_{L_r^2}^2. \end{aligned}$$

Note that we have (3.28). Thus using the formula (3.29) and integration-by-parts 10 times, we further control (3.67-ii-2a) as

$$\sum_{h=0}^{+\infty} \left\| \chi_{2^h}(r) \int_0^{+\infty} \int_0^{+\infty} \rho^{d-11} \chi_{2^k}(\rho) s^{d-11} \chi_{\geq 1}(s) \chi_{\geq 2^{h+2}}(s) \left| \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right| ds d\rho \right\|_{L^2(\mathbb{R}^d)}^2.$$

Hence, by the Hölder inequality, we get

$$(3.67\text{-ii-2a}) \lesssim \left\| \chi_{\geq 1}(r) \partial_r^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right\|_{L^2(\mathbb{R}^d)}^2 \lesssim 2^{2j_1 k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}^2.$$

For (3.67-ii-2b), thanks to the high-order decay of ρ and s , by Hölder's inequality, it is controlled by

$$\sum_{h=0}^{+\infty} 2^{-jh} 2^{(\frac{d}{2}-\beta)h} 2^{-9k} \left\| \chi_{\geq 2^{h+2}}(s) s^{\beta+d-10+j_1} \partial_s^{j_1} \left[P_{2^k}(\chi_{\geq 1} f) \right] \right\|_{L_s^2}^2.$$

Hence, it is also bounded by

$$2^{2j_1 k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}^2.$$

Therefore, we obtain that

$$(3.66\text{-ii-2}) \lesssim 2^{2j_1 k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}^2.$$

Collecting the estimates above, we finish the proof of the corollary. \square

4. ESTIMATES ON THE INCOMING/OUTGOING LINEAR FLOW

In this subsection, we present some important properties on the incoming/outgoing linear flow.

4.1. “Incoming/outgoing” decomposition of the linear flow. Let $N \geq 1$. First of all, we show that the linear flow $e^{it\Delta}(P_{\geq N}\chi_{\geq 1}f)_{out}$ is almost “outgoing” with the frequency dependent velocity. To do this, we use the Littlewood-Paley decomposition. Without loss of generality, we may assume that $N = 2^{k_0}$ for some $k_0 \in \mathbb{N}$, then

$$P_{\geq N}\chi_{\geq 1}f = \sum_{k=k_0}^{\infty} P_{2^k}(\chi_{\geq 1}f),$$

and thus

$$(P_{\geq N}\chi_{\geq 1}f)_{out/in} = \sum_{k=k_0}^{\infty} (P_{2^k}(\chi_{\geq 1}f))_{out/in}.$$

Now we consider $(P_{2^k}(\chi_{\geq 1}f))_{out/in}$, for which we only consider the outgoing part. First, from Proposition 3.11, we have

$$(P_{2^k}(\chi_{\geq 1}f))_{out} = (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} + h_k, \quad (4.1)$$

where h_k satisfies the following estimate,

$$\|h_k\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-10k} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}. \quad (4.2)$$

Due to this estimate, we only need to consider $(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}$. Further, we write

$$\begin{aligned} & (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \\ &= \chi_{\leq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} + \chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}. \end{aligned}$$

From Lemma 3.12, we have

$$\|\chi_{\leq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}\|_{H^{\mu(d)}(\mathbb{R}^d)} \lesssim 2^{-2k} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}. \quad (4.3)$$

So it is left to consider $\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}$. The first estimate is

Proposition 4.1. *Let $k \geq 0$ be an integer. Then there exists $\delta > 0$, such that for any triple (γ, q, r) satisfying that*

$$q \geq 2, \quad r > 2, \quad 0 \leq \gamma \leq 1, \quad \frac{2}{q} + \frac{2d-1}{r} < \frac{2d-1}{2}, \quad (4.4)$$

the following estimate holds,

$$\begin{aligned} & \left\| |\nabla|^\gamma \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \\ & \lesssim 2^{-\left(5-d-(\gamma-\frac{2}{q}-\frac{d}{r})k\right)} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

The same estimate holds when $e^{it\Delta}$ and $_{out}$ are replaced by $e^{-it\Delta}$ and $_{in}$, respectively.

Proof. We only consider the estimates on the “outgoing” part, since the “incoming” part can be treated in the same way.

First, from the definition,

$$\begin{aligned} & (P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}(r) \\ &= r^{-\beta} \int_0^{+\infty} (J(\rho r) - K(\rho r)) \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho. \end{aligned}$$

Here we use the notation $\chi_{2^k}(\rho) = \chi_{2^{k-1} \leq \cdot \leq 2^{k+1}}(\rho)$ again. We denote $\tilde{\eta}(\theta, r)$ as

$$\tilde{\eta}(\theta, r) = \chi_{\leq 2}(r) \chi_{\leq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta + \chi_{\geq 2}(r) \frac{1}{(2\pi i r)^5} \tilde{\eta}_d(\theta).$$

Then

$$\tilde{\eta}(\theta, r) = O(\langle r \rangle^{-5}). \quad (4.5)$$

Moreover, from Corollary 3.9,

$$J(r) - K(r) = \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta + \int_0^{\frac{\pi}{2}} e^{2\pi i r \sin \theta} \tilde{\eta}(\theta, r) d\theta.$$

Accordingly, we split $(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}(r)$ into two parts, which are denoted by F_1 and F_2 , as follows:

$$F_1(r) = r^{-\beta} \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \tilde{\eta}(\theta, \rho r) d\theta \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho;$$

and

$$F_2(r) = r^{-\beta} \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\rho) d\rho.$$

We consider F_1 first, and claim that for $s \in [0, 3]$,

$$\|\chi_{\geq \frac{1}{4}} F_1\|_{\dot{H}^s(\mathbb{R}^d)} \lesssim 2^{-(5-\frac{d}{2}-s)k} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}. \quad (4.6)$$

To prove (4.6), we need the following lemma.

Lemma 4.2. *Let $d = 3, 4, 5$, and $f \in L^2(\mathbb{R}^d)$, then*

$$\|\chi_{\geq 2^k}(\rho) \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))\|_{L^2_\rho} \lesssim 2^{-\frac{d-1}{2}k} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)}.$$

Proof. Note that

$$\begin{aligned} \|\chi_{\geq 2^k}(\rho) \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))\|_{L^2_\rho} &= \|\xi|^{-\frac{d-1}{2}} \chi_{\geq 2^k}(\xi) \mathcal{F}(P_{2^k}(\chi_{\geq 1}f))(\xi)\|_{L^2_\xi(\mathbb{R}^d)} \\ &\lesssim 2^{-\frac{d-1}{2}k} \|\mathcal{F}(P_{2^k}(\chi_{\geq 1}f))\|_{L^2_\xi(\mathbb{R}^d)}. \end{aligned} \quad (4.7)$$

From (3.3),

$$\mathcal{F}(P_{2^k}(\chi_{\geq 1}f)) = \mathcal{F}(|x|^\beta P_{2^k}(\chi_{\geq 1}f)).$$

Then by the Plancherel identity, we obtain

$$\begin{aligned} (4.7) &= 2^{-\frac{d-1}{2}k} \| |x|^\beta P_{2^k}(\chi_{\geq 1}f) \|_{L^2_x(\mathbb{R}^d)} \\ &\leq 2^{-\frac{d-1}{2}k} \| |x|^\beta \chi_{\leq \frac{1}{4}} P_{2^k}(\chi_{\geq 1}f) \|_{L^2_x(\mathbb{R}^d)} + 2^{-\frac{d-1}{2}k} \| |x|^\beta \chi_{\geq \frac{1}{4}} P_{2^k}(\chi_{\geq 1}f) \|_{L^2_x(\mathbb{R}^d)}. \end{aligned}$$

Note that $3 \leq d \leq 5$, we have that $-1 \leq \beta \leq 0$. Then it is further bounded by

$$2^{-\frac{d-1}{2}k} \|\chi_{\leq \frac{1}{4}} P_{2^k}(\chi_{\geq 1} f)\|_{L^\infty(\mathbb{R}^d)} + 2^{-\frac{d-1}{2}k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

From Lemma 2.4, it is controlled by $2^{-\frac{d-1}{2}k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}$. This gives the desired estimate. \square

By (4.5), Hölder's inequality and Lemma 4.2, we get

$$\begin{aligned} \chi_{\geq \frac{1}{4}}(r) |F_1(r)| &\lesssim r^{-\beta-5} \left\| \chi_{2^k}(\rho) \rho^{d-6} \right\|_{L_\rho^2} \cdot \left\| \chi_{\geq 2^k}(\rho) \mathcal{F}(P_{2^k}(\chi_{\geq 1} f)) \right\|_{L_\rho^2} \\ &\lesssim r^{-\beta-5} 2^{(d-\frac{11}{2})k} 2^{-\frac{d-1}{2}k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)} \\ &\lesssim r^{-\beta-5} 2^{(\frac{d}{2}-5)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

Therefore, we obtain

$$\|\chi_{\geq \frac{1}{4}} F_1\|_{L^2(\mathbb{R}^d)} \lesssim 2^{(\frac{d}{2}-5)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

For the estimate on the high-order derivatives, we note that for $j = 1, 2, 3$,

$$\partial_r^j \int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{2\pi i \rho r \sin \theta} \tilde{\eta}(\theta, \rho r) d\theta = O(r^{-5} \rho^{-5+j}).$$

Hence arguing similarly as above, we get that for $s = 1, 2, 3$,

$$\|\chi_{\geq \frac{1}{4}} F_1\|_{\dot{H}^s(\mathbb{R}^d)} \lesssim 2^{-(5-\frac{d}{2}-s)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

For general $s \in [0, 3]$, we obtain it by the interpolation. This proves (4.6).

Using Lemma 2.3, we obtain that

$$\left\| |\nabla|^\gamma \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta}(\chi_{\geq \frac{1}{4}} F_1) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim \left\| |\nabla|^\gamma e^{it\Delta}(\chi_{\geq \frac{1}{4}} F_1) \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)}.$$

Then from Lemma 2.9, it can be controlled by

$$\|\chi_{\geq \frac{1}{4}} F_1\|_{\dot{H}^s(\mathbb{R}^d)}, \quad \text{with} \quad s = \frac{d}{2} + \gamma - \frac{2}{q} - \frac{d}{r}.$$

Now applying (4.6), it gives

$$\left\| |\nabla|^\gamma \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta}(\chi_{\geq \frac{1}{4}} F_1) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim 2^{-(5-d-(\gamma-\frac{2}{q}-\frac{d}{r}))k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \quad (4.8)$$

Now we consider $e^{it\Delta}(\chi_{\geq \frac{1}{4}} F_2)$, which is equal to

$$\begin{aligned} &\int_0^{+\infty} \int_0^{\frac{\pi}{2}} e^{it\Delta} \left(r^{-\beta} \chi_{\geq \frac{1}{4}}(r) e^{2\pi i \rho r \sin \theta} \right) \chi_{\geq \frac{\pi}{6}}(\theta) \cos^{d-2} \theta d\theta \\ &\quad \cdot \chi_{2^k}(\rho) \rho^{d-1} \mathcal{F}(P_{2^k}(\chi_{\geq 1} f))(\rho) d\rho. \end{aligned} \quad (4.9)$$

To give its estimate, we consider the following term,

$$e^{it\Delta} \left(r^{-\beta} \chi_{\geq \frac{1}{4}}(r) e^{2\pi i \rho r \sin \theta} \right).$$

Using the formula in Lemma 2.7, we have

$$e^{it\Delta} \left(r^{-\beta} \chi_{\geq \frac{1}{4}}(r) e^{2\pi i \rho r \sin \theta} \right) = \frac{C}{t^{\frac{d}{2}}} \int_{\mathbb{R}^d} e^{i \frac{|x-y|^2}{4t} + 2\pi i \rho |y| \sin \theta} |y|^{-\beta} \chi_{\geq \frac{1}{4}}(y) dy.$$

We expand the phase and rewrite it as

$$\begin{aligned} & \frac{c}{t^{\frac{d}{2}}} e^{i\frac{|x|^2}{4t}} \int_{\mathbb{R}^d} e^{-i\frac{x \cdot y}{2t} + i\frac{|y|^2}{4t} + 2\pi i \rho |y| \sin \theta} |y|^{-\beta} \chi_{\geq \frac{1}{4}}(y) dy \\ &= \frac{c}{t^{\frac{d}{2}}} e^{i\frac{|x|^2}{4t}} \int_{|\omega|=1} \int_0^{+\infty} e^{i\phi(r)} r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) dr d\omega, \end{aligned} \quad (4.10)$$

where $\phi(r) = -\frac{x \cdot \omega}{2t} r + \frac{r^2}{4t} + 2\pi \rho r \sin \theta$. Then

$$\phi'(r) = -\frac{x \cdot \omega}{2t} + \frac{r}{2t} + 2\pi \rho \sin \theta.$$

Note that $\rho \sim 2^k$, $r \sim 1$ and $\sin \theta \geq \frac{1}{4}$, so choosing δ small enough, we obtain that when $|x| \leq \delta(1 + 2^k t)$,

$$\phi'(r) \geq \frac{1}{4} \left(\frac{r}{t} + \pi \rho \right). \quad (4.11)$$

Moreover,

$$\phi''(r) = \frac{1}{2t}, \quad \text{and} \quad \phi^{(j)}(r) = 0, \quad \text{for } j = 3, 4, \dots. \quad (4.12)$$

Hence we can use the formula,

$$e^{i\phi(r)} = \frac{1}{i\phi'(r)} \partial_r (e^{i\phi(r)}),$$

and integrate by parts to obtain that for any $K \in \mathbb{Z}^+$, there exists $c_K \in \mathbb{C}$, such that

$$\begin{aligned} & e^{it\Delta} \left(r^{-\beta} \chi_{\geq \frac{1}{4}}(r) e^{2\pi i \rho r \sin \theta} \right) \\ &= \frac{c_K}{t^{\frac{d}{2}}} e^{i\frac{|x|^2}{4t}} \int_{|\omega|=1} \int_0^{+\infty} e^{i\phi(r)} \partial_r \left(\frac{1}{\phi'(r)} \partial_r \right)^{K-1} \left[\frac{1}{\phi'(r)} r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) \right] dr d\omega. \end{aligned} \quad (4.13)$$

Using Lemma 2.6, we have

$$\begin{aligned} & \partial_r \left(\frac{1}{\phi'(r)} \partial_r \right)^{K-1} \left[\frac{1}{\phi'(r)} r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) \right] \\ &= \sum_{\substack{l_1, \dots, l_K \in \mathbb{N}, l' \in \mathbb{N}; \\ l_j \leq j; l_1 + \dots + l_K + l' = K}} C_{l_1, \dots, l_K, l'} \partial_r^{l_1} \left(\frac{1}{\phi'(r)} \right) \dots \partial_r^{l_K} \left(\frac{1}{\phi'(r)} \right) \partial_r^{l'} \left[r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) \right]. \end{aligned}$$

Since $\phi^{(j)}(r) = 0$ when $j \geq 3$, we find that for any integer $l \geq 0$,

$$\partial_r^l \left(\frac{1}{\phi'(r)} \right) = \frac{(\phi''(r))^l}{(\phi'(r))^{l+1}},$$

it implies that

$$\partial_r^{l_1} \left(\frac{1}{\phi'(r)} \right) \dots \partial_r^{l_K} \left(\frac{1}{\phi'(r)} \right) = \frac{(\phi''(r))^{l_1 + \dots + l_K}}{(\phi'(r))^{l_1 + \dots + l_K + K}}.$$

This together with (4.11), (4.12), gives

$$\left| \partial_r^{l_1} \left(\frac{1}{\phi'(r)} \right) \dots \partial_r^{l_K} \left(\frac{1}{\phi'(r)} \right) \right| \lesssim \frac{1}{(r + \rho t)^{l_1 + \dots + l_K} \left(\frac{r}{t} + \rho \right)^K}.$$

Moreover, we have

$$\left| \partial_r^{l'} \left[r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) \right] \right| \lesssim r^{d-1-\beta-l'} \chi_{\gtrsim 1}(r).$$

Combining the above two estimates, and noting that $l_1 + \dots + l_K + l' = K$, we obtain that

$$\left| \partial_r^{l_1} \left(\frac{1}{\phi'(r)} \right) \dots \partial_r^{l_K} \left(\frac{1}{\phi'(r)} \right) \right| \left| \partial_r^{l'} \left[r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) \right] \right| \lesssim \frac{1}{r^K \left(\frac{r}{t} + \rho \right)^K} r^{d-1-\beta} \chi_{\gtrsim 1}(r).$$

Note that when $0 \leq t \leq 1$, it is bounded by

$$\frac{t^3}{r^{K+3} \rho^{K-3}} r^{d-1-\beta} \chi_{\gtrsim 1}(r);$$

when $t \geq 1$, it is bounded by

$$\frac{1}{r^K \rho^K} r^{d-1-\beta} \chi_{\gtrsim 1}(r).$$

Hence, choosing K suitably large, we get

$$\left| \partial_r \left(\frac{1}{\phi'(r)} \partial_r \right)^K \left[\frac{1}{\phi'(r)} r^{d-1-\beta} \chi_{\geq \frac{1}{4}}(r) \right] \right| \lesssim t^3 \langle t \rangle^{-3} r^{-10} \rho^{-10} \chi_{\gtrsim 1}(r).$$

Inserting this estimate into (4.13), we obtain that

$$\left| \chi_{\leq \delta(1+2^k t)} e^{it\Delta} \left(r^{-\beta} \chi_{\geq \frac{1}{4}}(r) e^{2\pi i \rho r \sin \theta} \right) \right| \lesssim \langle t \rangle^{-\frac{d}{2}} \rho^{-10}.$$

Hence, from this estimate and (4.9), we obtain that

$$\left| \chi_{\leq \delta(1+2^k t)} e^{it\Delta} (\chi_{\geq \frac{1}{4}} F_2) \right| \lesssim \langle t \rangle^{-\frac{d}{2}} \int_0^{+\infty} |\chi_{2^k}(\rho) \rho^{d-11} \mathcal{F}(P_{2^k}(\chi_{\geq 1} f))(\rho)| d\rho.$$

Using Hölder's inequality, it is further controlled by

$$\langle t \rangle^{-\frac{d}{2}} \left\| \chi_{2^k}(\rho) \rho^{d-11} \right\|_{L_\rho^2} \cdot \left\| \chi_{\gtrsim 2^k}(\rho) \mathcal{F}(P_{2^k}(\chi_{\geq 1} f)) \right\|_{L_\rho^2}.$$

Now by Lemma 4.2, it is further bounded by

$$\langle t \rangle^{-\frac{d}{2}} 2^{(d-\frac{21}{2})k} 2^{-\frac{d-1}{2}k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

That is,

$$\left| \chi_{\leq \delta(1+2^k t)} e^{it\Delta} (\chi_{\geq \frac{1}{4}} F_2) \right| \lesssim \langle t \rangle^{-\frac{d}{2}} 2^{(\frac{d}{2}-10)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

Note that the structure of $e^{it\Delta}(\chi_{\geq \frac{1}{4}} F_2)$ does not essentially change when we consider the estimates on the high-order derivatives, hence in the same manner, we obtain that for any even integer $\tilde{s} \in [0, 10]$,

$$\left| |\nabla|^{\tilde{s}} \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta} (\chi_{\geq \frac{1}{4}} F_2) \right] \right| \lesssim \langle t \rangle^{-\frac{d}{2}} 2^{(\frac{d}{2}-10)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

Therefore, by Hölder's inequality, if

$$\frac{1}{q} + \frac{d}{r} < \frac{d}{2}, \quad q \geq 2, \quad r \geq 2$$

(note that this condition is implied by (4.4)), then we have

$$\left\| |\nabla|^{\tilde{s}} \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta} (\chi_{\geq \frac{1}{4}} F_2) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim 2^{(\frac{d}{2}-10+\frac{d}{r})k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

Since $d \leq 5$, by the interpolation, we obtain that for any $s \in [0, 10]$,

$$\left\| |\nabla|^s \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta} (\chi_{\geq \frac{1}{4}} F_2) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim 2^{-5k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \quad (4.14)$$

Now collecting the estimates in (4.8) and (4.14), we prove the proposition. \square

The following result is the incoming/outgoing linear flow's estimate related to the outside region.

Proposition 4.3. *Let $k \geq 0$ be an integer. Moreover, let r, γ_1, γ_2, s be the parameters satisfying*

$$r > 2, \quad \gamma_1 \geq 0, \quad \gamma_2 \geq 0, \quad s + \frac{1}{r} \geq \frac{1}{2}, \quad \gamma_1 + s = d\left(\frac{1}{2} - \frac{1}{r}\right).$$

Then for any $t > 0$,

$$\begin{aligned} & \left\| |\nabla|^{\gamma_2} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L^r(\mathbb{R}^d)} \\ & \lesssim (1 + 2^k t)^{-\gamma_1} 2^{(\gamma_2 + s)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

The same estimate holds when $e^{it\Delta}$ and $_{out}$ are replaced by $e^{-it\Delta}$ and $_{in}$, respectively.

Proof. As before, we only consider the estimate on the “outgoing” part. Moreover, we only need to consider the case of $r < \infty$. Indeed, when $r = \infty$, then by the Sobolev inequality,

$$\begin{aligned} & \left\| |\nabla|^{\gamma_2} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L^\infty(\mathbb{R}^d)} \\ & \lesssim \left\| |\nabla|^{\gamma_2} \langle \nabla \rangle^{0+} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L^{\infty-}(\mathbb{R}^d)}. \end{aligned}$$

Hence we only need to consider the case of $r < \infty$, by replacing γ_2 by $\gamma_2 +$ if necessary.

First, we split

$$|\nabla|^{\gamma_2} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right]$$

into the following three terms,

$$P_{\leq 1} |\nabla|^{\gamma_2} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \quad (4.15a)$$

$$+ \chi_{\leq \delta^2(1+2^k t)} P_{\geq 1} |\nabla|^{\gamma_2} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \quad (4.15b)$$

$$+ \chi_{\geq \delta^2(1+2^k t)} P_{\geq 1} |\nabla|^{\gamma_2} \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right]. \quad (4.15c)$$

Now we consider the first term (4.15a). From Lemma 2.1, under the assumptions, we have

$$\| |x|^{\gamma_1} g \|_{L^r(\mathbb{R}^d)} \lesssim \|g\|_{\dot{H}^s(\mathbb{R}^d)}.$$

Hence, this last estimate combined with the Bernstein inequality and Lemma 2.8, gives that

$$\begin{aligned} \|(4.15a)\|_{L^r(\mathbb{R}^d)} & \lesssim \left\| \chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right\|_{L^r(\mathbb{R}^d)} \\ & \lesssim (1 + 2^k t)^{-\gamma_1} \left\| e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right\|_{\dot{H}^s(\mathbb{R}^d)} \\ & \lesssim (1 + 2^k t)^{-\gamma_1} \left\| \chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right\|_{\dot{H}^s(\mathbb{R}^d)}. \end{aligned}$$

Therefore, by Corollary 3.15, we get

$$\|(4.15a)\|_{L^r(\mathbb{R}^d)} \lesssim (1 + 2^k t)^{-\gamma_1} 2^{sk} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

For the second term (4.15b), by Lemma 2.4, we have that for any $M > 0$, $K \geq 1$,

$$\left\| \chi_{\leq \delta K} P_{\geq 1} |\nabla|^{\gamma_2} (\chi_{\geq K} g) \right\|_{L^r(\mathbb{R}^d)} \lesssim_{\delta, M} K^{-M} \|\chi_{\geq K} g\|_{L^r(\mathbb{R}^d)}.$$

Then using the inequality above and treating similarly as (4.15a), we obtain that

$$\begin{aligned} \|(4.15b)\|_{L^r(\mathbb{R}^d)} &\lesssim_M (1 + 2^k t)^{-M} \left\| \chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right\|_{L^r(\mathbb{R}^d)} \\ &\lesssim_M (1 + 2^k t)^{-M-\gamma_1} \left\| \chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right\|_{\dot{H}^s(\mathbb{R}^d)} \\ &\lesssim_M (1 + 2^k t)^{-M-\gamma_1} 2^{sk} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

For the third term (4.15c), we use the similar argument as above, and obtain that

$$\|(4.15c)\|_{L^r(\mathbb{R}^d)} \lesssim (1 + 2^k t)^{-\gamma_1} \left\| \chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right\|_{\dot{H}^{\gamma_2+s}(\mathbb{R}^d)}.$$

From Lemma 2.3 and Lemma 2.8, it is further controlled by

$$(1 + 2^k t)^{-\gamma_1} \left\| \chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right\|_{H^{\gamma_2+s}(\mathbb{R}^d)}.$$

Now using Corollary 3.15, we get

$$\|(4.15c)\|_{L^r(\mathbb{R}^d)} \lesssim (1 + 2^k t)^{-\gamma_1} 2^{(\gamma_2+s)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}.$$

Combining the above estimates on (4.15a)–(4.15c), we finish the proof of the proposition. \square

Now an easy consequence of this proposition is the following space-time estimate.

Corollary 4.4. *Let (γ, q, r) be the triple satisfying*

$$\gamma \geq 0, \quad q \geq 1, \quad r > 2, \quad \frac{1}{q} < (d-1)\left(\frac{1}{2} - \frac{1}{r}\right), \quad (4.16)$$

then the following estimate holds,

$$\begin{aligned} &\left\| |\nabla|^\gamma \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \\ &\lesssim 2^{(-\frac{1}{q} - \frac{1}{r} + \frac{1}{2} + \gamma)k} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}. \end{aligned}$$

The same estimate holds when $e^{it\Delta}$ and out are replaced by $e^{-it\Delta}$ and in , respectively.

Proof. From Proposition 4.3, we have

$$\begin{aligned} &\left\| |\nabla|^\gamma \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}} (P_{2^k}(\chi_{\geq 1} f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \\ &\lesssim 2^{(\gamma+s)k} \|(1 + 2^k t)^{-\gamma_1}\|_{L_t^q(\mathbb{R}^+)} \|P_{2^k}(\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)}, \end{aligned} \quad (4.17)$$

where the parameters

$$s = \frac{1}{2} - \frac{1}{r}, \quad \gamma_1 = (d-1)\left(\frac{1}{2} - \frac{1}{r}\right).$$

By the condition (4.16), we have that $\frac{1}{q} < \gamma_1$, and thus

$$\|(1 + 2^k t)^{-\gamma_1}\|_{L_t^q(\mathbb{R}^+)} \lesssim 2^{-\frac{1}{q}k}.$$

Combining this estimate with (4.17), we prove the corollary. \square

4.2. Improved Strichartz's estimates. Now we collect the estimates above, and obtain the following results.

Proposition 4.5. *Let $N \geq 1, s_0 > 0$. Suppose that (s, q, r) is the triple satisfying that $(q, r) = (\infty, 2)$ or*

$$q \geq 2, \quad r \geq 2, \quad \frac{1}{q} < (d-1)\left(\frac{1}{2} - \frac{1}{r}\right);$$

$$0 \leq s \leq 1, \quad s + \frac{d}{2} - \left(\frac{2}{q} + \frac{d}{r}\right) \leq \mu(d);$$

and

$$\gamma_0 = s_0 - s - \max \left\{ (d-5) - \left(\frac{2}{q} + \frac{d}{r}\right), \frac{1}{2} - \frac{1}{q} - \frac{1}{r} \right\} > 0.$$

Then

$$\| |\nabla|^s e^{it\Delta} (P_{\geq N}(\chi_{\geq 1} f))_{out} \|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim N^{-\gamma_0+} \|P_{\geq N} \chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)}. \quad (4.18)$$

The same estimate holds when $e^{it\Delta}$ and $_{out}$ are replaced by $e^{-it\Delta}$ and $_{in}$, respectively.

Remark 4.6. Now we list some triples (s, q, r) which satisfy the conditions in Proposition 4.5 and will be used later. First, for any $r > \frac{2(d-1)}{d-2}, 0 \leq s \leq 1, s_0 > s - \frac{1}{r}$, we have

$$\| |\nabla|^s e^{it\Delta} (P_{\geq N}(\chi_{\geq 1} f))_{out} \|_{L_t^2 L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim N^{s-s_0-\frac{1}{r}+} \|P_{\geq N} \chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)}. \quad (4.19)$$

Second, for any $r > 2, s_0 > \frac{1}{2} - \frac{1}{r}$,

$$\| e^{it\Delta} (P_{\geq N}(\chi_{\geq 1} f))_{out} \|_{L_t^\infty L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim N^{-s_0+\frac{1}{2}-\frac{1}{r}+} \|P_{\geq N} \chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)}. \quad (4.20)$$

Third, for any $p \geq \max\{1, \frac{4}{d}\}$ and any $s_0 > \frac{1}{2} - \frac{d+2}{2dp}$,

$$\| e^{it\Delta} (P_{\geq N}(\chi_{\geq 1} f))_{out} \|_{L_t^{2p} L_x^{dp}(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim N^{-s_0+\frac{1}{2}-\frac{d+2}{2dp}+} \|P_{\geq N} \chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)}. \quad (4.21)$$

Proof of Proposition 4.5. Again, we only consider the estimates on the “outgoing” part, since the “incoming” part can be treated in the same way. Moreover, by choosing $s < s_0$, the case of $(q, r) = (\infty, 2)$ follows from Corollary 3.15. Hence, we only consider the case when $r > 2$. Let $N = 2^{k_0}$ for some $k_0 \in \mathbb{N}$.

First, we recall the reductions given at the beginning of Subsection 4.1. We write

$$\begin{aligned} e^{it\Delta}(P_{\geq N}\chi_{\geq 1}f)_{out} &= \sum_{k=k_0}^{\infty} e^{it\Delta}(P_{2^k}(\chi_{\geq 1}f))_{out} \\ &= \sum_{k=k_0}^{\infty} e^{it\Delta}\left(\chi_{\leq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} + h_k\right) \\ &\quad + \sum_{k=k_0}^{\infty} e^{it\Delta}\left(\chi_{\geq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1}\right). \end{aligned}$$

Let (s, q, r) be the triple satisfying

$$s \geq 0, \quad \frac{2}{q} + \frac{2d-1}{r} < \frac{2d-1}{2}, \quad s + \frac{d}{2} - \left(\frac{2}{q} + \frac{d}{r}\right) \leq \mu(d).$$

Then we use Lemma 2.9, (4.2) and (4.3), to get

$$\begin{aligned} &\sum_{k=k_0}^{\infty} \left\| e^{it\Delta} |\nabla|^s \left(\chi_{\leq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} + h_k \right) \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \\ &\lesssim \sum_{k=k_0}^{\infty} \left(\left\| \chi_{\leq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \right\|_{H^{\mu(d)}(\mathbb{R}^d)} + \|h_k\|_{H^{\mu(d)}(\mathbb{R}^d)} \right) \\ &\lesssim \sum_{k=k_0}^{\infty} 2^{-2k} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)} \lesssim \|P_{\geq N}\chi_{\geq 1}f\|_{H^{-1}(\mathbb{R}^d)}. \end{aligned} \tag{4.22}$$

Furthermore, we write

$$\begin{aligned} &\sum_{k=k_0}^{\infty} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \right) \\ &= \sum_{k=k_0}^{\infty} \chi_{\leq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \right) \\ &\quad + \sum_{k=k_0}^{\infty} \chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \right). \end{aligned}$$

On one hand, from Proposition 4.1, we have that for any (s, q, r) satisfying (4.4) (note that the condition (4.4) is implied by the conditions in this proposition),

$$\begin{aligned} &\sum_{k=k_0}^{\infty} \left\| |\nabla|^s \left[\chi_{\leq \delta(1+2^k t)} e^{it\Delta} \left(\chi_{\geq \frac{1}{4}}(P_{2^k}(\chi_{\geq 1}f))_{out, k-1 \leq \cdot \leq k+1} \right) \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \\ &\lesssim \sum_{k=k_0}^{\infty} 2^{-\left(5-d-(s-\frac{2}{q}-\frac{d}{r})k\right)} \|P_{2^k}(\chi_{\geq 1}f)\|_{L^2(\mathbb{R}^d)} \lesssim N^{-\gamma_0^+} \|P_{\geq N}\chi_{\geq 1}f\|_{H^{s_0}(\mathbb{R}^d)}. \end{aligned}$$

On the other hand, from Corollary 4.4, we obtain that

$$\begin{aligned} & \sum_{k=k_0}^{\infty} \left\| |\nabla|^s \left[\chi_{\geq \delta(1+2^k t)} e^{it\Delta} \left(P_{2^k} (\chi_{\geq 1} f) \right)_{out, k-1 \leq \cdot \leq k+1} \right] \right\|_{L_t^q L_x^r(\mathbb{R}^+ \times \mathbb{R}^d)} \\ & \lesssim \sum_{k=k_0}^{\infty} 2^{(-\frac{1}{q} - \frac{1}{r} + \frac{1}{2} + s +)k} \|P_{2^k} (\chi_{\geq 1} f)\|_{L^2(\mathbb{R}^d)} \lesssim N^{-\gamma_0 +} \|P_{\geq N} \chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)}. \end{aligned}$$

Then collecting the estimates above, we give the desired estimate and thus complete the proof of the proposition. \square

5. PROOF OF THE THEOREM 1.3

In this section, we are ready to prove Theorem 1.3. To do this, we need the following preliminary.

5.1. Definitions of the modified incoming and outgoing components. With the preparations in the previous sections, we can define the modified incoming and outgoing components, say f_+ and f_- , of the function f . First of all, we split the function f as follows,

$$f = P_{\leq N} f + P_{\geq N} \chi_{\leq 1} f + P_{\geq N} \chi_{\geq 1} f.$$

Definition 5.1. Let the radial function $f \in \mathcal{S}(\mathbb{R}^d)$. We define the modified outgoing component of f as

$$f_+ = \frac{1}{2} P_{\leq N} f + \frac{1}{2} P_{\geq N} \chi_{\leq 1} f + (P_{\geq N} \chi_{\geq 1} f)_{out};$$

the modified incoming component of f as

$$f_- = \frac{1}{2} P_{\leq N} f + \frac{1}{2} P_{\geq N} \chi_{\leq 1} f + (P_{\geq N} \chi_{\geq 1} f)_{in}.$$

From the definitions, we have

$$f = f_+ + f_-.$$

Moreover, Definition 5.1 combining with Proposition 4.5 gives the proof of Proposition 1.1.

To prove Theorem 1.3, we only consider the outgoing part, since the incoming part can be treated in the same way. That is, initial data $u_0 = f_+$.

Denote $v_L = e^{it\Delta} (P_{\geq N} \chi_{\geq 1} f)_{out}$. Let $w = u - v_L$, then w obeys the equation

$$\begin{cases} i\partial_t w + \Delta w = |u|^p u, \\ w(0, x) = w_0(x), \end{cases} \quad (5.1)$$

where

$$w_0 = \frac{1}{2} P_{\leq N} f + \frac{1}{2} P_{\geq N} \chi_{\leq 1} f.$$

Then by Bernstein's inequality, and (1.6) (in which $\varepsilon_0 = 1$), we have

$$\|w_0\|_{H^1(\mathbb{R}^d)} \lesssim N^{1-s_0}, \quad (5.2)$$

where the implicit constant depends on $\|\chi_{\leq 1} f\|_{H^1(\mathbb{R}^d)}$ and $\|\chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)}$.

The first result is the following local well-posedness theorem, in which the indices are not sharp but enough for this paper.

Proposition 5.2. *Let $d = 3, 4, 5$. Then there exists $p_1(d) < \frac{4}{d-2}$, such that for any $s_0 \geq \frac{1}{2}$ and any $p \in [p_1, \frac{4}{d-2})$, the following is true. Let f be the function under the same hypothesis on Theorem 1.3 with $\varepsilon_0 = 1$ and $w_0 \in H^1(\mathbb{R}^d)$, then there exists $T_{max} > 0$, such that the Cauchy problem (5.1) is locally well-posed on $[0, T_{max})$. Moreover, the blowup criterion holds: If $T_{max} < \infty$, then*

$$\lim_{t \rightarrow T_{max}} \|w(t)\|_{H^1(\mathbb{R}^d)} = +\infty.$$

Proof. It follows from the standard fixed point argument, and thus the proof is much brief. According to the Duhamel formula, we denote

$$\Phi(w)(t) = e^{it\Delta}w_0 + \int_0^t e^{i(t-s)\Delta}|u(s)|^p u(s) ds.$$

For short, we further denote the parameters

$$\frac{1}{\tilde{r}} = \frac{1}{2} - \frac{3d-2}{d(2d-1)} - \frac{\varepsilon}{d}; \quad \tilde{s} = 1 - \frac{d-1}{2d-1} - \varepsilon,$$

where ε is an arbitrary small positive constant. Fixing $T > 0$, we denote the norm

$$\|w\| := \|w\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^d)} + \|\langle \nabla \rangle^{\tilde{s}} w\|_{L_t^2 L_x^{\tilde{r}}([0, T] \times \mathbb{R}^d)}.$$

From Lemma 2.9 and Lemma 2.2, we have

$$\begin{aligned} \|\Phi(w)\| &\lesssim \|w_0\|_{H_x^1(\mathbb{R}^d)} + \|\langle \nabla \rangle^{\tilde{s}}(|u|^p u)\|_{L_t^2 L_x^{\frac{2d}{d+2\tilde{s}}}([0, T] \times \mathbb{R}^d)} \\ &\lesssim \|w_0\|_{H_x^1(\mathbb{R}^d)} + \|\langle \nabla \rangle^{\tilde{s}} v_L\|_{L_t^2 L_x^\infty([0, T] \times \mathbb{R}^d)} \left(\|v_L\|_{L_t^\infty L_x^{\frac{2dp}{d+2\tilde{s}}}([0, T] \times \mathbb{R}^d)}^p + \|w\|_{L_t^\infty L_x^{\frac{2dp}{d+2\tilde{s}}}([0, T] \times \mathbb{R}^d)}^p \right) \\ &\quad + \|\langle \nabla \rangle^{\tilde{s}} w\|_{L_t^\infty L_x^{r_1}([0, T] \times \mathbb{R}^d)} \left(\|v_L\|_{L_t^{2p} L_x^{dp}([0, T] \times \mathbb{R}^d)}^p + \|w\|_{L_t^{2p} L_x^{dp}([0, T] \times \mathbb{R}^d)}^p \right), \end{aligned} \quad (5.3)$$

where r_1 is the parameters satisfying

$$\frac{1}{r_1} = \frac{1}{2} - \frac{1}{d}(1 - \tilde{s}).$$

Note that when $p_1(d)$ is suitably close to $\frac{4}{d-2}$, we have $2 < r_1, \frac{2dp}{d+2\tilde{s}} < \frac{2d}{d-2}$. From (4.19)–(4.21), we have that for any $s_0 \geq \frac{1}{2}$,

$$\|\langle \nabla \rangle^{\tilde{s}} v_L\|_{L_t^2 L_x^\infty(\mathbb{R}^+ \times \mathbb{R}^d)} + \|v_L\|_{L_t^\infty L_x^{\frac{2dp}{d+2\tilde{s}}}(\mathbb{R}^+ \times \mathbb{R}^d)} + \|v_L\|_{L_t^{2p} L_x^{dp}(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim \|f\|_{H^{s_0}(\mathbb{R}^d)},$$

Hence, we get

$$\|\langle \nabla \rangle^{\tilde{s}} v_L\|_{L_t^2 L_x^\infty([0, T] \times \mathbb{R}^d)} + \|v_L\|_{L_t^\infty L_x^{\frac{2dp}{d+2\tilde{s}}}([0, T] \times \mathbb{R}^d)} + \|v_L\|_{L_t^{2p} L_x^{dp}([0, T] \times \mathbb{R}^d)} = r(T),$$

where $r(T) \rightarrow 0$ when $T \rightarrow 0$. Further, by the Sobolev inequality and interpolation, we have

$$\begin{aligned} \|w\|_{L_t^\infty L_x^{\frac{2dp}{d+2\tilde{s}}}([0, T] \times \mathbb{R}^d)} + \|\langle \nabla \rangle^{\tilde{s}} w\|_{L_t^\infty L_x^{r_1}([0, T] \times \mathbb{R}^d)} &\lesssim \|w\|_{L_t^\infty H_x^1([0, T] \times \mathbb{R}^d)}; \\ \|w\|_{L_t^{2p} L_x^{dp}([0, T] \times \mathbb{R}^d)} &\lesssim T^\theta \|w\|, \end{aligned}$$

where $\theta = \frac{1}{2}(1 - s_c) > 0$. Hence, inserting these estimates into (5.3), we obtain that

$$\|\Phi(w)\| \lesssim \|w_0\|_{H_x^1(\mathbb{R}^d)} + r(T)(1 + \|w\|^{p+1}).$$

Similarly, by choosing $T = T(\|f\|_{H^{s_0}(\mathbb{R}^d)}, \|w_0\|_{H^1(\mathbb{R}^d)}) > 0$ small enough, we have that for any w_j satisfying $\|w_j\| \leq 2\|w_0\|_{H^1(\mathbb{R}^d)}$ for $j = 1, 2$,

$$\|\Phi(w_1) - \Phi(w_2)\| \leq \frac{1}{2}\|w_1 - w_2\|.$$

Then the proposition follows from the fixed point theory. \square

In the following subsections, we will prove the boundedness of w in the energy space and some space-time spaces. To this end, we define the working space as follows. We denote $X_N(I)$ for $I \subset \mathbb{R}^+$ to be the space under the norms

$$\|h\|_{X_N(I)} := N^{3(s_0-1)}\|h\|_{L_t^\infty \dot{H}_x^1(I \times \mathbb{R}^d)} + N^{\alpha(d,p) \cdot (s_0-1)}\|h\|_{L_{tx}^{r_0}(I \times \mathbb{R}^d)},$$

where the positive constants

$$r_0 = p + 2 + \frac{2}{d-2}, \quad \text{and} \quad \alpha(d,p) = \frac{3d}{r_0(d-2)}.$$

In the following, we restrict our attention to 3, 4 and 5 dimensions. Moreover, fixing $\delta_0 > 0$, we set $N = N(\delta_0) > 0$, such that

$$\|P_{\geq N} \chi_{\geq 1} f\|_{H^{s_0}(\mathbb{R}^d)} \leq \delta_0. \quad (5.4)$$

First of all, we have the uniform boundedness of the $L_t^\infty L_x^2(I \times \mathbb{R}^d)$ norm of w . Indeed, choosing $s_0 > 0$, we have $u_0 \in L^2(\mathbb{R}^d)$. Using the mass conservation law, we have

$$\|u\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} = \|u_0\|_{L^2(\mathbb{R}^d)}.$$

Moreover, by Proposition 3.13,

$$\|v_L\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \lesssim \|f\|_{H^{s_0}(\mathbb{R}^d)}.$$

Hence, we obtain that

$$\|w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \lesssim \|f\|_{H^{s_0}(\mathbb{R}^d)}. \quad (5.5)$$

5.2. Space-time estimates. In this subsection, based on the $L_{tx}^{r_0}$ -norm in $\|w\|_{X_N(I)}$, we shall prove the general space-time estimates we will use below.

Lemma 5.3. *Let $p_1(d) > \max\{\frac{2}{d-2}, 1\}$, then there exists $s_0 \in (0, 1)$, such that for any $p \in [p_1(d), \frac{4}{d-2})$, there exist some constants $\alpha_1(d, p) > 0, \alpha_2(d, p) > 0$ such that*

$$\|w\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)} + \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \lesssim N^{\alpha_1(d,p) \cdot (1-s_0)} (1 + \|w\|_{X_N(I)}^{\alpha_2(d,p)}).$$

Proof. By Lemma 2.8, we have

$$\|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \lesssim \|\langle \nabla \rangle w_0\|_{L_x^2(\mathbb{R}^d)} + \|\langle \nabla \rangle (|u|^p u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)}. \quad (5.6)$$

From (5.2), we have

$$\|\langle \nabla \rangle w_0\|_{L_x^2(\mathbb{R}^d)} \lesssim N^{1-s_0}. \quad (5.7)$$

Moreover, using $u = w + v_L$, we have

$$\begin{aligned} \|\langle \nabla \rangle (|u|^p u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} &\lesssim \|\langle \nabla \rangle v_L\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \left(\|v_L\|_{L_t^\infty L_x^{\frac{dp}{2}}(I \times \mathbb{R}^d)}^p + \|w\|_{L_t^\infty L_x^{\frac{dp}{2}}(I \times \mathbb{R}^d)}^p \right) \\ &\quad + \|\langle \nabla \rangle w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \left(\|v_L\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p + \|w\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p \right). \end{aligned}$$

Since $\frac{dp}{2} > 2$, then by (4.19)–(4.21) and (5.4), we have

$$\|\langle \nabla \rangle v_L\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \lesssim \delta_0 N^{\frac{d+2}{2d} - s_0 +},$$

and

$$\|v_L\|_{L_t^\infty L_x^{\frac{dp}{2}}(I \times \mathbb{R}^d)} \lesssim \delta_0 N^{\frac{1}{2} - \frac{2}{dp} - s_0 +}; \quad \|v_L\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)} \lesssim \delta_0 N^{\frac{1}{2} - \frac{d+2}{2dp} - s_0 +}.$$

Moreover, by the interpolation,

$$\|w\|_{L_t^\infty L_x^{\frac{dp}{2}}(I \times \mathbb{R}^d)} \lesssim \|\nabla w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)}^{s_c} \|w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)}^{1-s_c} \lesssim N^{3(1-s_0)s_c} \|w\|_{X_N(I)}^{s_c}.$$

Hence, choosing $1 - s_0$ small enough such that for any $p \in [1, 4]$,

$$\frac{d+2}{2d} - s_0 + 3s_c p(1-s_0) < 0, \quad \left(\frac{1}{2} - \frac{d+2}{2dp} - s_0\right)p + 3(1-s_0) < 0,$$

we obtain that

$$\begin{aligned} \|\langle \nabla \rangle (|u|^p u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} &\lesssim 1 + \delta_0 (\|w\|_{X_N(I)} + \|w\|_{X_N(I)}^{s_c p}) \\ &\quad + \|\langle \nabla \rangle w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \|w\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p. \end{aligned}$$

Noting that $r_0 > 2p$ and $p > \frac{2}{d-2}$, then by the interpolation, we have

$$\begin{aligned} \|\langle \nabla \rangle w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \|w\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p &\lesssim \|\langle \nabla \rangle w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \|w\|_{L_{tx}^{r_0}(I \times \mathbb{R}^d)}^{\frac{r_0(p-1)}{r_0-2}} \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)}^{\frac{r_0-2p}{r_0-2}} \\ &\lesssim N \left[3 + \frac{r_0(p-1)}{r_0-2} \alpha(d, p) \right] \cdot (1-s_0) \|w\|_{X_N(I)}^{1+\frac{r_0(p-1)}{r_0-2}} \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)}^{\frac{r_0-2p}{r_0-2}}. \end{aligned}$$

Hence, we obtain that

$$\begin{aligned} \|\langle \nabla \rangle (|u|^p u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} &\lesssim 1 + \delta_0 (\|w\|_{X_N(I)} + \|w\|_{X_N(I)}^{s_c p}) \\ &\quad + N^{\widetilde{\alpha}_1(d, p) \cdot (1-s_0)} \|w\|_{X_N(I)}^{\widetilde{\alpha}_2(d, p)} \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)}^{\frac{r_0-2p}{r_0-2}}, \end{aligned}$$

where

$$\widetilde{\alpha}_1(d, p) = 3 + \frac{r_0(p-1)}{r_0-2} \alpha(d, p); \quad \widetilde{\alpha}_2(d, p) = 1 + \frac{r_0(p-1)}{r_0-2}.$$

Combining this estimate with (5.6) and (5.7), we get

$$\begin{aligned} \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} &\lesssim N^{1-s_0} + \delta_0 (\|w\|_{X_N(I)} + \|w\|_{X_N(I)}^{s_c p}) \\ &\quad + N^{\widetilde{\alpha}_1(d, p) \cdot (1-s_0)} \|w\|_{X_N(I)}^{\widetilde{\alpha}_2(d, p)} \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)}^{\frac{r_0-2p}{r_0-2}}. \end{aligned}$$

Note that by setting $p_1(d) > 1$, we have

$$\frac{r_0 - 2p}{r_0 - 2} < 1.$$

Hence, by the Cauchy-Schwarz inequality, we get

$$\|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \lesssim N^{\alpha_1(d, p) \cdot (1-s_0)} (1 + \|w\|_{X_N(I)}^{\alpha_2(d, p)}),$$

where

$$\alpha_1(d, p) = \max \left\{ 1, \widetilde{\alpha}_1(d, p) \frac{r_0 - 2}{2p - 2} \right\}, \quad \alpha_2(d, p) = \max \left\{ 1, s_c p, \widetilde{\alpha}_2(d, p) \frac{r_0 - 2}{2p - 2} \right\}.$$

Moreover, since

$$\|w\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p \lesssim \|\langle \nabla \rangle w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)}^{\frac{r_0 - 2p}{r_0 - 2}} \|w\|_{L_{tx}^{r_0}(I \times \mathbb{R}^d)}^{\frac{r_0(p-1)}{r_0 - 2}},$$

we get the desired estimate by resetting the constants $\alpha_1(d, p), \alpha_2(d, p)$ if necessary. \square

Remark 5.4. Taking $p_1(d) \geq \frac{11}{10}$, then we can roughly estimate that $\alpha_1(d, p) \leq 75, \alpha_2(d, p) \leq 150$. Hence, $\alpha_1(d, p), \alpha_2(d, p)$ have a uniform upper bound when $p_1(d)$ is close to $\frac{4}{d-2}$. Based on these facts, the parameters $\alpha_3(d, p), \alpha_4(d, p)$ defined later also have a uniform upper bound when $p_1(d)$ is close to $\frac{4}{d-2}$.

5.3. Morawetz estimates. In this subsection, we consider the Morawetz-type estimate of Lin-Strauss [35], see also [14] for the interaction Morawetz estimates. For convenience, we rewrite the equation of w in the following way,

$$i\partial_t w + \Delta w = |w|^p w + F(v_L, w),$$

where $F(v_L, w) = |v_L + w|^p(v_L + w) - |w|^p w$. Let

$$M(t) = \operatorname{Im} \int_{\mathbb{R}^d} \frac{x}{|x|} \cdot \nabla w(t, x) \bar{w}(t, x) dx.$$

Then we have the following lemma.

Lemma 5.5. *Under the same assumption as in Lemma 5.3, for any time interval I such that $0 \in I \subset \mathbb{R}^+$,*

$$\int_I \int_{\mathbb{R}^d} \frac{|w(t, x)|^{p+2}}{|x|} dx dt \lesssim N^{3(1-s_0)} \left(\|w\|_{X_N(I)} + \delta_0 \|w\|_{X_N(I)}^{1+p\alpha_2(d,p)} \right).$$

Proof. Note that

$$M'(t) = \operatorname{Im} \int \left(2 \frac{x}{|x|} \cdot \nabla w + \frac{d-1}{|x|} w \right) \bar{w}_t dx. \quad (5.8)$$

Since

$$\bar{w}_t = -i\Delta \bar{w} + i|w|^p \bar{w} + i\overline{F(v_L, w)},$$

we shall consider the following three terms,

$$\operatorname{Im} \int \left(2 \frac{x}{|x|} \cdot \nabla w + \frac{d-1}{|x|} w \right) (-i\Delta \bar{w}) dx; \quad (5.9a)$$

$$\operatorname{Im} \int \left(2 \frac{x}{|x|} \cdot \nabla w + \frac{d-1}{|x|} w \right) (i|w|^p \bar{w}) dx; \quad (5.9b)$$

and

$$\operatorname{Im} \int \left(2 \frac{x}{|x|} \cdot \nabla w + \frac{d-1}{|x|} w \right) \left(i\overline{F(v_L, w)} \right) dx. \quad (5.9c)$$

By a direct computation, we have

$$(5.9a) \geq 0. \quad (5.10)$$

Indeed, by integration-by-parts,

$$(5.9a) = \int \frac{1}{|x|} \left(|\nabla w|^2 - \left| \frac{x}{|x|} \cdot \nabla w \right|^2 \right) dx + G(w),$$

where

$$G(w) = \begin{cases} 2\pi |w(t, 0)|^2, & \text{if } d = 3, \\ \frac{(d-1)(d-3)}{4} \int \frac{|w(t, x)|^2}{|x|^3} dx, & \text{if } d > 3. \end{cases}$$

Hence, we obtain (5.10). Using integration-by-parts again, we find

$$(5.9b) = \frac{(d-1)p}{2(p+1)} \int \frac{|w(t, x)|^{p+2}}{|x|} dx. \quad (5.11)$$

For (5.9c), using the Hölder inequality and the Hardy inequality, we have

$$\begin{aligned} |(5.9c)| &\leq \left| \int \left(2 \frac{x}{|x|} \cdot \nabla w + \frac{d-1}{|x|} w \right) \left(i \overline{F(v_L, w)} \right) dx \right| \\ &\lesssim \|F(v_L, w)\|_{L_x^2(\mathbb{R}^d)} \left(\|\nabla w\|_{L_x^2(\mathbb{R}^d)} + \left\| \frac{w}{|x|} \right\|_{L_x^2(\mathbb{R}^d)} \right) \\ &\lesssim \|\nabla w\|_{L_x^2(\mathbb{R}^d)} \|F(v_L, w)\|_{L_x^2(\mathbb{R}^d)}. \end{aligned}$$

Hence, this last estimate combining (5.10) and (5.11), and integrating in time in (5.8), we obtain that

$$\int_I \int_{\mathbb{R}^d} \frac{|w(t, x)|^{p+2}}{|x|} dx dt \lesssim \max_{t \in I} M(t) + \|\nabla w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \|F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)}. \quad (5.12)$$

By the Hölder inequality and (5.5), we have

$$\max_{t \in I} M(t) \lesssim N^{3(1-s_0)} \|w\|_{X_N(I)}. \quad (5.13)$$

Now we claim that by choosing s_0 close enough to 1,

$$\|F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \lesssim 1 + \delta_0 \|w\|_{X_N(I)}^{p\alpha_2(d,p)}. \quad (5.14)$$

Indeed, we have

$$\|F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \lesssim \|v_L\|_{L_t^1 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \left(\|w\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p + \|v_L\|_{L_t^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p \right).$$

From (4.19), (4.21), (5.4) and Lemma 5.3, it is further bounded by

$$\delta_0 N^{-s_0 - \frac{d-2}{2d} + \left[N^{p\alpha_1(d,p) \cdot (1-s_0)} \left(1 + \|w\|_{X_N(I)}^{p\alpha_2(d,p)} \right) + \delta_0^p N^{-ps_0 + \frac{p}{2} - \frac{d+2}{2d} + \left[\right]} \right]}.$$

Choosing $1 - s_0$ small enough such that for any $p \in [1, 4]$,

$$(1 - s_0) \left[1 + p\alpha_1(d, p) \right] < 1 + \frac{d-2}{2d}, \quad (1 - s_0)(p+1) < \frac{p}{2} + 2,$$

we obtain (5.14).

Therefore, together with (5.12), (5.13) and (5.14), we get

$$\int_I \int_{\mathbb{R}^d} \frac{|w(t, x)|^{p+2}}{|x|} dx dt \lesssim N^{3(1-s_0)} \left(\|w\|_{X_N(I)} + \delta_0 \|w\|_{X_N(I)}^{1+p\alpha_2(d,p)} \right).$$

This finishes the proof of the lemma. \square

The following is a consequence of the previous lemma.

Corollary 5.6. *Under the same assumptions as in Lemma 5.3, there exists $\alpha_3(d, p) > 0$ such that*

$$\|w\|_{L_{tx}^{r_0}(I \times \mathbb{R}^d)} \lesssim N^{\alpha(d, p) \cdot (1-s_0)} \left(\|w\|_{X_N(I)}^{\frac{d}{r_0(d-2)}} + \delta_0^{\frac{1}{r_0}} \|w\|_{X_N(I)}^{\alpha_3(d, p)} \right).$$

Proof. By the Hölder inequality,

$$\int_I \int_{\mathbb{R}^d} |w(t, x)|^{r_0} dx dt \lesssim \| |x|^{\frac{d-2}{2}} w \|_{L_{tx}^\infty(I \times \mathbb{R}^d)}^{\frac{2}{d-2}} \int_I \int_{\mathbb{R}^d} \frac{|w(t, x)|^{p+2}}{|x|} dx dt.$$

From Lemma 2.1, we have

$$\| |x|^{\frac{d-2}{2}} w \|_{L_{tx}^\infty(I \times \mathbb{R}^d)} \lesssim \|\nabla w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \lesssim N^{3(1-s_0)} \|w\|_{X_N(I)}.$$

From Lemma 5.5, we have

$$\int_I \int_{\mathbb{R}^d} \frac{|w(t, x)|^{p+2}}{|x|} dx dt \lesssim N^{3(1-s_0)} \left(\|w\|_{X_N(I)} + \delta_0 \|w\|_{X_N(I)}^{1+p\alpha_2(d, p)} \right).$$

Hence, we obtain

$$\int_I \int_{\mathbb{R}^d} |w(t, x)|^{r_0} dx dt \lesssim N^{\frac{3d}{d-2}(1-s_0)} \left(\|w\|_{X_N(I)}^{\frac{d}{d-2}} + \delta_0 \|w\|_{X_N(I)}^{\frac{d}{d-2} + p\alpha_2(d, p)} \right).$$

Let

$$\alpha_3(d, p) = \frac{1}{r_0} \left[\frac{d}{d-2} + p\alpha_2(d, p) \right],$$

we have the desired estimate. This finishes the proof of the corollary. \square

5.4. Energy estimate. In this subsection, we consider the energy estimate for w . Since the energy of w is not conserved, the nonlinear estimates are needed in this subsection. The main result in this subsection is the following $\dot{H}^1(\mathbb{R}^d)$ -norm bound for w .

Lemma 5.7. *There exists $\alpha_4(d, p) > 1$ such that for any $0 \in I \subset \mathbb{R}^+$,*

$$\sup_{t \in I} E(w(t)) \lesssim N^{6(1-s_0)} \left(1 + \delta_0 \|w\|_{X_N(I)}^{2\alpha_4(d, p)} \right). \quad (5.15)$$

Proof. Let $t \in I$. Taking product with w_t on the equation (5.1) and integrating in space and in time from 0 to t , we get

$$E(w(t)) = E(w_0) + \operatorname{Im} \int_0^t \int_{\mathbb{R}^d} \nabla F(v_L, w) \cdot \nabla w dx dt' + \operatorname{Im} \int_0^t \int_{\mathbb{R}^d} F(v_L, w) |u|^p u dx dt'.$$

Since $p \leq 4$, by (5.2) and the Sobolev inequality, it follows that

$$E(w_0) \lesssim N^{(p+2)(1-s_0)} \lesssim N^{6(1-s_0)}.$$

Then by the Hölder inequality,

$$\begin{aligned} E(w(t)) &\lesssim N^{6(1-s_0)} + \|\nabla w\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \|\nabla F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \\ &\quad + \|F(v_L, w) |u|^p u\|_{L_{tx}^1(I \times \mathbb{R}^d)}. \end{aligned} \quad (5.16)$$

First, we claim that for any $0 \in I \subset \mathbb{R}^+$,

$$\|\nabla F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \lesssim \delta_0 N^{3(1-s_0)} N^{-\beta_0+} (1 + \|w\|_{X_N(I)}^{p\alpha_2(d, p)}), \quad (5.17)$$

where

$$\beta_0 = \min \left\{ s_0 - \frac{1}{2}; \frac{d-2}{2d} - (p\alpha_1(d, p) - 2)(1 - s_0); \frac{p}{2} + \frac{d+2}{d} - (\alpha_1(d, p) + p - 3)(1 - s_0) \right\}.$$

Choosing s_0 close enough to 1, then for any $p \in [1, 4]$, there exists $c(d) > 0$ such that $\beta_0 \geq c(d) > 0$. Since

$$|\nabla F(v_L, w)| \lesssim |w|^p |\nabla v_L| + |v_L|^p |\nabla v_L| + |v_L|^p |\nabla w| + |w|^{p-1} |\nabla w| |v_L|,$$

we have that

$$\begin{aligned} & \|\nabla F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \\ & \lesssim \|w\|_{L_x^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p \|\nabla v_L\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} + \|v_L\|_{L_x^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p \|\nabla v_L\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \\ & \quad + \|v_L\|_{L_x^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^p \|\nabla w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} + \|v_L\|_{L_x^{2p} L_x^{dp}(I \times \mathbb{R}^d)} \|w\|_{L_x^{2p} L_x^{dp}(I \times \mathbb{R}^d)}^{p-1} \|\nabla w\|_{L_t^2 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)}. \end{aligned}$$

Choose s_0 close enough to 1, such that for any $p \in [1, 4]$,

$$1 - s_0 \leq \frac{1}{2}, \quad (p\alpha_1(d, p) - 2) \cdot (1 - s_0) < \frac{d-2}{2d}, \quad (\alpha_1(d, p) + p - 3) \cdot (1 - s_0) < \frac{p}{2} + \frac{d+2}{d},$$

then by (4.19), (4.21), (5.4), Lemma 5.3 and Cauchy-Schwarz's inequality, we get

$$\begin{aligned} & \|\nabla F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \\ & \lesssim \delta_0 N^{p\alpha_1(d, p) \cdot (1-s_0)} N^{-s_0 + \frac{d+2}{2d}} + \left(1 + \|w\|_{X_N(I)}^{p\alpha_2(d, p)}\right) + \delta_0^{p+1} N^{-s_0 + \frac{d+2}{2d}} + N^{-ps_0 + \frac{p}{2} - \frac{d+2}{2d}} \\ & \quad + \delta_0^p N^{-ps_0 + \frac{p}{2} - \frac{d+2}{2d}} + N^{\alpha_1(d, p) \cdot (1-s_0)} \left(1 + \|w\|_{X_N(I)}^{\alpha_2(d, p)}\right) \\ & \quad + \delta_0 N^{-s_0 + \frac{1}{2} - \frac{d+2}{2dp}} + N^{p\alpha_1(d, p) \cdot (1-s_0)} \left(1 + \|w\|_{X_N(I)}^{p\alpha_2(d, p)}\right) \\ & \lesssim \delta_0 N^{3(1-s_0)} N^{-\beta_0 +} \left(1 + \|w\|_{X_N(I)}^{p\alpha_2(d, p)}\right). \end{aligned}$$

Hence we obtain (5.17).

Second, we claim that

$$\|F(v_L, w)|u|^p u\|_{L_{tx}^1(I \times \mathbb{R}^d)} \lesssim N^{6(1-s_0)} \left(1 + \delta_0 \|w\|_{X_N(I)}^{p\alpha_2(d, p) + p + 1}\right). \quad (5.18)$$

Indeed, by (5.17), it follows that

$$\begin{aligned} & \|F(v_L, w)|u|^p u\|_{L_{tx}^1(I \times \mathbb{R}^d)} \\ & \lesssim \|F(v_L, w)\|_{L_t^1 L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \| |u|^p u \|_{L_t^\infty L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} \\ & \lesssim \|\nabla F(v_L, w)\|_{L_t^1 L_x^2(I \times \mathbb{R}^d)} \|u\|_{L_t^\infty L_x^{\frac{2d(p+1)}{d+2}}(I \times \mathbb{R}^d)}^{p+1} \\ & \lesssim \delta_0 N^{3(1-s_0)} N^{-\beta_0 +} \left(1 + \|w\|_{X_N(I)}^{p\alpha_2(d, p)}\right) \left(\|w\|_{L_t^\infty L_x^{\frac{2d(p+1)}{d+2}}(I \times \mathbb{R}^d)}^{p+1} + \|v_L\|_{L_t^\infty L_x^{\frac{2d(p+1)}{d+2}}(I \times \mathbb{R}^d)}^{p+1} \right). \end{aligned}$$

Noting that $2 < \frac{2d(p+1)}{d+2} < \frac{2d}{d-2}$, then by the interpolation, we get

$$\|w\|_{L_t^\infty L_x^{\frac{2d(p+1)}{d+2}}(I \times \mathbb{R}^d)} \lesssim \|w\|_{L_t^\infty H_x^1(I \times \mathbb{R}^d)} \lesssim N^{3(1-s_0)} \|w\|_{X_N(I)}.$$

Moreover, from (4.20), we have

$$\|v_L\|_{L_t^\infty L_x^{\frac{2d(p+1)}{d+2}}(I \times \mathbb{R}^d)} \lesssim \delta_0.$$

Choose s_0 close enough to 1, such that for any $p \in [1, 4]$,

$$3p(1 - s_0) < \beta_0,$$

then collecting the estimates above, we obtain (5.18).

Inserting (5.17) and (5.18) into (5.16), we obtain that

$$E(w(t)) \lesssim N^{6(1-s_0)} \left(1 + \delta_0 \|w\|_{X_N(I)}^{p\alpha_2(d,p)+1} + \delta_0 \|w\|_{X_N(I)}^{p\alpha_2(d,p)+p+1} \right).$$

By Cauchy-Schwarz's inequality and setting

$$\alpha_4(d, p) = \frac{1}{2}(p\alpha_2(d, p) + p + 1),$$

we obtain the desired estimates. \square

5.5. The proofs. Now we are ready to prove Theorem 1.3. First, we show that for any I such that $0 \in I \subset \mathbb{R}^+$,

$$\|w\|_{X_N(I)} \lesssim 1. \quad (5.19)$$

Indeed, from Corollary 5.6 and Lemma 5.7, we have

$$\begin{aligned} & N^{-\alpha(d,p) \cdot (1-s_0)} \|w\|_{L_{tx}^{r_0}(I \times \mathbb{R}^d)} + N^{3(s_0-1)} \|w\|_{L_t^\infty \dot{H}_x^1(I \times \mathbb{R}^d)} \\ & \lesssim 1 + \|w\|_{X_N(I)}^{\frac{d}{r_0(d-2)}} + \delta_0^{\frac{1}{r_0}} \|w\|_{X_N(I)}^{\alpha_3(d,p)} + \delta_0^{\frac{1}{2}} \|w\|_{X_N(I)}^{\alpha_4(d,p)}. \end{aligned}$$

That is,

$$\|w\|_{X_N(I)} \lesssim 1 + \|w\|_{X_N(I)}^{\frac{d}{r_0(d-2)}} + \delta_0^{\frac{1}{r_0}} \|w\|_{X_N(I)}^{\alpha_3(d,p)} + \delta_0^{\frac{1}{2}} \|w\|_{X_N(I)}^{\alpha_4(d,p)}.$$

We note that

$$\frac{d}{r_0(d-2)} < 1, \quad \alpha_4(d, p) > 1.$$

Then using the Cauchy-Schwarz inequality, the second term can be absorbed by the term in the left-hand side, and thus we have

$$\|w\|_{X_N(I)} \lesssim 1 + \delta_0^{\frac{1}{r_0}} \|w\|_{X_N(I)}^{\alpha_3(d,p)} + \delta_0^{\frac{1}{2}} \|w\|_{X_N(I)}^{\alpha_4(d,p)}.$$

Furthermore, if $\alpha_3(d, p) \leq 1$, then choosing δ_0 suitably small, we have

$$\|w\|_{X_N(I)} \lesssim 1 + \delta_0^{\frac{1}{2}} \|w\|_{X_N(I)}^{\alpha_4(d,p)}.$$

Then using a continuity argument, we obtain (5.19). If $\alpha_3(d, p) > 1$, then choosing δ_0 suitably small, and using a continuity argument, we also obtain (5.19).

Since the estimate in (5.19) is uniform in the time interval I , we have $I = \mathbb{R}^+$. This proves the global existence in the forward time. Moreover,

$$\|w\|_{L_t^\infty \dot{H}_x^1(\mathbb{R}^+ \times \mathbb{R}^d)} + \|w\|_{L_{tx}^{r_0}(\mathbb{R}^+ \times \mathbb{R}^d)} \lesssim A(N), \quad (5.20)$$

for some constant A depending on N .

Now we set

$$u^+ = f_+ + \int_0^{+\infty} e^{-is\Delta} (|u|^p u) ds.$$

Then we have

$$u(t) - e^{it\Delta} u^+ = \int_t^{+\infty} e^{i(t-s)\Delta} (|u|^p u) ds.$$

Using Proposition 4.5 and (5.20), we have (the details are omitted here since similar treatment was presented above)

$$\left\| \int_t^{+\infty} e^{i(t-s)\Delta} (|u|^p u) ds \right\|_{H^1(\mathbb{R}^d)} \lesssim \|\langle \nabla \rangle (|u|^p u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}([t, +\infty) \times \mathbb{R})} \rightarrow 0,$$

as $t \rightarrow +\infty$. This proves the scattering statement.

ACKNOWLEDGEMENTS

Part of this work was done while M. Beceanu, A. Soffer and Y. Wu were visiting CCNU (C.C.Normal University) Wuhan, China. The authors thank the institutions for their hospitality and the support. M.B. is partially supported by NSF grant DMS 1700293. Q.D. is partially supported by NSFC 11661061, 11671163 and 11771165, and the Fundamental Research Funds for the Central Universities (CCNU18QN030, CCNU18CXTD04). A.S. is partially supported by NSF grant DMS 01600749. The research was also supported by NSFC 11971191 and 11671163. Y.W. is partially supported by NSFC 11771325 and National Youth Topnotch Talent Support Program in China, and A.S. and Y.W. would like to thank Ch. Miao for useful discussions.

REFERENCES

- [1] M. Beceanu, Q. Deng, A. Soffer and Y. Wu, *Large global solutions for nonlinear Schrödinger equations I, mass-subcritical cases*, arXiv:1809.09831.
- [2] M. Beceanu, and A. Soffer, *Large outgoing solutions to supercritical wave equations*, arXiv:1601.06335v1.
- [3] M. Beceanu, and A. Soffer, *Large initial data global well-posedness for a supercritical wave equation*, arXiv:1602.08163
- [4] J. Bourgain, *Periodic nonlinear Schrödinger equation and invariant measures*, Comm. Math. Phys. 166 (1994), no. 1, 1–26.
- [5] J. Bourgain, *Invariant measures for the 2D-defocusing nonlinear Schrödinger equation*, Comm. Math. Phys. 176 (1996), no. 2, 421–445.
- [6] J. Bourgain, *Refinements of Strichartz inequality and applications to 2d-NLS with critical nonlinearity*, Int. Math. Res. Not. (1998), 253–284.
- [7] J. Bourgain, *Global Solutions of Nonlinear Schrödinger Equations*, American Mathematical Society.
- [8] J. Bourgain and D. Li, *On an endpoint Kato-Ponce inequality*, Differential Integral Equations 27 (2014), no. 11-12, 1037–1072.
- [9] A. Bulut, *The defocusing energy-supercritical cubic nonlinear wave equation in dimension five*, arXiv: 1112.0629.
- [10] T. Cazenave, *Semilinear Schrödinger Equations*, Courant Lecture Notes in Mathematics, 10, American Mathematical Society, 2003.
- [11] T. Cazenave and F. Weissler, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* , Nonlinear Anal., Theory, Methods and Applications, **14** (1990), 807–836.
- [12] Y. Cho, S. Lee, *Strichartz estimates in spherical coordinates*. Indiana Univ. Math. J., 62 (2013), no.3, 991–1020.
- [13] M. Christ, J. Colliander, and T. Tao, *Ill-posedness for nonlinear Schrödinger and wave equations*, arXiv:math/0311048.

- [14] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on \mathbb{R}^3* , Comm. Pure Appl. Math. 57 (2004), no. 8, 987–1014.
- [15] J. Colliander, and T. Oh, *Almost sure well-posedness of the cubic nonlinear Schrödinger equation below $L^2(T)$* , Duke Math. J. 161 (2012), no. 3, 367–414.
- [16] B. Dodson, J. Lührmann, and D. Mendelson, *Almost sure scattering for the 4D energy-critical defocusing nonlinear wave equation with radial data*, to appear in Amer. J. Math.
- [17] B. Dodson, C. Miao, J. Murphy, and J. Zheng, *The defocusing quintic NLS in four space dimensions*, Ann. Inst. H. Poincaré Anal. Non Linéaire 34 (2017), no. 3, 759–787.
- [18] T. Duyckaerts, J. Holmer, S. Roudenko, *Scattering for the non-radial 3D cubic nonlinear Schrödinger equation*, Math. Res. Lett. 15 (2008), no. 6, 1233–1250.
- [19] T. Duyckaerts, C. Kenig and F. Merle, *Scattering for radial, bounded solutions of focusing supercritical wave equations*, International Mathematics Research Notices 2014 (2014), 224–258.
- [20] T. Duyckaerts, and T. Roy, *Blow-up of the critical Sobolev norm for nonscattering radial solutions of supercritical wave equations on \mathbb{R}^3* , arXiv:1506.00788 (2015).
- [21] D. Ferreira, W. Staubach, *Global and local regularity for Fourier integral operators on weighted and unweighted spaces*, Mem. Amer. Math. Soc., 229 (2014), Memo 1074.
- [22] J. Ginibre and G. Velo, *Scattering theory in the energy space for a class of nonlinear Schrödinger equations*, J. Math. Pures Appl. 64 (1985), 363–401.
- [23] Z. Guo and Y. Wang, *Improved Strichartz estimates for a class of dispersive equations in the radial case and their applications to nonlinear Schrödinger and wave equations*, J. Anal. Math. 124 (2014), 1–38.
- [24] M. Keel and T. Tao, *Endpoint Strichartz Estimates*, Amer. J. Math. 120 (1998), 955–980.
- [25] C. Kenig, F. Merle, *Scattering for $\dot{H}^{\frac{1}{2}}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions*, Trans. Amer. Math. Soc. 362 (2010), 1937–1962.
- [26] C. Kenig, F. Merle, *Nondispersive radial solutions to energy supercritical non-linear wave equations, with applications*, Amer. J. Math., 133 (2011), 1029–1065.
- [27] C. E. Kenig, G. Ponce, and L. Vega, *On the ill-posedness of some canonical dispersive equations*, Duke Math. J. 106 (2001), no. 3, 617–633.
- [28] R. Killip, D. Li, M. Visan and X. Zhang, *Characterization of minimal-mass blowup solutions to the focusing mass-critical NLS*, SIAM J. Math. Anal. 41 (2009), 219–236.
- [29] R. Killip, S. Masaki, J. Murphy, and M. Visan, *The radial mass-subcritical NLS in negative order Sobolev spaces*, arXiv:1804.06753v1.
- [30] R. Killip, J. Murphy, and M. Visan, *Almost sure scattering for the energy-critical NLS with radial data below $H^1(\mathbb{R}^4)$* , arXiv:1707.09051.
- [31] R. Killip, T. Tao, and M. Visan, *The cubic nonlinear Schrödinger equation in two dimensions with radial data*, J. Eur. Math. Soc. (JEMS) 11 (2009), no. 6, 1203–1258.
- [32] D. Li, *On Kato-Ponce and fractional Leibniz*, Rev. Mat. Iberoam., 35 (2019), no. 1, 23–100.
- [33] D. Li, X. Zhang, *Regularity of almost periodic modulo scaling solutions for mass-critical NLS and applications*, Anal. PDE, 3 (2010), no 2, 175–195.
- [34] D. Li, X. Zhang, *On the rigidity of solitary waves for the focusing mass-critical NLS in dimensions $d \geq 2$* , Sci. China Math. 55 (2012), no. 2, 385–434.
- [35] J. E. Lin, W. A. Strauss, *Decay and scattering of solutions of a nonlinear Schrödinger equation*, J. Funct. Anal. 30 (1978), no. 2, 245–263.
- [36] C. Miao, J. Murphy and J. Zheng, *The defocusing energy-supercritical NLS in four space dimensions*, J. Funct. Anal. 267 (2014), no. 6, 1662–1724.
- [37] C. Miao, Y. Wu and X. Zhang, *The defocusing energy-supercritical nonlinear wave equation when $d = 4$* , preprint.
- [38] S. Miao, *On large future-global-in-time solutions to energy-supercritical nonlinear wave equation*, Non-linear dispersive waves and fluids, 187–214, Contemp. Math., 725, Amer. Math. Soc., Providence, RI, 2019.
- [39] E. Mourre, *Absence of singular continuous spectrum for certain selfadjoint operators*, Comm. Math. Phys. 78 (1980), no. 3, 391–408.
- [40] A. Nahmod, and G. Staffilani, *Almost sure well-posedness for the periodic 3D quintic nonlinear Schrödinger equation below the energy space*, J. Eur. Math. Soc. 17 (2015), no. 7, 1687–1759.

- [41] T. Oh, M. Okamoto, and O. Pocovnicu, *On the probabilistic well-posedness of the nonlinear Schrödinger equations with non-algebraic nonlinearities*, arXiv:1708.01568.
- [42] S. Shao, *Sharp linear and bilinear restriction estimates for paraboloids in the cylindrically symmetric case*, Rev. Mat. Iberoam., 25 (2009), 1127–1168.
- [43] E. Stein, *Harmonic Analysis: Real-variable Methods, Orthogonality and Oscillatory Integrals*. Princeton University Press. (1993).
- [44] T. Tao, *On the asymptotic behavior of large radial data for a focusing non-linear Schrödinger equation*, Dynamics of PDE, V.1 (2004), 1-47,
- [45] T. Tao, M. Visan, X. Zhang, *Global well-posedness and scattering for the defocusing mass-critical nonlinear Schrödinger equation for radial data in high dimensions*, Duke Math. J., 140 (2007), no. 1, 165–202.
- [46] L. Thomann, *Random data Cauchy problem for supercritical Schrödinger equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire 26 (2009), no. 6, 2385–2402.
- [47] I. M. Sigal and A. Soffer, *The N -particle scattering problem: asymptotic completeness for short-range systems*, Annals of Mathematics, Second Series, Vol. 126, No. 1 (1987), 35–108.
- [48] A. Soffer, *Monotonic local decay estimates*, arXiv:1110.6549.
- [49] J. Xie and D. Fang, *Global well-posedness and scattering for the defocusing \dot{H}^s -critical NLS*, Chin. Ann. Math. Ser. B 34 (2013), no. 6, 801–842.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY AT ALBANY SUNY, EARTH SCIENCE
110, ALBANY, NY, 12222, USA,

Email address: mbeceanu@albany.edu

DEPARTMENT OF MATHEMATICS, HUBEI KEY LABORATORY OF MATHEMATICAL SCIENCE, CENTRAL
CHINA NORMAL UNIVERSITY, WUHAN 430079, CHINA.,

Email address: dengq@mail.ccnu.edu.cn

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, 110 FRELINGHUYSEN RD., PISCATAWAY,
NJ, 08854, USA,

DEPARTMENT OF MATHEMATICS, HUBEI KEY LABORATORY OF MATHEMATICAL SCIENCE, CENTRAL
CHINA NORMAL UNIVERSITY, WUHAN 430079, CHINA.,

Email address: soffer@math.rutgers.edu

CENTER FOR APPLIED MATHEMATICS, TIANJIN UNIVERSITY, TIANJIN 300072, CHINA

Email address: yerfmath@gmail.com