

Sun's log-concavity conjecture on the Catalan-Larcombe-French sequence

James J.Y. Zhao

Center for Applied Mathematics
Tianjin University, Tianjin 300072, P. R. China

Email: jjyzhao@tju.edu.cn

Abstract

Let $\{P_n\}_{n \geq 0}$ denote the Catalan-Larcombe-French sequence, which naturally came from the series expansion of the complete elliptic integral of the first kind. In this paper, we prove the strict log-concavity of the sequence $\{\sqrt[n]{P_n}\}_{n \geq 1}$, which was originally conjectured by Z.W. Sun. We also obtain the strict log-concavity of the sequence $\{\sqrt[n]{V_n}\}_{n \geq 1}$, where $\{V_n\}_{n \geq 0}$ is the Fennessey-Larcombe-French sequence arising from the series expansion of the complete elliptic integral of the second kind.

AMS Classification 2010: Primary 05A20

Keywords: The Catalan-Larcombe-French sequence, the Fennessey-Larcombe-French sequence, log-concavity, three-term recurrence relation

1 Introduction

The main purpose of this paper is to prove the log-concavity of the sequences $\{\sqrt[n]{P_n}\}_{n \geq 1}$ and $\{\sqrt[n]{V_n}\}_{n \geq 1}$, where $\{P_n\}_{n \geq 0}$ and $\{V_n\}_{n \geq 0}$, known as the Catalan-Larcombe-French sequence and the Fennessey-Larcombe-French sequence respectively, are given by

$$(n+1)^2 P_{n+1} = 8(3n^2 + 3n + 1)P_n - 128n^2 P_{n-1}, \quad (1.1)$$

$$n(n+1)^2 V_{n+1} = 8n(3n^2 + 5n + 1)V_n - 128(n-1)(n+1)^2 V_{n-1}, \quad (1.2)$$

with the initial values $P_0 = V_0 = 1$ and $P_1 = V_1 = 8$. These two sequences came naturally from the series expansions of the complete elliptic integrals. For more information on $\{P_n\}_{n \geq 0}$ and $\{V_n\}_{n \geq 0}$, see [1, 3, 4, 5, 6].

Let us first give an overview of some background. Recall that a sequence $\{a_n\}_{n \geq 0}$ of real numbers is said to be log-concave (resp. log-convex) if

$$a_n^2 \geq a_{n-1}a_{n+1} \quad (\text{resp. } a_n^2 \leq a_{n-1}a_{n+1})$$

for all $n \geq 1$, and it is strictly log-concave (resp. strictly log-convex) if the inequality is strict. Clearly, for some integer $N > 0$, the positive sequence $\{a_n\}_{n \geq N}$ is log-concave (resp. log-convex) if and only if the sequence $\{a_{n+1}/a_n\}_{n \geq N}$ is decreasing (resp. increasing).

The strict log-concavity of $\{\sqrt[n]{P_n}\}_{n \geq 1}$ was first conjectured by Sun [10], who also conjectured that the sequences $\{P_{n+1}/P_n\}_{n \geq 0}$ and $\{\sqrt[n]{P_n}\}_{n \geq 1}$ are strictly increasing. In fact, motivated by Firoozbakht's conjecture on the strictly decreasing property of the sequence $\{\sqrt[n]{p_n}\}_{n \geq 1}$, where p_n is the n -th prime number (see [7, p.185]), Sun [10] studied the monotonicity of many number theoretical and combinatorial sequences.

The strictly increasing property of the sequence $\{P_{n+1}/P_n\}_{n \geq 0}$ and $\{\sqrt[n]{P_n}\}_{n \geq 1}$ has been confirmed by Xia and Yao [12], and independently by Zhao [14]. We would like to point out that Zhao [14] only proved the log-convexity of $\{P_n\}_{n \geq 0}$, which also implies the monotonicity of $\{\sqrt[n]{P_n}\}_{n \geq 1}$ by using a result due to Wang and Zhu [11].

For a positive sequence $\{a_n\}_{n \geq 0}$ satisfying a three-term recurrence relation, Chen, Guo and Wang [2] obtained a useful criterion to determine the log-concavity of $\{\sqrt[n]{a_n}\}_{n \geq N}$ for some positive integer N . While, their criterion does not apply to the Catalan-Larcombe-French sequence, namely not on Sun's conjecture. The first main result of this paper is as follows.

Theorem 1.1. *The sequence $\{\sqrt[n]{P_n}\}_{n \geq 1}$ is strictly log-concave, that is, for $n \geq 2$,*

$$\left(\sqrt[n]{P_n}\right)^2 > \sqrt[n-1]{P_{n-1}} \cdot \sqrt[n+1]{P_{n+1}}. \quad (1.3)$$

Since the Fennessey-Larcombe-French sequence $\{V_n\}_{n \geq 0}$ is closely related to the sequence $\{P_n\}_{n \geq 0}$, we are led to study the log-behavior of $\{\sqrt[n]{V_n}\}_{n \geq 1}$. The second main result of this paper is as follows.

Theorem 1.2. *The sequence $\{\sqrt[n]{V_n}\}_{n \geq 1}$ is strictly log-concave, that is, for $n \geq 2$,*

$$\left(\sqrt[n]{V_n}\right)^2 > \sqrt[n-1]{V_{n-1}} \cdot \sqrt[n+1]{V_{n+1}}. \quad (1.4)$$

This paper is organized as follows. In Section 2, we give a proof of Theorem 1.1 by establishing a lower bound and an upper bound for the ratio P_n/P_{n-1} . The proof of Theorem 1.2 is similar to that of Theorem 1.1, which will be given in Section 3. Finally, we give a new proof of the monotonicity of $\{\sqrt[n]{P_n}\}_{n \geq 1}$ by using Theorem 1.1, and obtain the monotonicity of $\{\sqrt[n]{V_n}\}_{n \geq 1}$ by using Theorem 1.2.

2 Proof of Theorem 1.1

In this section we aim to prove Theorem 1.1. To this end, we first establish a lower bound and an upper bound for the ratio P_n/P_{n-1} that will lead to the strict log-concavity of the sequence $\{\sqrt[n]{P_n}\}_{n \geq 1}$.

Lemma 2.1. *For any integer $n \geq 1$, let*

$$f(n) = \frac{16(n-1)}{n}. \quad (2.1)$$

Then for $n \geq 5$, we have

$$f(n-1) < \frac{P_n}{P_{n-1}} < f(n). \quad (2.2)$$

Proof For notational convenience, let $v(n) = P_n/P_{n-1}$. We first use induction on n to prove $v(n) > f(n-1)$ for $n \geq 5$. By the recurrence (1.1), we have

$$v(n+1) = \frac{8(3n^2 + 3n + 1)}{(n+1)^2} - \frac{128n^2}{(n+1)^2v(n)}, \quad n \geq 1, \quad (2.3)$$

with the initial value $v(1) = 8$. Clearly, $v(5) = 2152/169 > 12 = f(4)$. Assume that $v(n) > f(n-1)$, and we proceed to prove that $v(n+1) > f(n)$ for $n \geq 5$. Observe that

$$\begin{aligned} v(n+1) - f(n) &= \frac{8(3n^2 + 3n + 1)}{(n+1)^2} - \frac{128n^2}{(n+1)^2v(n)} - \frac{16(n-1)}{n} \\ &= \frac{8(n^3 + n^2 + 3n + 2)v(n) - 128n^3}{n(n+1)^2v(n)}. \end{aligned}$$

By the induction hypothesis, we have $v(n) > f(n-1) > 0$ for $n \geq 5$, therefore

$$v(n+1) - f(n) > \frac{8(n^3 + n^2 + 3n + 2)f(n-1) - 128n^3}{n(n+1)^2v(n)} = \frac{128(n^2 - 4n - 4)}{(n-1)n(n+1)^2v(n)} > 0$$

for $n \geq 5$, since $n^2 - 4n - 4 = (n+1)(n-5) + 1 > 0$. This proves that $v(n) > f(n-1)$ for $n \geq 5$. Moreover, it is clear that $v(n)$ is positive for $n \geq 1$.

The inequality $v(n) < f(n)$ for $n \geq 5$ can be obtained in the same manner, so the detailed proof is omitted here. This completes the proof. \square

With the bounds given in Lemma 2.1 we are now able to prove Theorem 1.1.

Proof of Theorem 1.1 Note that for $n \geq 2$, the inequality (1.3) can be rewritten as

$$\frac{\sqrt[n]{P_n}}{\sqrt[n-1]{P_{n-1}}} > \frac{\sqrt[n+1]{P_{n+1}}}{\sqrt[n]{P_n}},$$

or equivalently,

$$\left(\frac{P_n}{P_{n-1}}\right)^{n(n+1)} > P_n^2 \left(\frac{P_{n+1}}{P_n}\right)^{n(n-1)}. \quad (2.4)$$

By the recurrence (1.1), it is easy to verify that (2.4) holds for $2 \leq n \leq 6$. We proceed to prove that (2.4) is true for $n \geq 7$. By Lemma 2.1 we have

$$\left(\frac{P_n}{P_{n-1}}\right)^{n(n+1)} - P_n^2 \left(\frac{P_{n+1}}{P_n}\right)^{n(n-1)} > (f(n-1))^{n(n+1)} - P_n^2 (f(n+1))^{n(n-1)}$$

$$\begin{aligned}
&= \left(\frac{16(n-2)}{n-1} \right)^{n(n+1)} - P_n^2 \left(\frac{16n}{n+1} \right)^{n(n-1)} \\
&= 16^{n(n-1)} \left(16^{2n} \left(\frac{n-2}{n-1} \right)^{n(n+1)} - P_n^2 \left(\frac{n}{n+1} \right)^{n(n-1)} \right).
\end{aligned}$$

It suffices to prove that

$$16^{2n} \left(\frac{n-2}{n-1} \right)^{n(n+1)} - P_n^2 \left(\frac{n}{n+1} \right)^{n(n-1)} > 0,$$

or equivalently

$$\frac{P_n^2}{16^{2n}} < \left(\frac{(n-2)(n+1)}{(n-1)n} \right)^{n(n-1)} \left(\frac{n-2}{n-1} \right)^{2n}.$$

Thus we only need to show that, for $n \geq 7$,

$$\frac{P_n}{16^n} < \left(\frac{(n-2)(n+1)}{(n-1)n} \right)^{\frac{n(n-1)}{2}} \left(\frac{n-2}{n-1} \right)^n. \quad (2.5)$$

Let l_n denote the term on the left hand side, and r_n denote the term on the right hand side. We claim that

- (i) the sequence $\{l_n\}_{n \geq 5}$ is strictly decreasing, and
- (ii) the sequence $\{r_n\}_{n \geq 5}$ is strictly increasing.

By Lemma 2.1, we see that

$$0 < \frac{l_n}{l_{n-1}} = \frac{P_n/P_{n-1}}{16} < 1$$

for $n \geq 5$, which implies (i).

We proceed to prove (ii). Note that

$$r_n = \left(1 - \frac{1}{\binom{n}{2}} \right)^{\binom{n}{2}} \cdot \left(1 - \frac{1}{n-1} \right)^{n-1} \cdot \left(1 - \frac{1}{n-1} \right).$$

The increasing property of the sequence $\{r_n\}_{n \geq 5}$ immediately follows from the well-known fact that the sequence $\left\{ \left(1 - \frac{1}{n} \right)^n \right\}_{n \geq 1}$ is strictly increasing.

It is easy to verify that $l_7 < r_7$. Combining (i) and (ii), we get that $l_n < r_n$ for $n \geq 7$, namely (2.5) holds. This completes the proof. \square

3 Proof of Theorem 1.2

In this section, we complete the proof of Theorem 1.2 in a similar way of that of Theorem 1.1. For this purpose, we need a lower bound and an upper bound for the ratio V_n/V_{n-1} . For integer $n \geq 2$, let

$$h(n) = \frac{16(n^3 - n^2 + 1)}{n^3 - n^2}, \quad (3.1)$$

which was introduced by Yang and Zhao [13] in their study of the log-concavity of the sequence $\{V_n\}_{n \geq 1}$. They [13] showed that $h(n)$ is a lower bound for the ratio V_n/V_{n-1} , precisely,

$$\frac{V_n}{V_{n-1}} > h(n), \quad (3.2)$$

for $n \geq 4$. We further show that $h(n-1)$ is an upper bound for the ratio V_n/V_{n-1} .

Lemma 3.1. *Let $h(n)$ be given by (3.1). Then for $n \geq 11$, we have*

$$\frac{V_n}{V_{n-1}} < h(n-1). \quad (3.3)$$

Proof Let $g(n) = V_n/V_{n-1}$, by (1.2) it is clear that

$$g(n+1) = \frac{8(3n^2 + 5n + 1)}{(n+1)^2} - \frac{128(n-1)}{ng(n)}, \quad n \geq 1, \quad (3.4)$$

with $g(1) = 8$. The inequality (3.3) can be proved inductively based on the recurrence (3.4). The proof is similar to that of Lemma 2.1, and hence is omitted here. \square

We proceed to prove Theorem 1.2.

Proof of Theorem 1.2 Note that for $n \geq 2$, the inequality (1.4) can be rewritten as

$$\frac{\sqrt[n]{V_n}}{\sqrt[n-1]{V_{n-1}}} > \frac{\sqrt[n+1]{V_{n+1}}}{\sqrt[n]{V_n}},$$

or equivalently,

$$\left(\frac{V_n}{V_{n-1}}\right)^{n(n+1)} > V_n^2 \left(\frac{V_{n+1}}{V_n}\right)^{n(n-1)}. \quad (3.5)$$

By the recurrence (1.2), it is easy to verify that (3.5) holds for $2 \leq n \leq 9$. We proceed to prove that (3.5) is true for $n \geq 10$. By (3.2) we have

$$\frac{V_n}{V_{n-1}} > h(n),$$

for $n \geq 4$. By Lemma 3.1, we have

$$\frac{V_{n+1}}{V_n} < h(n),$$

for $n \geq 10$. Then for $n \geq 10$, it follows that

$$\begin{aligned} \left(\frac{V_n}{V_{n-1}}\right)^{n(n+1)} - V_n^2 \left(\frac{V_{n+1}}{V_n}\right)^{n(n-1)} &> (h(n))^{n(n+1)} - V_n^2 (h(n))^{n(n-1)} \\ &= (h(n))^{n(n-1)} ((h(n))^n + V_n) ((h(n))^n - V_n). \end{aligned}$$

Clearly, both $(h(n))^{n(n-1)}$ and $((h(n))^n + V_n)$ are positive for $n \geq 10$. Thus it suffices to show that for $n \geq 10$,

$$(h(n))^n - V_n > 0. \quad (3.6)$$

Note that the relation (24) in [5] gave an upper bound of V_n , that is, for $n \geq 1$,

$$V_n < (2n+1) \binom{2n}{n}^2. \quad (3.7)$$

Sasvári [9, Corollary 1] showed that for $n \geq 1$,

$$\binom{2n}{n} < \frac{4^n}{\sqrt{\pi n}} e^{-\frac{1}{8n} + \frac{1}{192n^3}}. \quad (3.8)$$

It is clear that for $n \geq 1$,

$$0 < e^{-\frac{1}{8n} + \frac{1}{192n^3}} < 1.$$

Then it follows from (3.8) that for $n \geq 1$

$$\binom{2n}{n} < \frac{4^n}{\sqrt{\pi n}}. \quad (3.9)$$

Combining (3.7) and (3.9), for $n \geq 1$ we have

$$V_n < \frac{2n+1}{\pi n} 16^n < 16^n. \quad (3.10)$$

Note that for $n \geq 2$

$$h(n) = \frac{16(n^3 - n^2 + 1)}{n^3 - n^2} > 16. \quad (3.11)$$

Thus by (3.10) and (3.11), we obtain (3.6). This completes the proof. \square

4 The monotonicity of $\{\sqrt[n]{P_n}\}_{n \geq 1}$ and $\{\sqrt[n]{V_n}\}_{n \geq 1}$

In this section, we aim to derive the monotonicity of both $\{\sqrt[n]{P_n}\}_{n \geq 1}$ and $\{\sqrt[n]{V_n}\}_{n \geq 1}$ from their log-concavity obtained above. The main result of this section is as follows.

Proposition 4.1. *Both $\{\sqrt[n]{P_n}\}_{n \geq 1}$ and $\{\sqrt[n]{V_n}\}_{n \geq 1}$ are strictly increasing.*

Proof Recall that for a real sequence $\{a_n\}_{n \geq 0}$ with positive numbers, it was shown that

$$\liminf_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} \leq \liminf_{n \rightarrow \infty} \sqrt[n]{a_n},$$

and

$$\limsup_{n \rightarrow \infty} \sqrt[n]{a_n} \leq \limsup_{n \rightarrow \infty} \frac{a_{n+1}}{a_n},$$

see Rudin [8, §3.37]. These two inequalities imply a well-known criterion, that is, if $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = c$, then $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = c$, where c is a real number.

Note that it was proved in [5, Eq. (30)] that

$$\lim_{n \rightarrow \infty} \frac{P_n}{P_{n-1}} = \lim_{n \rightarrow \infty} \frac{V_n}{V_{n-1}} = 16.$$

Then it follows that

$$\lim_{n \rightarrow \infty} \sqrt[n]{P_n} = \lim_{n \rightarrow \infty} \sqrt[n]{V_n} = 16.$$

Consequently,

$$\lim_{n \rightarrow \infty} \sqrt[n+1]{P_{n+1}} / \sqrt[n]{P_n} = \lim_{n \rightarrow \infty} \sqrt[n+1]{V_{n+1}} / \sqrt[n]{V_n} = 1.$$

By Theorems 1.1 and 1.2, we see that both $\{\sqrt[n+1]{P_{n+1}} / \sqrt[n]{P_n}\}_{n \geq 1}$ and $\{\sqrt[n+1]{V_{n+1}} / \sqrt[n]{V_n}\}_{n \geq 1}$ are strictly decreasing. Thus for all $n \geq 1$, we have

$$\sqrt[n+1]{P_{n+1}} / \sqrt[n]{P_n} > 1, \quad \text{and} \quad \sqrt[n+1]{V_{n+1}} / \sqrt[n]{V_n} > 1.$$

This completes the proof. \square

Remark. By employing a criterion due to Wang and Zhu [11, Theorem 2.1], Yang and Zhao [13] showed that the sequence $\{\sqrt[n]{V_{n+1}}\}_{n \geq 1}$ is strictly decreasing.

Acknowledgements. I would like to thank the referees for their helpful comments. This work was supported by the 863 Program and the National Science Foundation of China.

References

- [1] Catalan, E.: Sur les Nombres de Segner. *Rend. Circ. Mat. Palermo.*, **1**, 190–201 (1887)

- [2] Chen, W. Y. C., Guo, J. J. F., Wang, L. X. W.: Infinitely log-monotonic combinatorial sequences. *Adv. Appl. Math.*, **52**, 99–120 (2014)
- [3] Jarvis, F., Verrill, H. A.: Supercongruences for the Catalan-Larcombe-French numbers. *Ramanujan J.*, **22**, 171–186 (2010)
- [4] Larcombe, P. J., French, D. R.: On the “other” Catalan numbers: a historical formulation re-examined. *Congr. Numer.*, **143**, 33–64 (2000)
- [5] Larcombe, P. J., French, D. R., Fennessey, E. J.: The Fennessey-Larcombe-French sequence $\{1, 8, 144, 2432, 40000, \dots\}$: formulation and asymptotic form. *Congr. Numer.*, **158**, 179–190 (2002)
- [6] Larcombe, P. J., French, D. R., Fennessey, E. J.: The Fennessey-Larcombe-French sequence $\{1, 8, 144, 2432, 40000, \dots\}$: a recursive formulation and prime factor decomposition. *Congr. Numer.*, **160**, 129–137 (2003)
- [7] Ribenboim, P.: The Little Book of Bigger Primes, Second Edition, Springer-Verlag, New York, 2004
- [8] Rudin, W.: Principles of Mathematical Analysis, Third Edition, McGraw-Hill, Inc., New York, 1976
- [9] Sasvári, Z.: Inequalities for Binomial Coefficients. *J. Math. Anal. Appl.*, **236**, 223–226 (1999)
- [10] Sun, Z. W.: Conjectures involving arithmetical sequences, in Numbers Theory: Arithmetic in Shangri-La (eds., S. Kanemitsu, H. Li and J. Liu), Proc. 6th China-Japan Seminar (Shanghai, August 15–17, 2011), World Sci., Singapore, 2013, pp. 244–258. arXiv: 1208.2683v9.
- [11] Wang, Y., Zhu, B. X.: Proofs of some conjectures on monotonicity of number-theoretic and combinatorial sequences. *Sci. China Math.*, **57(11)**, 2429–2435 (2014)
- [12] Xia, E. X. W., Yao, O. X. M.: A criterion for the log-convexity of combinatorial sequences. *Electron. J. Combin.*, **20(4)**, #P3 (2013)
- [13] Yang, A. L. B., Zhao, J. J. Y.: Log-concavity of the Fennessey-Larcombe-French Sequence. preprint, arXiv: 1503.02151v1
- [14] Zhao, F.-Z.: The log-behavior of the Catalan-Larcombe-French sequences. *Int. J. Number Theory*, **10**, 177–182 (2014)