

Optimal error estimates of compact finite difference discretizations for the Schrödinger-Poisson system

Yong Zhang*

Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, P.R. China.

Abstract. We study compact finite difference methods for the Schrödinger-Poisson equation in a bounded domain and establish their optimal error estimates under proper regularity assumptions on wave function ψ and external potential $V(x)$. The Crank-Nicolson compact finite difference method and the semi-implicit compact finite difference method are both of order $O(h^4 + \tau^2)$ in discrete l^2, H^1 and l^∞ norms with mesh size h and time step τ . For the errors of compact finite difference approximation to the second derivative and Poisson potential are nonlocal, thus besides the standard energy method and mathematical induction method, the key technique in analysis is to estimate the nonlocal approximation errors in discrete l^∞ and H^1 norm by discrete maximum principle of elliptic equation and properties of some related matrix. Also some useful inequalities are established in this paper. Finally, extensive numerical results are reported to support our error estimates of the numerical methods.

AMS subject classifications: 35Q55, 65M06, 65M12, 65M22, 81-08

Key words: Schrödinger-Poisson system, Crank-Nicolson scheme, semi-implicit scheme, compact finite difference method, Gronwall inequality, the maximum principle

1 Introduction

The Schrödinger-Poisson system (SPS) is a local single particle approximation of the time-dependent Hartree-Fock system. It reads, in dimensionless form,

$$i\partial_t\psi(\mathbf{x},t) = \left[-\frac{1}{2}\Delta + V(\mathbf{x}) + \beta\Phi(\mathbf{x},t) \right] \psi(\mathbf{x},t), \quad \mathbf{x} \in \mathbb{R}^d, \quad t > 0, \quad (1.1)$$

$$\nabla^2\Phi(\mathbf{x},t) = -|\psi(\mathbf{x},t)|^2, \quad \mathbf{x} \in \mathbb{R}^d, \quad (1.2)$$

$$\psi(\mathbf{x},t=0) = \psi_0(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^d. \quad (1.3)$$

The complex-valued function $\psi(\mathbf{x},t)$ stands for the single particle wave function with $\lim_{|\mathbf{x}|\rightarrow\infty} |\psi(\mathbf{x},t)| = 0$, $V(\mathbf{x})$ is a given external potential, $\Phi(\mathbf{x},t)$ denotes the Poisson potential subject to free boundary condition, and β is coupling constant. The attractive case ($\beta < 0$) is usually called the Schrödinger-Newton (SN) system and it describes the particle moving in its own gravitational potential, while the repulsive case ($\beta > 0$) describing electrons travelling in its own Coulomb potential is named as Schrödinger-Poisson (SP) system.

The SPS can be rewritten as nonlinear Schrödinger equation (NLS) as

$$i\partial_t\psi(\mathbf{x},t) = \left[-\frac{1}{2}\Delta + V(\mathbf{x}) + \beta\Phi(|\psi|^2,t) \right] \psi(\mathbf{x},t), \quad \mathbf{x} \in \mathbb{R}^d, \quad t > 0. \quad (1.4)$$

*Corresponding author. *Email address:* zhangyong07@mails.tsinghua.edu.cn

Here, the Poisson potential is equivalent to $G_d(|\mathbf{x}|) * |\psi|^2$ with $G_d(|\mathbf{x}|)$ representing the Green function of Poisson equation on \mathbb{R}^d , which is specified as,

$$G_d(|\mathbf{x}|) = \begin{cases} -\frac{1}{2}|\mathbf{x}|, & d=1, \\ -\frac{1}{2\pi}\ln(|\mathbf{x}|), & d=2, \\ \frac{1}{4\pi}|\mathbf{x}|^{-1}, & d=3. \end{cases} \quad (1.5)$$

There are at least two important invariants of (1.4): the *mass of particles*

$$N(\psi) := \|\psi\|^2 = \int_{\mathbb{R}^d} |\psi(\mathbf{x})|^2 d\mathbf{x}, \quad (1.6)$$

and the *total energy*

$$E(\psi) := \int_{\mathbb{R}^d} \frac{1}{2} |\nabla \psi|^2 + V(\mathbf{x})|\psi|^2 + \frac{\beta}{2} \Phi(|\psi|^2)|\psi|^2 d\mathbf{x}. \quad (1.7)$$

The NLS has been studied mathematically and numerically extensively. Mathematically, for the well-posedness, smoothing effects and time behavior of with SPS with/without local term (exchange term), we refer to [4, 7, 14, 20, 21] and references therein. Numerically, different efficient and accurate numerical methods had been proposed to solve NLS, such as the time-splitting spectral/pseudospectral method [3, 8], finite difference method [5, 6, 10, 24] and finite element method [16, 19] and so on. Specially, for the Schrödinger-Poisson equation, we refer the reader to [2, 30] for the time splitting pseudospectral method, to [11, 15, 23] for difference method and etc.

Finite difference method is simplest among them, however, the standard central difference discretization of the Laplacian operator is only of second order accuracy. If combined with the partial differential equation, by carefully designating the finite difference coefficients, one could get higher accuracy with fewer adjacent stencil points such as the compact finite difference method. For details about compact finite difference method, we refer to [25, 26, 28]. Compact finite difference method was popular and had been applied to different models, such as the cubic nonlinear Schrödinger equation, Helmholtz equation and Navier-Stokes equation [18, 22, 27].

Up to our knowledge, compact finite difference method has not yet been applied to SPS. In this paper, we first present Crank-Nicolson compact finite difference scheme (CNCFD) which preserves the conservation laws of energy and mass on the discrete level. While applying CNCFD to SPS, we have to solve a fully nonlinear equation each step which is quite expensive in the view of computation time. Therefore, we propose a semi-implicit compact finite difference scheme (SICFD) in the following and also establish optimal error estimates for both of the schemes.

The paper is organized as follows. In section 2, we present two compact finite difference schemes and their corresponding error estimates in Theorem 2.1 and Theorem 2.2. In section 3, optimal error estimate of CNCFD method for Schrödinger-Poisson is presented by energy method and some related *a priori* bound inequalities are given. The optimal error estimate of SICFD method is established by energy method and mathematical induction in section 4. Extensive numerical results are reported to support our error estimates in the section 5. Finally, some conclusions are made in the last section. Through out the paper, we adopt the standard Sobolev spaces and their corresponding norms and the commonly used constant C does not depend on mesh size h or time step τ if not stated otherwise.

2 Numerical methods and main results

In this section, we introduce CNCFD and SICFD in 1- d for the sake of simplicity. Extensions of CNCFD and SICFD to higher dimensions are possible and similar.

For the wave function ψ decays exponentially fast and constant shift of self-consistent Poisson potential does not affect the physical observation $n = |\psi|^2$, therefore in computational practice we could always truncate the whole space to bounded domain. The 1-d truncated SPS on bounded domain $[a, b]$ reads as

$$i\partial_t\psi(x,t) = \left[-\frac{1}{2}\partial_x^2 + V(x) + \beta\Phi(x,t) \right] \psi(x,t), \quad x \in (a,b), \quad (2.1)$$

$$-\partial_x^2\Phi(x,t) = |\psi(x,t)|^2, \quad x \in (a,b), \quad (2.2)$$

$$\psi(x,0) = \psi_0(x), \quad x \in [a,b], \quad (2.3)$$

subject to boundary condition

$$\psi(a,t) = \psi(b,t) = 0, \quad \Phi(a,t) = \Phi(b,t) = 0. \quad (2.4)$$

The 1-d computational domain $\Omega = [a, b]$ is discretized as $x_j = a + jh, j=0, 1, \dots, M-1, M$ with $h = \frac{b-a}{M}$ and M being a positive integer. Define function space

$$X_M = \{u = (u_j)_{j \in \mathcal{T}_M^0} \mid u_0 = u_M = 0\} \subset \mathbb{C}^{M+1}$$

where $\mathcal{T}_M^0 = \{j \mid j = 0, 1, \dots, M-1, M\}$. Choose time step $\tau := \Delta t$ and denote the time $t_n = n\tau, n=0, 1, \dots$. Let ψ_j^n be the numerical approximation of $\psi(x_j, t_n)$ and $\psi^n \in X_M$ be the numerical solution at time t_n . The standard finite difference operators are listed as follows:

$$\begin{aligned} \delta_x^+ u_j^n &= \frac{u_{j+1}^n - u_j^n}{h}, & \delta_x^- u_j^n &= \frac{u_j^n - u_{j-1}^n}{h}, & \delta_x u_j^n &= \frac{u_{j+1}^n - u_{j-1}^n}{2h}, \\ \delta_t^+ u_j^n &= \frac{u_j^{n+1} - u_j^n}{\tau}, & \delta_t^- u_j^n &= \frac{u_j^n - u_j^{n-1}}{\tau}, & \delta_t u_j^n &= \frac{u_j^{n+1} - u_j^{n-1}}{2\tau}, \\ \mu_t u_j^n &= \frac{u_j^n + u_j^{n+1}}{2}, & \delta_x^2 u_j^n &= \frac{u_{j+1}^n - 2u_j^n + u_{j-1}^n}{h^2}. \end{aligned}$$

2.1 Compact finite difference method

Before presenting the compact finite difference scheme for SPS, we would first make a brief introduction of compact finite difference approximation to the Poisson equation.

For Poisson equation with homogeneous Dirichlet boundary condition

$$\partial_x^2\phi(x) = f(x), \quad x \in (a,b) \quad \text{with} \quad \phi(a) = \phi(b) = 0. \quad (2.5)$$

By Taylor expansion, $\delta_x^2\phi(x_j) = \frac{\phi(x_j+h) - 2\phi(x_j) + \phi(x_j-h)}{h^2}$ approximates $\partial_x^2\phi(x)$ as

$$\begin{aligned} \delta_x^2\phi(x_j) &= (\partial_x^2\phi)(x_j) + \frac{h^2}{12}(\partial_x^4\phi)(x_j) + O(h^4) \\ &= f(x_j) + \frac{h^2}{12}(\partial_x^2(\partial_x^2\phi))(x_j) + O(h^4) \\ &= f(x_j) + \frac{h^2}{12}(\partial_x^2 f)(x_j) + O(h^4) \\ &= f(x_j) + \frac{h^2}{12}(\delta_x^2 f)(x_j) + O(h^4). \end{aligned}$$

Then the Poisson equation could be approximated by $\delta_x^2\phi_j = (1 + \frac{h^2}{12}\delta_x^2)f_j$ with fourth order accuracy. Let $\phi_h, f_h \in X_M$ and denote $A_h = (a_{ij})_{(M-1) \times (M-1)}, \Delta_h = B_h^{-1}A_h$ as the standard central finite difference and fourth order approximation of ∂_x^2 respectively, where

$B_h = \mathbb{I} + \frac{h^2}{12} A_h$, $a_{ii} = -\frac{2}{h^2}$, $a_{i,i+1} = a_{i,i-1} = \frac{1}{h^2}$. Let $I_h: \mathbb{C}^{M+1} \rightarrow \mathbb{C}^{M-1}$ be the standard identity projection operator, i.e.,

$$(I_h u) = (u_1, \dots, u_{M-1})^T \in \mathbb{C}^{M-1}, \quad \forall u = (u_0, \dots, u_M)^T \in \mathbb{C}^{M+1}.$$

Thus the numerical Poisson potential ϕ_h and the second order derivative $f_h = (\partial_x^2 \phi)_h$ are approximated as

$$I_h f_h = \Delta_h (I_h \phi_h), \quad I_h \phi_h = \Delta_h^{-1} (I_h f_h). \quad (2.6)$$

2.2 Numerical methods

Based on the fourth order compact finite difference discretization, the conservative Crank-Nicolson compact finite difference scheme (CNCFD) for SPS reads as follows:

$$i \delta_t^+ \psi_j^n = -\frac{1}{2} \left(\Delta_h I_h \mu_t \psi^n \right)_j + V_j \mu_t \psi_j^n + \beta \mu_t \Phi_j^n \mu_t \psi_j^n, \quad j \in \mathcal{T}_M, \quad n \geq 0, \quad (2.7)$$

where

$$-\Delta_h I_h \Phi^n = I_h (|\psi^n|^2), \quad -\Delta_h I_h \Phi^{n+1} = I_h (|\psi^{n+1}|^2), \quad (2.8)$$

and the index set \mathcal{T}_M is defined as $\mathcal{T}_M = \{j \mid j = 1, 2, \dots, M-1\}$.

The boundary conditions (2.4) are discretized as

$$\psi_0^n = \psi_M^n = 0, \quad \Phi_0^n = \Phi_M^n = 0, \quad n = 0, 1, \dots, \quad (2.9)$$

and the initial value is discretized as

$$\psi_j^0 = \psi_0(x_j), \quad j \in \mathcal{T}_M^0. \quad (2.10)$$

We apply iteration method to solve the nonlinear equation. Given $\psi^n \in X_M$, to solve ψ^{n+1} in (2.7), one could solve its linearized equation, i.e.,

$$i \frac{\psi_j^{*,s+1} - \psi_j^n}{\tau} = -\frac{1}{4} \left[\Delta_h I_h (\psi^{*,s+1} + \psi^n) \right]_j + \left[V_j + \beta \frac{\Phi_j^{*,s} + \Phi_j^n}{2} \right] \frac{\psi_j^{*,s+1} + \psi_j^n}{2}, \quad (2.11)$$

until the $\psi^{*,s}$ converges up to given accuracy that is sufficiently small so as to preserve the conservation of mass and energy on discrete level.

As stated in section 3, the above CNCFD scheme conserves the mass and energy. However, due to the nonlinearity of Poisson potential, one has to solve a fully nonlinear equation which is quite expensive in computation time. Therefore we come up with semi-implicit compact finite difference method (SICFD).

The semi-implicit compact finite difference method, a three-level scheme, reads:

$$i \delta_t \psi_j^n = -\frac{1}{2} \left[\Delta_h I_h \left(\frac{\psi^{n+1} + \psi^{n-1}}{2} \right) \right]_j + V_j \frac{\psi_j^{n+1} + \psi_j^{n-1}}{2} + \beta \Phi_j^n \psi_j^n, \quad n \geq 1, \quad (2.12)$$

and the Poisson potential, boundary condition and the initial value are determined the same way as in CNCFD. The first step value ψ_j^1 could be computed by an at least second order in time accurate scheme, for example, a second order modified Euler method, i.e.,

$$\psi_j^1 = \psi_j^0 - i\tau \left[-\frac{1}{2} \left(\Delta_h I_h \psi^{(1)} \right)_j + V_j \psi_j^{(1)} + \beta \Phi_j^{(1)} \psi_j^{(1)} \right], \quad j \in \mathcal{T}_M, \quad (2.13)$$

$$\psi_j^{(1)} = \psi_j^0 - i\frac{\tau}{2} \left[-\frac{1}{2} \left(\Delta_h I_h \psi^0 \right)_j + V_j \psi_j^0 + \beta \Phi_j^0 \psi_j^0 \right], \quad j \in \mathcal{T}_M, \quad (2.14)$$

where Poisson potential $\Phi^{(1)}$ and Φ^0 are evaluated by scheme (2.8).

Thanks to the equally spaced stencils, equation (2.12) and (2.12) can be accelerated by discrete sine transform (DST), we refer to [17] on DST, which would help reduce the computational cost from $O(M^3)$ (direct linear system solver) to $O(M \log(M))$. Extensions of DST acceleration method to higher dimensions are similar and straightforward.

Remark 2.1. In higher dimension, if the wave function ψ and Poisson potential φ are both specified with homogeneous Dirichlet boundary condition on bounded computation domain, higher order compact finite difference discretizations of $\Delta\psi$ and φ are still applicable, we refer the reader to [25, 26, 28] for more details. Numerical methods proposed here, i.e., CNCFD and SICFD methods, can be generalized to higher dimensions similarly while the error analysis therein is different. However, in practice, we have better accurate and efficient solver in higher dimensions, we refer to [29, 30].

2.3 Main error estimate results

Before presenting the main error estimates, we would like first to introduce some notations and definitions. For any $u, v \in X_M, w \in C^{M-1}$, we define inner product and norms as

$$\begin{aligned} \|u\|_{l^2}^2 &= h \sum_{k=1}^{M-1} |u_k|^2, \quad \|\delta_x^+ u\|_{l^2}^2 = h \sum_{k=0}^{M-1} |\delta_x^+ u_k|^2, \quad \|u\|_{l^\infty} = \max_{j \in \mathcal{T}_M^0} |u_j|, \\ (\delta_x^+ u, \delta_x^+ v) &= h \sum_{k=0}^{M-1} (\delta_x^+ u_k)(\delta_x^+ \bar{v}_k), \quad \langle u, w \rangle = \overline{\langle w, u \rangle} = h \sum_{k=1}^{M-1} u_k \bar{w}_k \\ E_h(u) &= \frac{1}{2} \langle -\Delta_h I_h u, u \rangle + \frac{\beta}{2} \langle \Delta_h^{-1} I_h |u|^2, |u|^2 \rangle + \langle V u, u \rangle, \\ \mathcal{E}(u) &= \frac{1}{2} \langle -\Delta_h I_h u, u \rangle + \langle V u, u \rangle, \end{aligned}$$

where \bar{u}_k denotes the conjugate of u_k .

We make the following assumptions:

$$(A): \quad \psi \in C^0([0, T]; W^{7, \infty}(\Omega) \cap H_0^2(\Omega)) \cap C^1([0, T]; W^{4, \infty}(\Omega)) \cap C^2([0, T]; W^{3, \infty}(\Omega)) \cap C^3([0, T]; L^\infty(\Omega)) \cap C^4([0, T]; L^\infty(\Omega)),$$

where $0 \leq T \leq T_{\max}$ with T_{\max} being the maximal existing time [20, 21].

$$(B): \quad \text{The external potential } V(x) \text{ is smooth.}$$

Denote $M_1 = \max_{0 \leq t \leq T} \|\psi\|_{L^\infty}$ and define the error function $e^n \in X_M$ as

$$e_j^n = \psi(x_j, t_n) - \psi_j^n, \quad j \in \mathcal{T}_M^0, \quad n = 0, 1, \dots.$$

Theorem 2.1. For the CNCFD method, under assumptions (A) and (B), there exists h_0 and τ_0 such that for any $0 < h < h_0$ and $0 < \tau < \tau_0$, the error function satisfies that

$$\|e^n\|_{l^2} + \|\delta_x^+ e^n\|_{l^2} \leq C(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau} \quad (2.15)$$

where the constant C depends on Ω but not on h or τ .

Theorem 2.2. For the SICFD method, under assumptions (A) and (B), there exists h_0 and τ_0 such that for any $0 < h < h_0, 0 < \tau < \tau_0$ and $\tau \leq h$, the error function satisfies that

$$\|e^n\|_{l^2} + \|\delta_x^+ e^n\|_{l^2} \leq C(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau} \quad (2.16)$$

where the constant C depends on Ω but not on h or τ .

3 Error estimates for the CNCFD method

In this section, we will give detailed proof of the main results by energy method with inequalities presented in the following lemma.

Let $u, v \in X_M$, then u, v satisfy the following inequalities:

Lemma 3.1.

$$\langle u, \delta_x^2 v \rangle = -(\delta_x^+ u, \delta_x^+ v) = \langle \delta_x^2 u, v \rangle, \quad \langle A_h I_h u, v \rangle = \langle u, \delta_x^2 v \rangle, \quad (3.1)$$

$$\|u\|_{l^\infty} \leq \sqrt{b-a} \|\delta_x^+ u\|_{l^2}, \quad \|u^2\|_{l^\infty} \leq \sqrt{2} \|u\|_{l^2} \|\delta_x^+ u\|_{l^2}, \quad (3.2)$$

$$\langle -\Delta_h^{-1} I_h |u^2|, |u^2| \rangle \leq \varepsilon \|\delta_x^+ u\|_{l^2}^2 + C_\varepsilon \|u\|_{l^2}^6. \quad (3.3)$$

Proof. The equality (3.1) can be verified using summation by parts as

$$\begin{aligned} \langle u, \delta_x^2 v \rangle &= h \sum_{k=1}^{M-1} u_k \frac{\bar{v}_{k+1} - 2\bar{v}_k + \bar{v}_{k-1}}{h^2} \\ &= -h \sum_{k=0}^{M-1} \left(\frac{u_{k+1} - u_k}{h} \right) \left(\frac{\bar{v}_{k+1} - \bar{v}_k}{h} \right) \\ &= -(\delta_x^+ u, \delta_x^+ v) = \langle \delta_x^2 u, v \rangle. \end{aligned}$$

$\langle A_h I_h u, v \rangle = \langle \delta_x^2 u, v \rangle$ can be proved by definition.

Proof of (3.2). For any $u \in X_M$,

$$\begin{aligned} |u_j| &= \left| \sum_{k=1}^j (u_k - u_{k-1}) \right| = \left| \sum_{k=1}^j \delta_x^+ u_{k-1} \sqrt{h} \sqrt{h} \right| \\ &\leq \left(h \sum_{k=1}^j |\delta_x^+ u_{k-1}|^2 \right)^{\frac{1}{2}} (hj)^{\frac{1}{2}} \leq \sqrt{b-a} \|\delta_x^+ u\|_{l^2}, \end{aligned}$$

take maximum over index $j \in \mathcal{T}_M^0$, we could get

$$\|u\|_{l^\infty} \leq \sqrt{b-a} \|\delta_x^+ u\|_{l^2}.$$

Similarly,

$$u_j^2 = \sum_{k=1}^j u_k^2 - u_{k-1}^2 = \sum_{k=1}^j \left(\frac{u_k - u_{k-1}}{h} \sqrt{h} \right) \left((u_k + u_{k-1}) \sqrt{h} \right),$$

take absolute value on both sides, we have

$$\begin{aligned} |u_j^2| &\leq \left(h \sum_{k=1}^j |\delta_x^+ u_{k-1}|^2 \right)^{1/2} \left(h \sum_{k=1}^j |u_k + u_{k-1}|^2 \right)^{1/2} \\ &\leq \sqrt{2} \|u\|_{l^2} \|\delta_x^+ u\|_{l^2}. \end{aligned}$$

Then by taking maximum over $j \in \mathcal{T}_M^0$, we could get $\|u^2\|_{l^\infty} \leq \sqrt{2} \|u\|_{l^2} \|\delta_x^+ u\|_{l^2}$.

The inequality $\langle -\Delta_h^{-1} I_h |u^2|, |u^2| \rangle \leq \varepsilon \|\delta_x^+ u\|_{l^2}^2 + C_\varepsilon \|u\|_{l^2}^6$ is useful in estimating the Poisson potential energy. Herein the Δ_h^{-1} approximates inverse Poisson operator.

$$\begin{aligned} |\langle -\Delta_h^{-1} I_h |u^2|, |u^2| \rangle| &= \left| \sum_{k=1}^{M-1} (-\Delta_h^{-1} I_h |u^2|)_k |u_k|^2 h \right| \\ &\leq \|u\|_{l^2}^2 \|\Delta_h^{-1} I_h |u^2|\|_{l^\infty} \leq C \|u\|_{l^2}^2 \|u\|_{l^\infty}^2 \\ &\leq C \|u\|_{l^2}^3 \|\delta_x^+ u\|_{l^2} \leq \varepsilon \|\delta_x^+ u\|_{l^2}^2 + C_\varepsilon \|u\|_{l^2}^6, \end{aligned}$$

where we use discrete maximum principle in $\|\Delta_h^{-1} I_h |u^2|\|_{l^\infty}$ estimate and in last equality we apply Young inequality with ε being any positive real number. \square

In order to analyze the H^1 error, we need to investigate some related matrix in detail. The following lemma establish some useful properties of approximation matrix $A_h, B_h, \Delta_h \in \mathbb{R}^{M-1 \times M-1}$.

Lemma 3.2. (Properties of related approximation matrix) For any $u, v \in X_M$ and matrix A_h, B_h, Δ_h , we have

$$\|A_h^{-1} I_h u\|_{l^\infty} \leq C \|u\|_{l^\infty}, \quad \|A_h^{-1} I_h u\|_{l^2} \leq \frac{(b-a)^2}{\pi^2} \|u\|_{l^2}, \quad (3.4)$$

$$\|B_h I_h u\|_{l^\infty} \leq \|u\|_{l^\infty}, \quad \|B_h^{-1} I_h u\|_{l^\infty} \leq \frac{3}{2} \|u\|_{l^\infty}, \quad (3.5)$$

$$\|B_h I_h u\|_{l^2} \leq \|u\|_{l^2}, \quad \|B_h^{-1} I_h u\|_{l^2} \leq \frac{3}{2} \|u\|_{l^2}, \quad (3.6)$$

$$\langle -A_h I_h u, u \rangle \leq \langle -\Delta_h I_h u, u \rangle \leq \frac{3}{2} \langle -A_h I_h u, u \rangle, \quad (3.7)$$

where the constant C depends on Ω but not on h or u .

Proof. By applying discrete maximum principle in [13], one can get $\|A_h^{-1} I_h u\|_{l^\infty} \leq C \|u\|_{l^\infty}$ where the constant C depends on Ω but not on h or u . Standard contradiction argument would lead us to $\|B_h^{-1} I_h u\|_{l^\infty} \leq \frac{3}{2} \|u\|_{l^\infty}$. If examining entries of B_h carefully, one can get

$$\|B_h\|_{l^\infty} = \max_{1 \leq i \leq M-1} \sum_{j=1}^{M-1} |(B_h)_{ij}| \leq 1, \text{ thus we finished the proof of } \|B_h I_h u\|_{l^\infty} \leq \|u\|_{l^\infty}.$$

Note that $-A_h, B_h$ are commutable positive definite matrix, then they have the same eigenvectors. The j -th eigenvalue of $-A_h$ equals to $\lambda_j(-A_h) = \frac{2}{h^2} (1 - \cos(\frac{\pi j}{M}))$, $j = 1, 2, \dots, M-1$ and the j -th eigenvalue of B_h is $\lambda_j(B_h) = 1 - \frac{1}{6} (1 - \cos(\frac{\pi j}{M})) \in (\frac{2}{3}, 1)$, then we have $\lambda_j(-A_h^{-1}) = \frac{h^2}{2} \frac{1}{1 - \cos(\frac{\pi j}{M})}$ which implies that $\|A_h^{-1}\|_{l^2} = \max_{1 \leq j \leq M-1} |\lambda_j(-A_h^{-1})| \leq$

$$\frac{(b-a)^2}{\pi^2}. \text{ Thus we prove } \|A_h^{-1} I_h u\|_{l^2} \leq \frac{(b-a)^2}{\pi^2} \|u\|_{l^2}.$$

Similarly, we have

$$\|B_h\|_{l^2} = \max_{1 \leq j \leq M-1} |\lambda_j(B_h)| \leq 1, \quad \|B_h^{-1}\|_{l^2} = \max_{1 \leq j \leq M-1} |\lambda_j(B_h^{-1})| \leq \frac{3}{2}.$$

Therefore we get $\|B_h I_h u\|_{l^2} \leq \|u\|_{l^2}$, $\|B_h^{-1} I_h u\|_{l^2} \leq \frac{3}{2} \|u\|_{l^2}$.

The last inequality states the equivalence of two kinetic energies. Notice $\langle -A_h I_h u, u \rangle = \|\sqrt{-A_h} I_h u\|_{l^2}^2$, $\langle -\Delta_h I_h u, u \rangle = \|\sqrt{-\Delta_h} I_h u\|_{l^2}^2$, and denote $S = \sqrt{-A_h}$, $T = \sqrt{-\Delta_h}$ with $ST = TS$. Then we have

$$\begin{aligned} \|\sqrt{-A_h} I_h u\|_{l^2} &= \|S I_h u\|_{l^2} = \|T(T^{-1}S) I_h u\|_{l^2} \\ &\leq \|T\|_{l^2} \|(T^{-1}S) I_h u\|_{l^2} \leq \|T\|_{l^2} \|\sqrt{-\Delta_h} I_h u\|_{l^2}, \\ \|\sqrt{-\Delta_h} I_h u\|_{l^2} &= \|(T^{-1}S) I_h u\|_{l^2} \leq \|T^{-1}\|_{l^2} \|S I_h u\|_{l^2} \leq \|T^{-1}\|_{l^2} \|\sqrt{-A_h} I_h u\|_{l^2}, \end{aligned}$$

which implies

$$\begin{aligned}\langle -A_h I_h u, u \rangle &= \|\sqrt{-A_h} I_h u\|_{l^2}^2 \leq \|T\|_{l^2}^2 \langle -\Delta_h I_h u, u \rangle, \\ \langle -\Delta_h I_h u, u \rangle &= \|\sqrt{-\Delta_h} I_h u\|_{l^2}^2 \leq \|T^{-1}\|_{l^2}^2 \langle -A_h I_h u, u \rangle.\end{aligned}$$

For $\|B_h\|_{l^2}$ is bounded, we can conclude that $\|T^{-1}\|_{l^2}^2 = \|B_h^{-1}\|_{l^2} \leq \frac{3}{2}$, $\|T\|_{l^2}^2 \leq 1$. Thus we prove the equivalence inequality. \square

Since the fourth order compact finite difference scheme has been proposed before, the following lemma completes detailed error analysis.

Lemma 3.3. (Error estimates of compact finite difference scheme) For Poisson equation (2.5), let $f_{\text{ext}} = (f(x_0), f(x_1), \dots, f(x_M))^T$, $\phi_{\text{ext}} = (\phi(x_0), \phi(x_1), \dots, \phi(x_M))^T$ be the exact solution and $f_h = (f_0, f_1, \dots, f_M)^T$, $\phi_h = (\phi_0, \phi_1, \dots, \phi_M)^T$ be the numerical approximation obtained by the fourth order compact finite difference scheme.

Assume $\phi(x) \in C^6([a, b])$, $f(x) \in C^4([a, b])$, if $f(x)$ is known, we get $\|\phi_h - \phi_{\text{ext}}\|_{l^\infty} = O(h^4, \partial_x^4 f)$. Inversely, with $\phi(x)$ being known, $\|f_h - f_{\text{ext}}\|_{l^\infty} = O(h^4, \partial_x^6 \phi)$.

Proof. This proof is mainly based on Taylor formula with integral remainder and discrete maximum principle of elliptic equation.

Firstly, if $\phi(x) \in C^6([a, b])$, $f(x) \in C^4([a, b])$, by Taylor formula, we have

$$\begin{aligned}\delta_x^2 \phi(x_j) &= (\partial_x^2 \phi)(x_j) + \frac{h^2}{12} (\partial_x^4 \phi)(x_j) + \mathcal{R}_j^I \\ &= f(x_j) + \frac{h^2}{12} (\partial_x^2 f)(x_j) + \mathcal{R}_j^I \\ &= f(x_j) + \frac{h^2}{12} \delta_x^2 f_j + \mathcal{R}_j^I + \mathcal{R}_j^{\text{II}},\end{aligned}$$

where

$$\begin{aligned}\mathcal{R}_j^I &= \frac{1}{5!h^2} \int_0^h (h-s)^5 [\phi^{(6)}(x_j+s) + \phi^{(6)}(x_j-s)] ds, \\ \mathcal{R}_j^{\text{II}} &= -\frac{1}{72} \int_0^h (h-s)^3 [f^{(4)}(x_j+s) + f^{(4)}(x_j-s)] ds.\end{aligned}$$

To solve the Poisson potential in (2.5), we apply the compact finite difference method as $-\Delta_h I_h \phi_h = I_h f_{\text{ext}}$. Then $A_h I_h (\phi_{\text{ext}} - \phi_h) = \mathcal{R}^I + \mathcal{R}^{\text{II}}$, by discrete maximum principle, it can be concluded that

$$\|\phi_h - \phi_{\text{ext}}\|_{l^\infty} \leq C \|\mathcal{R}^I + \mathcal{R}^{\text{II}}\|_{l^\infty} \leq Ch^4 \|\partial_x^4 f\|_{L^\infty},$$

where $\mathcal{R}^I, \mathcal{R}^{\text{II}} \in C^{M-1}$ and the constant C depends on Ω but not on $f(x)$ or h .

Reversely, the second derivative of $\phi(x)$ was approximated by $I_h f_h = \Delta_h I_h \phi_{\text{ext}}$ and the equation $B_h I_h (f_h - f_{\text{ext}}) = (\mathcal{R}^I + \mathcal{R}^{\text{II}})$ holds. Then we have

$$\|f_h - f_{\text{ext}}\|_{l^\infty} \leq \frac{3}{2} \|\mathcal{R}^I + \mathcal{R}^{\text{II}}\|_{l^\infty} \leq Ch^4 \|\partial_x^6 \phi\|_{L^\infty}.$$

\square

Remark 3.1. One must have noticed that the approximation error in f_{ext} , ϕ_{ext} are both globally dependent on $\phi(x)$ or $f(x)$, and this is quite different from the cubic nonlinear or other local nonlinear Schrödinger equations.

Lemma 3.4. (Conservation of mass and energy) For the CNCFD method (2.7) with (2.9) and (2.10), for any time step $\tau > 0$ and mesh size $h > 0$ and initial data ψ_0 . It conserves the mass and energy in the discretized level, i.e.,

$$\|\psi^n\|_{l^2} = \|\psi^0\|_{l^2}, \quad E_h(\psi^n) = E_h(\psi^0). \quad (3.8)$$

Proof. One can apply a similar process as in [5], so we omit it for brevity. \square

Lemma 3.5. (Solvability of the difference equation) For any given ψ^n , under assumptions-(A) and (B), there exists a solution $\psi^{n+1} \in X_M$ satisfies (2.7). There exists $\tau_0 > 0$, such that, the solution is unique for $0 < \tau < \tau_0$.

Proof. Proof for existence and uniqueness of CNCFD are similar to those in [1]. Rewrite (2.7) as

$$\psi^{n+1/2} = \psi^n + i \frac{\tau}{2} F^n(\psi^{n+1/2}), \quad n=0,1,\dots,$$

where $F^n : X_M \rightarrow X_M$ is defined as

$$(F^n(u))_j = -\frac{1}{2}(A_h I_h u)_j + V_j u_j + \beta \left[-A_h^{-1} I_h \left(\frac{|\psi^n|^2 + |2u - \psi^n|^2}{2} \right) \right]_j u_j.$$

Define the map $G^n : X_M \rightarrow X_M$ as

$$G^n(u) = u - \psi^n - i \frac{\tau}{2} F^n(u), \quad u \in X_M,$$

and it is continuous. Moreover,

$$\operatorname{Re} \langle G^n(u), u \rangle = \|u\|_{l^2}^2 - \operatorname{Re} \langle \psi^n, u \rangle \geq \|u\|_{l^2} (\|u\|_{l^2} - \|\psi^n\|_{l^2}), \quad u \in X_M,$$

which implies

$$\lim_{\|u\|_{l^2} \rightarrow \infty} \frac{|\langle G^n(u), u \rangle|}{\|u\|_{l^2}} = \infty.$$

Thus G^n is surjective according to theorem in [12], that is to say, there exists a solution $u_0 \in X_M$ satisfy $G^n(u_0) = 0$.

We can use standard energy argument to prove uniqueness of (2.7). Assume $u, v \in X_M$ satisfies equation (2.7) for given ψ^n . Denote $w = u - v \in X_M$, we have

$$i \frac{w_j}{\tau} = -\frac{1}{2} \frac{(\Delta_h I_h w)_j}{2} + \frac{V_j w_j}{2} + \beta \chi_j, \quad j \in \mathcal{T}_M,$$

where

$$\chi_j := \frac{1}{4} \left[-\Delta_h^{-1} I_h (|u|^2 + |\psi^n|^2) \right]_j (u_j + \psi_j^n) - \frac{1}{4} \left[-\Delta_h^{-1} I_h (|v|^2 + |\psi^n|^2) \right]_j (v_j + \psi_j^n).$$

Multiply both sides (3.9) by $\overline{w_j} h$ and then take imaginary part of the summation over $j \in \mathcal{T}_M$, we can get

$$\begin{aligned} \frac{\|w\|_{l^2}^2}{\tau} &= \left| \frac{\beta}{4} h \sum_{j=1}^{M-1} \left(\Delta_h^{-1} I_h (|u|^2 + |\psi^n|^2) \right)_j |w_j|^2 + \left(\Delta_h^{-1} I_h (|u|^2 - |v|^2) \right)_j (v_j + \psi_j^n) \overline{w_j} \right| \\ &\leq C \|w\|_{l^2}^2 (\|u\|_{l^\infty}^2 + \|\psi^n\|_{l^\infty}^2) + C (\|v\|_{l^\infty} + \|\psi^n\|_{l^\infty}) \|w\|_{l^2} \|\Delta_h^{-1} I_h (|u|^2 - |v|^2)\|_{l^2} \\ &\leq C (\|u\|_{l^\infty}^2 + \|v\|_{l^\infty}^2 + \|\psi^n\|_{l^\infty}^2) (\|w\|_{l^2}^2 + \| |u|^2 - |v|^2 \|_{l^2}^2). \end{aligned}$$

As stated before, the CNCFD scheme preserves the mass and energy in the discretized level, by applying inequalities in (3.2)-(3.3), we have

$$\begin{aligned} E_h(u) &= \langle -\Delta_h I_h u, u \rangle + \frac{\beta}{2} \langle \Delta_h^{-1} I_h |u|^2, |u|^2 \rangle + h \sum_{k=1}^{M-1} V_k u_k \bar{u}_k \\ &\geq \|\delta_x^+ u\|_{l_2}^2 - |\beta| \varepsilon \|\delta_x^+ u\|_{l_2}^2 - C_\varepsilon \|u\|_{l_2}^6, \end{aligned}$$

so $\|\delta_x^+ u\|_{l_2}^2 \leq E_h(\psi^0) + C_\varepsilon \|\psi^0\|_{l_2}^6$. By simple calculations, we have

$$\| |u|^2 - |v|^2 \|_{l_2}^2 = h \sum_{j=1}^{M-1} [u_j(\bar{u}_j - \bar{v}_j) + (u_j - v_j)\bar{v}_j]^2 \leq C (\|u\|_{l_\infty}^2 + \|v\|_{l_\infty}^2) \|w\|_{l_2}^2.$$

Put together all the inequalities listed above, we have

$$\begin{aligned} \|w\|_{l_2}^2 &\leq C \tau \|w\|_{l_2}^2 (\|u\|_{l_\infty}^2 + \|v\|_{l_\infty}^2 + \|\psi^n\|_{l_\infty}^2 + 1)^2, \\ &\leq C \tau \|w\|_{l_2}^2 (\|\delta_x^+ u\|_{l_2}^2 + \|\delta_x^+ v\|_{l_2}^2 + \|\delta_x^+ \psi^n\|_{l_2}^2 + C_0)^2, \\ &\leq C \tau \|w\|_{l_2}^2 (E_h(\psi^0) + C_\varepsilon \|\psi^0\|_{l_2}^6 + C_0)^2, \end{aligned}$$

where C_0 is a constant that does not depend on h or ψ . There exists τ_0 such that $|C\tau(E_h(\psi^0) + C_\varepsilon \|\psi^0\|_{l_2}^6 + C_0)| \leq \frac{1}{2}$ for any $0 \leq \tau \leq \tau_0$, which implies that

$$\|w\|_{l_2}^2 = \|u - v\|_{l_2}^2 = 0 \quad \implies \quad u = v,$$

thus we finish the proof of the uniqueness of solution to (2.7). \square

Denote the local truncation error of CNCFD scheme by η^n , which is defined as

$$\begin{aligned} \eta_j^n &:= i \delta_t^+ \psi(x_j, t_n) + \frac{1}{2} [\Delta_h I_h \mu_t \psi(\cdot, t_n)]_j - V(x_j) \mu_t \psi(x_j, t_n) \\ &\quad - \beta \left[-\Delta_h^{-1} I_h \mu_t |\psi(\cdot, t_n)|^2 \right]_j \mu_t \psi(x_j, t_n), \quad j \in \mathcal{T}_M, \quad n \geq 0, \end{aligned}$$

where $\psi(\cdot, t_n) = (\psi(x_0, t_n), \psi(x_1, t_n), \dots, \psi(x_M, t_n))^T \in X_M$.

Lemma 3.6. (Local truncation error) Under assumptions (A) and (B), the local truncation error for CNCFD is

$$\|\eta^n\|_{l_\infty} = O(h^4 + \tau^2), \quad \|\eta^n\|_{l_2} = O(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau},$$

and

$$\|\delta_x^+ \eta^n\|_{l_2} = O(h^4 + \tau^2), \quad \|\delta_x^+ \eta^n\|_{l_\infty} = O(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau}.$$

Proof. Firstly, by Taylor formula with integral remainder, we have

$$\begin{aligned} \delta_t^+ \psi(x_j, t_n) &= (\partial_t \psi)(x_j, t_{n+\frac{1}{2}}) + \mathcal{Q}_j^{n+\frac{1}{2}}(\psi), \\ V(x_j) \mu_t \psi(x_j, t_n) &= V(x_j) \psi(x_j, t_{n+\frac{1}{2}}) + V(x_j) \mathcal{P}_j^{n+\frac{1}{2}}(\psi), \\ \Delta_h I_h \mu_t \psi(x_j, t_n) &= (\partial_x^2 \psi)(x_j, t_{n+\frac{1}{2}}) + \mathcal{P}_j^{n+\frac{1}{2}}(\partial_x^2 \psi) + \left[B_h^{-1} \mu_t \mathcal{R}^n(\psi) \right]_j, \end{aligned}$$

where $\mathcal{Q}_j^{n+\frac{1}{2}}, \mathcal{P}_j^{n+\frac{1}{2}}, \mathcal{R}_j^n$ are defined as

$$\begin{aligned} \mathcal{Q}_j^{n+\frac{1}{2}}(f(x, t)) &= \frac{1}{4\tau} \int_0^\tau (\tau-s)^2 \left[\partial_t^3 f(x_j, t_{n+\frac{1}{2}}+s) + \partial_t^3 f(x_j, t_{n+\frac{1}{2}}-s) \right] ds, \\ \mathcal{P}_j^{n+\frac{1}{2}}(f(x, t)) &= \int_0^\tau (\tau-s) \left[\partial_t^2 f(x_j, t_{n+\frac{1}{2}}+s) + \partial_t^2 f(x_j, t_{n+\frac{1}{2}}-s) \right] ds, \\ \mathcal{R}_j^n(f(x, t)) &= \frac{1}{5!h^2} \int_0^h (h-s)^5 \left[f^{(6)}(x_j+s, t_n) + f^{(6)}(x_j-s, t_n) \right] ds \\ &\quad - \frac{1}{72} \int_0^h (h-s)^3 \left[f^{(6)}(x_j+s, t_n) + f^{(6)}(x_j-s, t_n) \right] ds. \end{aligned}$$

Let $\Phi(\cdot, t_n) = (\Phi(x_0, t_n), \Phi(x_1, t_n), \dots, \Phi(x_M, t_n))^T \in X_M$, then

$$\left[-\Delta_h^{-1} I_h \mu_t |\psi(\cdot, t_n)|^2 \right]_j = \Phi(x_j, t_{n+\frac{1}{2}}) + \mathcal{P}_j^{n+\frac{1}{2}}(\Phi) - \left[A_h^{-1} \mu_t \mathcal{R}^n(\Phi) \right]_j,$$

thus

$$\begin{aligned} & \left[-\Delta_h^{-1} I_h \mu_t |\psi(\cdot, t_n)|^2 \right]_j \mu_t \psi(x_j, t_n) \\ &= \left[\Phi(x_j, t_{n+\frac{1}{2}}) + \mathcal{P}_j^{n+\frac{1}{2}}(\Phi) - \left(A_h^{-1} \mu_t \mathcal{R}^n(\Phi) \right)_j \right] \left[\psi(x_j, t_{n+\frac{1}{2}}) + \mathcal{P}_j^{n+\frac{1}{2}}(\psi) \right]. \end{aligned}$$

By discrete maximum principle of elliptic equation, we have

$$\|\Phi\|_{L^\infty} \leq C \|\psi\|^2_{L^\infty}, \quad \|\partial_t^2 \Phi\|_{L^\infty} \leq C \|\partial_t^2 \psi\|^2_{L^\infty},$$

therefore,

$$\begin{aligned} & \left| \left[-\Delta_h^{-1} I_h \mu_t |\psi(\cdot, t_n)|^2 \right]_j \mu_t \psi(x_j, t_n) - (\Phi \psi)(x_j, t_{n+\frac{1}{2}}) \right| \\ &= \left| \Phi(x_j, t_{n+\frac{1}{2}}) \mathcal{P}_j^{n+\frac{1}{2}}(\psi) + \left[\psi(x_j, t_{n+\frac{1}{2}}) + \mathcal{P}_j^{n+\frac{1}{2}}(\psi) \right] \left[\mathcal{P}_j^{n+\frac{1}{2}}(\Phi) - \left(A_h^{-1} \mu_t \mathcal{R}^n(\Phi) \right)_j \right] \right| \\ &\leq h^4 (\|\partial_x^6 \Phi\|_{L^\infty} \|\psi\|_{L^\infty}) + \tau^2 (\|\psi\|_{L^\infty}^2 \|\partial_t^2 \psi\|_{L^\infty} + \|\psi\|_{L^\infty} \|\partial_t^2 \Phi\|_{L^\infty}) + O(h^4 \tau^2 + \tau^4) \\ &\leq h^4 (\|\partial_x^4 |\psi|^2\|_{L^\infty} \|\psi\|_{L^\infty}) + \tau^2 (\|\psi\|_{L^\infty}^2 \|\partial_t^2 \psi\|_{L^\infty} + \|\psi\|_{L^\infty} \|\partial_t^2 \psi\|^2_{L^\infty}) + O(h^4 \tau^2 + \tau^4). \end{aligned}$$

Finally, local truncation error η_j^n can be written in integral form as

$$\begin{aligned} \eta_j^n &= i \mathcal{Q}_j^{n+\frac{1}{2}}(\psi) + \frac{1}{2} \mathcal{P}_j^{n+\frac{1}{2}}(\partial_x^2 \psi) + \frac{1}{2} \left[B_h^{-1} \mu_t \mathcal{R}^n(\psi) \right]_j - \left[V(x_j) + \beta \Phi(x_j, t_{n+\frac{1}{2}}) \right] \mathcal{P}_j^{n+\frac{1}{2}}(\psi) \\ &\quad - \beta \left[\psi(x_j, t_{n+\frac{1}{2}}) + \mathcal{P}_j^{n+\frac{1}{2}}(\psi) \right] \left[\mathcal{P}_j^{n+\frac{1}{2}}(\Phi) - \left(A_h^{-1} \mu_t \mathcal{R}^n(\Phi) \right)_j \right]. \end{aligned}$$

Thus we can get

$$\begin{aligned} |\eta_j^n| &\leq C \tau^2 (\|\partial_t^2 \psi\|_{L^\infty} + \|\partial_t^3 \psi\|_{L^\infty} + \|\partial_t^2 \partial_x^2 \psi\|_{L^\infty} + \|\psi\|_{L^\infty}^2 \|\partial_t^2 \psi\|_{L^\infty} + \|\psi\|_{L^\infty} \|\partial_t^2 \psi\|^2_{L^\infty}) \\ &\quad + C h^4 (\|\partial_x^6 \psi\|_{L^\infty} + \|\partial_x^4 |\psi|^2\|_{L^\infty} \|\psi\|_{L^\infty}) + O(h^4 \tau^2 + \tau^4), \end{aligned}$$

and this would lead us to $\|\eta^n\|_{l^\infty} = O(h^4 + \tau^2)$ and $\|\eta^n\|_{l^2} = O(h^4 + \tau^2)$.

To evaluate $\delta_x^+ \eta_j^n$, we just need to estimate the nonlocal term error, i.e., $\left(B_h^{-1} \mu_t \mathcal{R}^n(\psi) \right)_j$ and $\left(A_h^{-1} \mu_t \mathcal{R}^n(\Phi) \right)_j$, because the local term could be dealt with by standard technique [1]. We just take $\delta_x^+ \left(B_h^{-1} \mu_t \mathcal{R}^n(\psi) \right)_j$ as an example and the other term is estimated the same way.

Notice $B_h^{-1} \mu_t \mathcal{R}^n(\psi) \in X_M$ and denote

$$\begin{aligned} \mathcal{R}^n(\psi) &= (\mathcal{R}_1^n(\psi), \dots, \mathcal{R}_{M-1}^n(\psi))^T, \quad \mathcal{R}_x^n(\psi) = (\delta_x^+ \mathcal{R}_1^n(\psi), \dots, \delta_x^+ \mathcal{R}_{M-2}^n(\psi))^T, \\ g_j &= \left[B_h^{-1} \mu_t \mathcal{R}^n(\psi) \right]_j, \quad \mathcal{G} = (g_1, \dots, g_{M-1})^T, \quad \mathcal{G}_x = (\delta_x^+ g_1, \dots, \delta_x^+ g_{M-2})^T. \end{aligned}$$

Reformulate the equation, we get

$$\mathcal{G}_x = B_h^{-1} \mu_t \mathcal{R}_x^n + B_h^{-1} \mathbf{b}, \quad (3.9)$$

with $\mathbf{b} = \frac{1}{12}(\delta_x^+ g_0, 0, \dots, 0, \delta_x^+ g_{M-1})_{M-2}^T$.

Then by taking l^2 norm on both sides, we have

$$\sum_{j=1}^{M-2} |\delta_x^+ g_j|^2 h \leq C \sum_{j=1}^{M-2} (|\delta_x^+ \mathcal{R}_j^n|^2 + |\delta_x^+ \mathcal{R}_j^{n+1}|^2) h + (|\delta_x^+ g_0|^2 + |\delta_x^+ g_{M-1}|^2) h,$$

while $\delta_x^+ g_0$ can be estimated as

$$\begin{aligned} |\delta_x^+ g_0| &= \left| \frac{g_1 - g_0}{h} \right| = \left| \frac{g_1}{h} \right| = \left| \frac{[B_h^{-1} \mu_t \mathcal{R}^n(\psi)]_1}{h} \right| = \left| \frac{\sum_{j=1}^{M-1} (B_h^{-1})_{1,j} \mu_t \mathcal{R}_j^n(\psi)}{h} \right| \\ &\leq C \sum_{j=1}^{M-1} d_j \frac{\mu_t \mathcal{R}_j^n(\psi)}{h} \leq C \sum_{j=1}^{M-1} \frac{12}{(4+2\sqrt{6})^j} \frac{\mu_t \mathcal{R}_j^n(\psi)}{h} \leq Ch^4 \|\partial_x^7 \psi\|_{L^\infty}, \end{aligned}$$

where $|d_j| = [B_h^{-1}]_{1,j} \leq \frac{12}{(4+2\sqrt{6})^j}$, $j = 1, 2, \dots, M-1$ and the last inequality holds because $\partial_x^6 \psi(a) = \partial_x^6 \psi(b) = 0$ under regularity assumptions (A) and (B).

Similarly, $|\delta_x^+ g_{M-1}| \leq Ch^4 \|\partial_x^7 \psi\|_{L^\infty}$. Therefore $\|\delta_x^+ g\|_{l^2} \leq Ch^4 \|\partial_x^7 \psi\|_{L^\infty}$. Taking maximum norm of (3.9) on both sides and using inequality (3.5), we can have $\|\delta_x^+ g\|_{l^\infty} \leq Ch^4 \|\partial_x^7 \psi\|_{L^\infty}$.

Thus, we obtain

$$\|\delta_x^+ \eta^n\|_{l^2} = O(h^4 + \tau^2), \quad \|\delta_x^+ \eta^n\|_{l^\infty} = O(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau}.$$

□

Theorem 3.1. (*l² norm estimate*) Under assumptions (A) and (B), there exist h_0 and τ_0 such that for any $0 < h < h_0$ and $0 < \tau < \tau_0$, the error function satisfies that

$$\|e^n\|_{l^2} \leq C(\tau^2 + h^4), \quad 0 \leq n \leq \frac{T}{\tau}, \quad (3.10)$$

where the constant C does not depend on h or τ .

Proof. Choose a smooth function $\alpha \in C^\infty([0, \infty))$, which is defined as

$$\alpha(\rho) = \begin{cases} 1 & , \quad 0 \leq \rho \leq 1, \\ \in [0, 1] & , \quad 1 \leq \rho \leq 2, \\ 0 & , \quad \rho \geq 2. \end{cases}$$

Define truncating function $F_{M_0}(\rho)$ as

$$F_{M_0}(\rho) = \alpha\left(\frac{\rho}{M_0}\right) \rho, \quad 0 \leq \rho \leq \infty, \quad M_0 = 2(1 + M_1^2) \geq 0,$$

so that $F_{M_0}(\rho)$ satisfy the following Lipschitz condition, that is,

$$|F_{M_0}(\rho_1) - F_{M_0}(\rho_2)| \leq C|\sqrt{\rho_1} - \sqrt{\rho_2}|, \quad 0 \leq \rho_1, \rho_2 \leq \infty.$$

Introduce an auxiliary scheme of $\phi \in X_M$, taking $\phi^0 = \psi^0$ as the initial value, which is given by

$$i\delta_t^+ \phi_j^n = -\frac{1}{2} [\Delta_h I_h \mu_t \phi^n]_j + V_j \mu_t \phi_j^n + \beta \mu_t \left[-\Delta_h^{-1} I_h F_{M_0}(|\phi^n|^2) \right]_j \mu_t \phi_j^n. \quad (3.11)$$

Herein, ϕ_j^n can be viewed as another approximation of $\psi(x_j, t_n)$. Define the 'auxiliary error' function $\tilde{e}^n \in X_M$

$$\tilde{e}_j^n = \psi(x_j, t_n) - \phi_j^n, \quad j \in \mathcal{T}_M^0, \quad n \geq 0.$$

The corresponding local truncation error $\tilde{\eta}^n$ is defined as

$$\begin{aligned} \tilde{\eta}_j^n &= i\delta_t^+ \psi(x_j, t_n) + \frac{1}{2} [\Delta_h I_h \mu_t \psi(\cdot, t_n)]_j - V(x_j) \mu_t \psi(x_j, t_n) \\ &\quad - \beta \mu_t \left[-\Delta_h^{-1} I_h F_{M_0} (|\psi(\cdot, t_n)|^2) \right]_j \mu_t \psi(x_j, t_n). \end{aligned} \quad (3.12)$$

Similarly, the local truncation error $\|\tilde{\eta}^n\|_{l^\infty} = O(h^4 + \tau^2)$, $0 \leq n < \frac{T}{\tau}$. Subtracting (3.12) from (3.11), we have

$$i\delta_t^+ \tilde{e}_j^n = -\frac{1}{2} [\Delta_h I_h \mu_t \tilde{e}^n]_j + V_j \mu_t \tilde{e}_j^n + \tilde{\xi}_j^n + \tilde{\eta}_j^n, \quad (3.13)$$

with

$$\tilde{\xi}_j^n = \beta \mu_t \left[-\Delta_h^{-1} I_h F_{M_0} (|\psi(\cdot, t_n)|^2) \right]_j \mu_t \psi(x_j, t_n) - \beta \mu_t \left[-\Delta_h^{-1} I_h F_{M_0} (|\phi^n|^2) \right]_j \mu_t \phi_j^n.$$

Let $\Phi_j^n = \left[-\Delta_h^{-1} I_h F_{M_0} (|\phi^n|^2) \right]_j$ and $\tilde{\Phi}_j^n = \left[-\Delta_h^{-1} I_h F_{M_0} (|\psi(\cdot, t_n)|^2) \right]_j$, we have

$$\begin{aligned} \left| \text{Im} \langle \tilde{\xi}_j^n, \tilde{e}_j^n + \tilde{e}_j^{n+1} \rangle \right| &= |\beta| \left| \text{Im} \langle \mu_t (\tilde{\Phi}_j^n - \Phi_j^n) \mu_t \psi(x_j, t_n) - \mu_t \Phi_j^n \mu_t \tilde{e}_j^n, 2\mu_t \tilde{e}_j^n \rangle \right| \\ &= |\beta| \left| \text{Im} \langle \mu_t (\tilde{\Phi}_j^n - \Phi_j^n) \mu_t \psi(x_j, t_n), 2\mu_t \tilde{e}_j^n \rangle \right|. \end{aligned}$$

Denote $v, w \in \mathbb{C}^{M-1}$ as

$$v_j = F_{M_0} (|\psi(x_j, t_n)|^2) - F_{M_0} (|\phi_j^n|^2), \quad w_j = \tilde{e}_j^n + \tilde{e}_j^{n+1}, \quad j = 1, 2, \dots, M-1.$$

Then by the Lipschitz condition of F_{M_0} , we have

$$|v_j| \leq C \left| |\psi(x_j, t_n)| - |\phi_j^n| \right| \leq C |\psi(x_j, t_n) - \phi_j^n| = C |\tilde{e}_j^n|, \quad j = 1, 2, \dots, M-1.$$

Therefore, by Cauchy-Schwartz inequality, we obtain

$$\begin{aligned} \left| \text{Im} \langle \mu_t (\tilde{\Phi}_j^n - \Phi_j^n) \mu_t \psi(x_j, t_n), 2\mu_t \tilde{e}_j^n \rangle \right| &\leq C \rho(A_h^{-1}) \sum_{j=1}^{M-1} (|v_j|^2 + |w_j|^2) h \\ &\leq C (\|\tilde{e}^n\|_{l_2}^2 + \|\tilde{e}^{n+1}\|_{l_2}^2), \end{aligned}$$

and

$$\begin{aligned} \left| \langle \tilde{\eta}_j^n, \mu_t \tilde{e}_j^n \rangle \right| &\leq C \|\tilde{\eta}^n\|_{l_2} \|\mu_t \tilde{e}^n\|_{l_2} \leq C (\|\tilde{\eta}^n\|_{l_2}^2 + \|\mu_t \tilde{e}^n\|_{l_2}^2) \\ &\leq C (\|\tilde{\eta}^n\|_{l^\infty}^2 + \|\tilde{e}^n\|_{l_2}^2 + \|\tilde{e}^{n+1}\|_{l_2}^2). \end{aligned}$$

Multiply both sides of equation (3.13) by $2h\mu_t \tilde{e}_j^n$ and sum up over index $j \in \mathcal{T}_M$, after taking imaginary parts, we have

$$\|\tilde{e}^{n+1}\|_{l_2}^2 - \|\tilde{e}^n\|_{l_2}^2 \leq C\tau (\|\tilde{\eta}^n\|_{l^\infty}^2 + \|\tilde{e}^{n+1}\|_{l_2}^2 + \|\tilde{e}^n\|_{l_2}^2), \quad 0 \leq n < \frac{T}{\tau}$$

Applying discrete Gronwall inequality ([27]), we have

$$\|\tilde{e}^n\|_{l_2} \leq C(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau},$$

and

$$\|\tilde{e}^n\|_{l^\infty} \leq \|\tilde{e}^n\|_{l^2} h^{-1/2} \leq C(h^{\frac{7}{2}} + \frac{\tau^2}{\sqrt{h}}) \leq Ch, \quad \tau \leq h,$$

thus

$$|\phi_j^n| \leq |\psi(x_j, t_n)| + |\tilde{e}_j^n| \leq \frac{\sqrt{M_0}}{2} + Ch \leq \sqrt{M_0},$$

which infers $F_{M_0}(|\phi_j^n|^2) = |\phi_j^n|^2$ and

$$\phi_j^n = \psi_j^n, \quad e_j^n = \tilde{e}_j^n, \quad j=1,2,\dots,M-1, \quad 0 \leq n \leq \frac{T}{\tau}.$$

□

Next, we prove Theorem 2.1.

Proof. The local truncation equation (3.13) is reduced as

$$i\delta_t^+ e_j^n = -\frac{1}{2}[\Delta_h I_h \mu_t e^n]_j + V_j \mu_t e_j^n + \xi_j^n + \eta_j^n, \quad (3.14)$$

where

$$\xi_j^n = \beta \left[-\Delta_h^{-1} I_h \mu_t |\psi(\cdot, t_n)|^2 \right]_j \mu_t \psi(x_j, t_n) - \beta \left[-\Delta_h^{-1} I_h \mu_t |\psi^n|^2 \right]_j \mu_t \phi_j^n, \quad (3.15)$$

satisfies

$$\|\delta_x^+ \xi^n\|_{l^2} \leq C((h^4 + \tau^2)^2 + \|e^n\|_{l^2} + \|e^{n+1}\|_{l^2} + \|\delta_x^+ e^n\|_{l^2} + \|\delta_x^+ e^{n+1}\|_{l^2}). \quad (3.16)$$

Rewrite (3.14) as

$$e_j^{n+1} - e_j^n = -i\tau(\mathcal{L}_j^n + \xi_j^n + \eta_j^n),$$

where \mathcal{L}_j^n is defined as

$$\mathcal{L}_j^n = -\frac{1}{2}[\Delta_h I_h \mu_t e^n]_j + V_j \mu_t e_j^n, \quad j=1,2,\dots,M-1, \quad n \geq 0.$$

Multiply both sides of equation (3.14) by $2h \overline{e_j^{n+1} - e_j^n}$ and sum up over index $j \in \mathcal{T}_M$, after taking real parts, we have

$$\begin{aligned} \mathcal{E}(e^{n+1}) - \mathcal{E}(e^n) &= -2 \operatorname{Re} \langle \xi^n + \eta^n, e^{n+1} - e^n \rangle \\ &= -2 \operatorname{Re} \langle \xi^n + \eta^n, -i\tau(\mathcal{L}^n + \xi^n + \eta^n) \rangle \\ &= 2\tau \operatorname{Im} \langle \xi^n + \eta^n, \mathcal{L}^n \rangle. \end{aligned}$$

By Cauchy-Schwartz inequality and equivalent energies inequality, we have

$$\begin{aligned} |\langle \eta^n, \mathcal{L}^n \rangle| &\leq |\langle \eta^n, -\frac{1}{2} \Delta_h I_h \mu_t e^n \rangle| + |\langle \eta^n, V \mu_t e^n \rangle| \\ &\leq C(|\langle (-\Delta_h)^{\frac{1}{2}} I_h \eta^n, (-\Delta_h)^{\frac{1}{2}} I_h \mu_t e^n \rangle| + \|e^n\|_{l^2}^2 + \|e^{n+1}\|_{l^2}^2 + (h^4 + \tau^2)^2) \\ &\leq C(\|(-A_h)^{\frac{1}{2}} I_h \eta^n\|_{l^2}^2 + \|(-A_h)^{\frac{1}{2}} I_h \mu_t e^n\|_{l^2}^2 + (h^4 + \tau^2)^2) \\ &\leq C(\|\delta_x^+ \eta^n\|_{l^2}^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|e^{n+1}\|_{l^2}^2 + (h^4 + \tau^2)^2) \\ &\leq C((h^4 + \tau^2)^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|\delta_x^+ e^{n+1}\|_{l^2}^2). \end{aligned}$$

Similarly, we have

$$\begin{aligned} |\langle \bar{\zeta}^n, \mathcal{L}^n \rangle| &\leq |\langle \bar{\zeta}^n, -\frac{1}{2} \Delta_h I_h \mu_t e^n \rangle| + |\langle \bar{\zeta}^n, V \mu_t e^n \rangle| \\ &\leq C(\|\delta_x^+ \bar{\zeta}^n\|_{l^2}^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|\delta_x^+ e^{n+1}\|_{l^2}^2 + \|e^n\|_{l^2}^2 + \|e^{n+1}\|_{l^2}^2) \\ &\leq C((h^4 + \tau^2)^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|\delta_x^+ e^{n+1}\|_{l^2}^2 + \|e^n\|_{l^2}^2 + \|e^{n+1}\|_{l^2}^2), \end{aligned}$$

where the last inequality holds because of (3.16).

Thus, we have

$$|\operatorname{Im} \langle \bar{\zeta}^n + \eta^n, \mathcal{L}^n \rangle| \leq C(h^4 + \tau^2)^2 + \mathcal{E}(e^{n+1}) + \mathcal{E}(e^n),$$

Combing the above inequalities, we get

$$\mathcal{E}(e^{n+1}) - \mathcal{E}(e^n) \leq C\tau \left[(h^4 + \tau^2)^2 + \mathcal{E}(e^{n+1}) + \mathcal{E}(e^n) \right].$$

Then there exists $\tau_0 > 0$, when $0 < \tau \leq \tau_0$, by Gronwall inequality,

$$\mathcal{E}(e^n) \leq C(h^4 + \tau^2)^2, \quad 0 \leq n \leq \frac{T}{\tau},$$

which, together with theorem (3.1), finish the proof of Theorem 2.1. \square

Remark 3.2. If we consider the general SPS with local term $|\psi|^{\frac{2}{d}}$, Theorem 2.1 still holds true. We refer to [1] on how to deal with the local term.

4 Error estimates for the SICFD method

In this section, we present optimal error estimate for the SICFD method (2.12)-(2.14) with initial value (2.10) and Dirichlet boundary condition (2.9) in discrete l^2 and H^1 norm.

First, we prove the solvability and uniqueness of the SICFD method solution.

Lemma 4.1. (*Solvability and uniqueness of the SICFD method*) Under assumptions (A) and (B), for any initial value $\psi^0 \in X_M$, there exists a unique solution $\psi^n \in X_M$ of (2.13)-(2.14) for $n=1$ and (2.12) for $n > 1$.

Proof. The lemma holds true for $n=1$. First, we prove the uniqueness of (2.12). Given $\psi^n, \psi^{n-1} \in X_M$, suppose there exist two solutions $u, v \in X_M$, i.e.,

$$i \frac{u_j - \psi_j^{n-1}}{2\tau} = -\frac{1}{2} \left[\Delta_h I_h \left(\frac{u + \psi^{n-1}}{2} \right) \right]_j + V_j \frac{u_j + \psi_j^{n-1}}{2} + \beta \Phi_j^n \psi_j^n, \quad (4.1)$$

$$i \frac{v_j - \psi_j^{n-1}}{2\tau} = -\frac{1}{2} \left[\Delta_h I_h \left(\frac{v + \psi^{n-1}}{2} \right) \right]_j + V_j \frac{v_j + \psi_j^{n-1}}{2} + \beta \Phi_j^n \psi_j^n. \quad (4.2)$$

Set $w = u - v$ and subtract (4.2) from (4.1), we have

$$i \frac{w_j}{\tau} = -\frac{1}{2} [\Delta_h I_h w]_j + V_j w_j. \quad (4.3)$$

Multiply both sides of (4.3) by $\bar{w}_j h$ and take imaginary parts of summation over $j \in \mathcal{T}_M$, we get $\|w\|_{l^2} = 0$ which implies $u = v$, thus the uniqueness of the solution is proved.

For the solvability, we follow similar process as CNCFD method. Equation (2.12) can be rewritten as

$$i \psi_j^{n+1} + \frac{\tau}{2} \left[\Delta_h I_h \psi^{n+1} \right]_j - \tau V_j \psi_j^{n+1} + \chi_j = 0,$$

where

$$\chi_j = -i\psi_j^{n-1} + \frac{\tau}{2} [\Delta_h I_h \psi^{n-1}]_j - \tau V_j \psi_j^{n-1} - 2\beta\tau \Phi_j^n \psi_j^n.$$

Define map $G: u \in X_M \rightarrow G(u) \in X_M$ as

$$G(u) = iu_j + \frac{\tau}{2} [\Delta_h I_h u]_j - \tau V_j u_j + \chi_j.$$

The map G is continuous from X_M to X_M and satisfies

$$|\operatorname{Im} \langle G(u), u \rangle| = \|u\|_{l^2}^2 + \operatorname{Im} \langle \chi, u \rangle \geq \|u\|_{l^2}^2 - \|\chi\|_{l^2} \|u\|_{l^2},$$

from which infers

$$\lim_{\|u\|_{l^2} \rightarrow \infty} \frac{|\langle G^n(u), u \rangle|}{\|u\|_{l^2}} = \infty.$$

Thus G^n is surjective according to theorem in [12], that is to say, there exists a solution $u_0 \in X_M$ satisfy $G^n(u_0) = 0$. \square

Define local truncation error function η^n for SICFD method as

$$\begin{aligned} \eta_j^n &= i\delta_t \psi(x_j, t_n) + \frac{1}{2} \left[\Delta_h I_h \left(\frac{\psi(\cdot, t_{n+1}) + \psi(\cdot, t_{n-1})}{2} \right) \right]_j \\ &\quad - V(x_j) \frac{\psi(x_j, t_{n+1}) + \psi(x_j, t_{n-1})}{2} - \beta \left[-\Delta_h^{-1} I_h |\psi(\cdot, t_n)|^2 \right]_j \psi(x_j, t_n), \quad n > 1 \end{aligned} \quad (4.4)$$

and for $n=0$

$$\eta_j^0 = i\delta_t^+ \psi(x_j, 0) - \left[-\frac{1}{2} \left(\Delta_h I_h \psi^{(1)} \right)_j + V(x_j) \psi_j^{(1)} + \beta \Phi_j^{(1)} \psi_j^{(1)} \right], \quad j \in \mathcal{T}_M, \quad (4.5)$$

with

$$\psi_j^{(1)} = \psi_0(x_j) - i\frac{\tau}{2} \left[-\frac{1}{2} \left(\Delta_h I_h \psi(\cdot, 0) \right)_j + V(x_j) \psi_0(x_j) + \beta \Phi_j^0 \psi_0(x_j) \right], \quad j \in \mathcal{T}_M.$$

Then we have

Lemma 4.2. (Local truncation error) Under assumptions (A) and (B), we have local truncation error η^n as

$$\|\eta^n\|_{l^\infty} \leq C(h^4 + \tau^2), \quad 0 \leq n \leq \frac{T}{\tau} - 1, \quad \|\delta_x^+ \eta^0\|_{l^\infty} \leq C(h^3 + \tau), \quad \text{for } \tau \leq Ch \quad (4.6)$$

$$\|\delta_x^+ \eta^n\|_{l^\infty} \leq C(h^4 + \tau^2), \quad 1 \leq n \leq \frac{T}{\tau} - 1. \quad (4.7)$$

Proof. When $n=0$, we have

$$\psi_j^{(1)} = \psi(x_j, \tau/2) + i\frac{\tau}{2} \eta_j^{(1)}, \quad j \in \mathcal{T}_M,$$

where truncation error $\eta_j^{(1)}, j \in \mathcal{T}_M$ satisfies

$$\begin{aligned} \eta_j^{(1)} &= i \frac{\psi(x_j, \tau/2) - \psi_0(x_j)}{\tau/2} + \frac{1}{2} [\Delta_h \psi_0]_j - V_j \psi_0(x_j) - \beta \Phi_j^0 \psi_0(x_j) \\ &= \frac{i}{\tau/2} \int_0^{\tau} (\tau-s) \partial_t^2 \psi(x_j, s) ds + \left[B_h^{-1} I_h \mathcal{R}^0(\psi) \right]_j - \beta \left[A_h^{-1} I_h \mathcal{R}^0(\Phi) \right]_j \psi_0(x_j). \end{aligned}$$

We can conclude that $\|\eta^{(1)}\|_{l^\infty} = O(h^4 + \tau)$. Then

$$\begin{aligned}\eta_j^0 &= i \frac{\psi(x_j, \tau) - \psi_0(x_j)}{\tau} + \frac{1}{2} \left[\Delta_h I_h \psi^{(1)} \right]_j - V(x_j) \psi_j^{(1)} - \beta \Phi_j^{(1)} \psi_j^{(1)} \\ &= \mathcal{Q}_j^{\frac{1}{2}}(\psi) + \frac{i\tau}{4} \left[\Delta_h I_h \eta^{(1)} \right]_j + \frac{1}{2} \left[B_h^{-1} I_h \mathcal{R}^{\frac{1}{2}}(\psi) \right]_j - \frac{i\tau}{2} V(x_j) \eta_j^{(1)} \\ &\quad - \frac{i\tau}{2} \beta \Phi_j^{(1)} \eta_j^{(1)} + \beta \psi(x_j, \tau/2) (\Phi(x_j, \tau/2) - \Phi_j^{(1)}).\end{aligned}$$

By similar argument in CNCFD, we can obtain $\|\eta^0\|_{l^\infty} \leq C(h^4 + \tau^2)$ and

$$|\delta_x^+ \eta_j^0| \leq C \frac{\|\eta_j^0\|_{l^\infty}}{h} \leq C(h^3 + \tau), \quad j=0, 1, \dots, M-,$$

for $\tau \leq Ch$. Thus, we prove local truncation error for $n=0$. While for $n > 1$, by Taylor expansion with integral remainder, we get

$$\begin{aligned}\eta_j^n &= \mathcal{Q}_j^{n+\frac{1}{2}}(\psi) + \frac{1}{2} \left[B_h^{-1} I_h \frac{\mathcal{R}^{n+1}(\psi) + \mathcal{R}^{n-1}(\psi)}{2} \right]_j + \frac{1}{2} \mathcal{P}_j^{n+\frac{1}{2}}(\partial_x^2 \psi) \\ &\quad - V(x_j) \mathcal{P}_j^{n+\frac{1}{2}}(\psi) - \beta \left[A_h^{-1} I_h \mathcal{R}^n(\Phi) \right]_j \psi(x_j, t_n).\end{aligned}$$

Thus

$$|\eta_j^n| \leq Ch^4 (\|\partial_x^6 \psi\|_{L^\infty} + \|\partial_x^4 \psi\|_{L^\infty} \|\psi\|_{L^\infty}) + C\tau^2 (\|\partial_t^2 \psi\|_{L^\infty} + \|\partial_t^3 \psi\|_{L^\infty} + \|\partial_t^2 \partial_x^2 \psi\|_{L^\infty}),$$

which implies $\|\eta^n\|_{l^\infty} \leq C(h^4 + \tau^2)$, $1 \leq n \leq \frac{T}{\tau} - 1$. Similarly, we get

$$\begin{aligned}\delta_x^+ \eta_j^n &= \delta_x^+ \mathcal{Q}_j^{n+\frac{1}{2}}(\psi) + \frac{1}{2} \delta_x^+ \left[B_h^{-1} I_h \frac{\mathcal{R}^{n+1}(\psi) + \mathcal{R}^{n-1}(\psi)}{2} \right]_j + \frac{1}{2} \delta_x^+ \mathcal{P}_j^{n+\frac{1}{2}}(\partial_x^2 \psi) \\ &\quad + \delta_x^+ (V(x_j) \mathcal{P}_j^{n+\frac{1}{2}}(\psi)) - \beta \delta_x^+ \left(\left[A_h^{-1} I_h \mathcal{R}^n(\Phi) \right]_j \psi(x_j, t_n) \right),\end{aligned}$$

from which we obtain

$$\begin{aligned}|\delta_x^+ \eta_j^n| &\leq Ch^4 (\|\partial_x^7 \psi\|_{L^\infty} + \|\partial_x^5 \psi\|_{L^\infty} \|\psi\|_{L^\infty} + \|\partial_x^4 \psi\|_{L^\infty} \|\partial_x \psi\|_{L^\infty} + \|\partial_x^3 \psi\|_{L^\infty} \|\partial_x^2 \psi\|_{L^\infty}) \\ &\quad + C\tau^2 (\|\partial_x \partial_t^2 \psi\|_{L^\infty} + \|\partial_x \partial_t^3 \psi\|_{L^\infty} + \|\partial_t^2 \partial_x^3 \psi\|_{L^\infty}), \quad 1 \leq n \leq \frac{T}{\tau} - 1,\end{aligned}$$

where the nonlocal term $A_h^{-1} I_h \mathcal{R}^n(\Phi)$ and $B_h^{-1} I_h (\mathcal{R}^{n+1}(\psi) + \mathcal{R}^{n-1}(\psi))$ are dealt with similarly. Therefore, we prove the local error estimate. \square

Theorem 4.1. (*l^2 norm estimate*) Under assumptions (A) and (B), there exist $h_0 > 0$ and $\tau > 0$, when $0 < h < h_0$ and $0 < \tau < \tau_0, \tau \leq h$, we have

$$\|e^n\|_{l^2} \leq C(h^4 + \tau^2), \quad \|\psi^n\|_{l^\infty} \leq 1 + M_1, \quad 1 \leq n \leq \frac{T}{\tau}. \quad (4.8)$$

Proof. If $\|e^n\|_{l^2} \leq C(h^4 + \tau^2)$ is known, from $e_j^n = \psi(x_j, t_n) - \psi_j^n$, we can get

$$\begin{aligned}\|\psi^n\|_{l^\infty} &\leq \|\psi(\cdot, t_n)\|_{L^\infty} + \|e^n\|_{l^\infty} \leq M_1 + h^{-\frac{1}{2}} \|e^n\|_{l^2} \\ &\leq M_1 + C(h^{\frac{7}{2}} + h^{\frac{3}{2}}) \leq M_1 + 1,\end{aligned}$$

with sufficiently small h and τ . Therefore we only need to verify $\|e^n\|_{l^2} \leq C(h^4 + \tau^2)$.

We will use mathematical induction to prove the l^2 norm estimate. The initial inequality $\|e^1\|_{l^2} \leq C(h^4 + \tau^2)$ holds true spontaneously. Now we assume that (4.8) is

valid for $0 \leq n \leq m-1 \leq \frac{T}{\tau} - 1$, then we need to confirm that it is still valid when $n = m$. Subtracting (4.4) from (2.12), we have

$$i\delta_i^+ e_j^n = \mathcal{L}_j^n + \zeta_j^n + \eta_j^n, \quad (4.9)$$

where

$$\begin{aligned} \mathcal{L}_j^n &= -\frac{1}{2} \left[\Delta_h I_h \frac{e^{n+1} + e^{n-1}}{2} \right]_j + V_j \frac{e_j^{n+1} + e_j^{n-1}}{2}, \quad j=1,2,\dots,M-1, \quad n \geq 0. \\ \zeta_j^n &= \beta \left[-\Delta_h^{-1} I_h |\psi(\cdot, t_n)|^2 \right]_j \psi(x_j, t_n) - \beta \left[-\Delta_h^{-1} I_h |\psi^n|^2 \right]_j \psi_j^n, \quad j=1,2,\dots,M-1, \quad n \geq 0. \end{aligned}$$

Similarly, noticing (3.4) and (3.16), we have

$$\begin{aligned} \|\zeta^n\|_{l^2}^2 &\leq C \|e^n\|_{l^2}^2, \quad 1 \leq n \leq m-1, \\ \|\delta_x^+ \zeta^n\|_{l^2}^2 &\leq C \left((h^4 + \tau^2)^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|e^n\|_{l^2}^2 \right), \quad 1 \leq n \leq m-1. \end{aligned} \quad (4.10)$$

Take imaginary parts of the summation on each sides after multiplying (4.9) by $\overline{e_j^{n+1} + e_j^{n-1}}$, by Cauchy-Schwartz inequality, we have

$$\begin{aligned} \|e^{n+1}\|_{l^2}^2 - \|e^{n-1}\|_{l^2}^2 &= 2\tau \operatorname{Im} \langle \zeta^n + \eta^n, e^{n+1} + e^{n-1} \rangle \\ &\leq 2\tau \left[\|e^{n+1}\|_{l^2}^2 + \|e^{n-1}\|_{l^2}^2 + \|\zeta\|_{l^2}^2 + \|\eta^n\|_{l^2}^2 \right] \\ &\leq C\tau(h^4 + \tau^2)^2 + 2\tau (\|e^{n+1}\|_{l^2}^2 + \|e^{n-1}\|_{l^2}^2) + C\tau \|e^n\|_{l^2}^2. \end{aligned}$$

Noticing $e^0 = 0$ and $\|e^1\|_{l^2}^2 \leq C(h^4 + \tau^2)^2$, when $\tau \leq \frac{1}{4}$, sum up the above inequality from $n=1$ to $n=m-1$ and rewrite the inequality, we have

$$\frac{1}{2} (\|e^m\|_{l^2}^2 + \|e^{m-1}\|_{l^2}^2) \leq C(m-1)\tau(h^4 + \tau^2)^2 + \|e^1\|_{l^2}^2 + (C\tau + 4\tau) \sum_{k=1}^{m-1} \|e^k\|_{l^2}^2,$$

which implies

$$\|e^m\|_{l^2}^2 \leq 2C(T+1)(h^4 + \tau^2)^2 + 2(C\tau + 4\tau) \sum_{k=1}^{m-1} \|e^k\|_{l^2}^2. \quad (4.11)$$

Apply Gronwall inequality to (4.11), we get

$$\begin{aligned} \|e^m\|_{l^2}^2 &\leq 2C(T+1)(h^4 + \tau^2)^2 e^{2(C\tau + 4\tau)m} \\ &\leq 2C(T+1)e^{2(C+4)T} (h^4 + \tau^2)^2. \end{aligned}$$

Hence, the proof is finished. \square

Then we continue to prove Theorem (2.2).

Proof. As is stated before, $e^0 = 0$ and thus Theorem 2.2 holds true.

$$|\delta_x^+ e_j^1| = |\delta_x^+ (\psi(x_j, \tau) - \psi_j^1)| = |-i\tau \delta_x^+ \eta_j^0| \leq C\tau(h^3 + \tau) \leq C(h^4 + \tau^2),$$

which implies Theorem (2.2) is true for $n=1$. Rewrite equation (4.9) as

$$e_j^{n+1} - e_j^{n-1} = -2i\tau (\mathcal{L}_j^n + \zeta_j^n + \eta_j^n).$$

Multiply both sides of (4.9) by $2h \overline{e_j^{n+1} - e_j^{n-1}}$ and sum up over index $j \in \mathcal{T}_M$, after taking real parts, we have

$$\begin{aligned} \mathcal{E}(e^{n+1}) - \mathcal{E}(e^{n-1}) &= -2\operatorname{Re}\langle \zeta^n + \eta^n, e^{n+1} - e^{n-1} \rangle \\ &= -2\operatorname{Re}\langle \zeta^n + \eta^n, -2i\tau(\mathcal{L}^n + \zeta^n + \eta^n) \rangle \\ &= 4\tau \operatorname{Im}\langle \zeta^n + \eta^n, \mathcal{L}^n \rangle. \end{aligned}$$

By Cauchy-Schwartz inequality and equivalent energy inequalities, we have

$$\begin{aligned} |\langle \zeta^n, \mathcal{L}^n \rangle| &\leq C((h^4 + \tau^2)^2 + \|\delta_x^+ e^{n-1}\|_{l^2}^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|\delta_x^+ e^{n+1}\|_{l^2}^2), \\ |\langle \eta^n, \mathcal{L}^n \rangle| &\leq C((h^4 + \tau^2)^2 + \|\delta_x^+ e^{n-1}\|_{l^2}^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|\delta_x^+ e^{n+1}\|_{l^2}^2) \end{aligned}$$

holds true for any $1 \leq n \leq \frac{T}{\tau} - 1$. Thus, we can get

$$\begin{aligned} \mathcal{E}(e^{n+1}) - \mathcal{E}(e^{n-1}) &\leq C\tau(h^4 + \tau^2)^2 + \tau \left[\|\delta_x^+ e^{n-1}\|_{l^2}^2 + \|\delta_x^+ e^n\|_{l^2}^2 + \|\delta_x^+ e^{n+1}\|_{l^2}^2 \right] \\ &\leq C\tau(h^4 + \tau^2)^2 + \tau \left[\mathcal{E}(e^{n-1}) + \mathcal{E}(e^n) + \mathcal{E}(e^{n+1}) \right], 1 \leq n \leq \frac{T}{\tau} - 1. \end{aligned}$$

Then sum up the above inequality from $n=1$ to $n=m-1 \leq \frac{T}{\tau} - 1$, we have, for $\tau \leq \frac{1}{4}$ and $\tau \leq h$,

$$\frac{1}{2}(\mathcal{E}(e^m) - \mathcal{E}(e^{m-1})) \leq C(T+1)(h^4 + \tau^2)^2 + C\tau \sum_{k=1}^{m-1} \mathcal{E}(e^k), \quad 1 \leq m \leq \frac{T}{\tau}.$$

Apply discrete Gronwall inequality to the above inequality, we get

$$\frac{1}{2} \|\delta_x^+ e^n\|_{l^2}^2 \leq \mathcal{E}(e^n) \leq C(T+1)e^{CT} (h^4 + \tau^2)^2.$$

The proof for theorem (2.2) is complete. \square

Lemma 4.3. (*l^∞ norm estimate*) For the SICFD method, under assumptions (A) and (B), the l^∞ norm is of order $O(h^4 + \tau^2)$.

Proof. Since we have obtained $\|\delta_x^+ e^n\|_{l^2} = O(h^4 + \tau^2)$, then the l^∞ norm is valid immediately by inequality (3.2). \square

Remark 4.1. If we consider the general SPS with local term $|\psi|^{\frac{2}{\alpha}}$, Theorem 2.2 of the l^2 and H^1 error estimates holds true. We refer to [1] on how to deal with the local term.

Remark 4.2. Compared with (2.7), the SICFD scheme does not require to solve any nonlinear equation, thus it takes up less computational cost and more appropriate to be extended to higher dimensions.

5 Numerical results

In this section, we present numerical results to verify the error estimates. We choose symmetric computation domain with/without external potential $V(x) = \frac{x^2}{2}$. Let ψ_e be the numerical 'exact' solution obtained with fine mesh h and small time step $\tau = h^2$ at time $T = 0.5$ and $\psi_{h,\tau}$ be the numerical solution with mesh size h and time step τ . Let $e_{h,\tau} = \psi_e - \psi_{h,\tau}$ denote the error function. The initial value is chosen as $\psi_0(x) = e^{-x^2/2} \pi^{-1/4}$. The numerical 'exact' solution is obtained with $h = \frac{1}{256}, \tau = h^2$ on interval $[-16, 16]$.

Table 1: Temporal error analysis of CNCFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x) = \frac{x^2}{2}$.

		$\tau=2^{-8}$	$\tau=2^{-9}$	$\tau=2^{-10}$	$\tau=2^{-11}$
$\beta=5$	$\ e\ _{l^2}$	3.907E-03	9.774E-04	2.443E-04	6.104E-05
	Rate	1.999	2.000	2.001	
	$\ \delta_x^+ e\ _{l^2}$	3.857E-03	9.650E-04	2.412E-04	6.0268e-05
	Rate	1.999	2.000	2.001	
	$\ e\ _{l^\infty}$	2.650E-03	6.627E-04	1.657E-04	4.138E-05
	Rate	2.000	2.000	2.001	
$\beta=-5$	$\ e\ _{l^2}$	3.564E-03	8.914E-04	2.228E-04	5.567E-05
	Rate	1.999	2.000	2.001	
	$\ \delta_x^+ e\ _{l^2}$	3.618E-03	9.050E-04	2.2623E-04	5.653E-05
	Rate	1.999	2.000	2.001	
	$\ e\ _{l^\infty}$	3.177E-03	7.946E-04	1.987E-04	4.963E-05
	Rate	1.999	2.000	2.001	

First, we test the second order temporal accuracy by choosing different time step τ with uniform fine mesh size $h = 1/256$ on interval $[-16,16]$ and the temporal convergence rate is defined as $\log_2(\|e_{h,\tau}\|/\|e_{h,\frac{\tau}{2}}\|)$ with corresponding discrete norm. Table 1, 2, 3 and 4 show errors $\|e\|_{l^2}$, $\|\delta_x^+ e\|_{l^2}$ and $\|e\|_{l^\infty}$ for CNCFD and SICFD schemes with $\beta = \pm 5$ and external potential $V(x) = x^2/2$ or $V(x) = 0$.

Next, we verify the fourth order spatial accuracy by choosing appropriate mesh size and time step (h,τ) and the convergence rate is defined as $\log_2(\|e_{h,\tau}\|/\|e_{\frac{h}{2},\frac{\tau}{4}}\|)$ with corresponding norm. Table 5 and 6 show errors $\|e\|_{l^2}$, $\|\delta_x^+ e\|_{l^2}$ and $\|e\|_{l^\infty}$ for the CNCFD method with Poisson coupling constants $\beta = \pm 5$ and external potential $V(x) = 0$ or $V(x) = \frac{x^2}{2}$. Table 7 and 8 show errors $\|e\|_{l^2}$, $\|\delta_x^+ e\|_{l^2}$ and $\|e\|_{l^\infty}$ for the SICFD method with coupling Poisson constants $\beta = \pm 5$ and external potential $V(x) = 0$ or $V(x) = \frac{x^2}{2}$.

Table 9 lists computation time cost by CNCFD and SICFD with different mesh sizes and time step for repulsive SPS ($\beta=5$) at time $T=1.0$ with/without external potential. The algorithm was carried out by Linux (version 3.0.0-16) MATLAB (version 7.8.0.347 (R2009a)) on Intel CPU (i3-530). Figure 1 depicts the long time evolution of the errors of discrete mass and energy for SICFD scheme with mesh size $h=1/16$ when external potential $V(x)=0$ and Poisson constant $\beta=5$.

Based on numerical results shown and not shown here, we can draw the following conclusions: (i) Tables 5, 6, 7 and 8 confirm the $O(h^4 + \tau^2)$ convergence estimates for the CNCFD and SICFD scheme in l^2 , H^1 and l^∞ discrete norms. (ii) The CNCFD scheme conserves discrete mass and energy analytically, however, as stated in [1], we have to solve the nonlinear equation accurate enough (up to machine accuracy or nearby) so as to keep the conservation laws numerically. Extensive numerical experiments not shown here confirms that. (iii) From Figure 1, we can observe oscillatory error evolution phenomenon which is similar to that in [1]. (iv) Table 9 shows that the efficiency of SICFD is superior to that of CNCFD with/without external potential. All the linear systems involved in both schemes are accelerated by DST and this make the computation time depends almost linearly on mesh size h . The DST accelerated linear system solver can be extended to higher dimensions directly which would still preserve the linear dependency of computation time and mesh size.

Table 2: Temporal error analysis of CNCFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x)=0$.

		$\tau=2^{-8}$	$\tau=2^{-9}$	$\tau=2^{-10}$	$\tau=2^{-11}$
$\beta=5$	$\ e\ _{l^2}$	3.720E-03	9.305E-04	2.326E-04	5.811E-05
	Rate	1.999	2.000	2.001	
	$\ \delta_x^+ e\ _{l^2}$	4.752E-03	1.189E-03	2.972E-04	7.426E-05
	Rate	1.999	2.000	2.001	
	$\ e\ _{l^\infty}$	2.447E-03	6.119E-04	1.530E-04	3.821E-05
	Rate	2.000	2.000	2.001	
$\beta=-5$	$\ e\ _{l^2}$	3.633E-03	9.088E-04	2.272E-04	5.676E-05
	Rate	1.999	2.000	2.001	
	$\ \delta_x^+ e\ _{l^2}$	3.131E-03	7.833E-04	1.958E-04	4.892E-05
	Rate	1.999	2.000	2.001	
	$\ e\ _{l^\infty}$	3.024E-03	7.565E-04	1.891E-04	4.724E-05
	Rate	1.999	2.000	2.001	

Table 3: Temporal error analysis of SICFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x)=\frac{x^2}{2}$.

		$\tau=2^{-8}$	$\tau=2^{-9}$	$\tau=2^{-10}$	$\tau=2^{-11}$
$\beta=5$	$\ e\ _{l^2}$	6.780E-03	1.698E-03	4.243E-04	1.060E-04
	Rate	2.002	2.001	2.001	
	$\ \delta_x^+ e\ _{l^2}$	6.127E-03	1.529E-03	3.820E-04	9.542E-05
	Rate	2.003	2.001	2.001	
	$\ e\ _{l^\infty}$	4.840E-03	1.209E-03	3.021E-04	7.548E-05
	Rate	2.001	2.000	2.001	
$\beta=-5$	$\ e\ _{l^2}$	8.206E-03	2.045E-03	5.104E-04	1.272E-04
	Rate	2.005	2.002	2.005	
	$\ \delta_x^+ e\ _{l^2}$	8.872E-03	2.210E-03	5.517E-04	1.375E-04
	Rate	2.005	2.002	2.005	
	$\ e\ _{l^\infty}$	7.535E-03	1.878E-03	4.687E-04	1.168E-04
	Rate	2.005	2.002	2.005	

6 Conclusions

We presented error analysis of two compact finite difference methods, i.e., the conservative Crank-Nicolson compact finite difference scheme and the semi-implicit compact finite difference scheme, for the Schrödinger-Poisson equation in a bounded domain under proper regularity assumptions on wave function ψ and external potential $V(x)$. Both of the schemes are of order $O(h^4 + \tau^2)$ in discrete l^2 , H^1 and l^∞ norms with mesh size h and time step τ . We analyzed the local truncation error of the compact finite difference for the second derivative and Poisson potential, which is not nonlocal, by the discrete maximum principle of the elliptic equation and properties of related approximation matrix. We analyzed the nonlocal approximation term in the local truncation error and global error with Taylor formula with integral remainders. In the proof of CNCFD scheme, we used a Lipschitz function to approximate the nonlinearity so as to obtain l^2 norm estimate and by inverse inequality to get a *priori* bound for l^∞ ; for SICFD

Table 4: Temporal error analysis of SICFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x)=0$.

		$\tau=2^{-8}$	$\tau=2^{-9}$	$\tau=2^{-10}$	$\tau=2^{-11}$
$\beta=5$	$\ e\ _{l^2}$	6.522E-03	1.629E-03	4.069E-04	1.014E-04
	Rate	2.001	2.001	2.004	
	$\ \delta_x^+ e\ _{l^2}$	7.098E-03	1.771E-03	4.423E-04	1.102E-04
	Rate	2.003	2.002	2.005	
	$\ e\ _{l^\infty}$	4.455E-03	1.113E-03	2.780E-04	6.931E-05
	Rate	2.001	2.001	2.004	
$\beta=-5$	$\ e\ _{l^2}$	7.930E-03	1.977E-03	4.935E-04	1.230E-04
	Rate	2.004	2.002	2.005	
	$\ \delta_x^+ e\ _{l^2}$	7.382E-03	1.840E-03	4.593E-04	1.145E-04
	Rate	2.004	2.002	2.005	
	$\ e\ _{l^\infty}$	6.889E-03	1.717E-03	4.287E-04	1.069E-04
	Rate	2.004	2.002	2.005	

Table 5: Error analysis of CNCFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x)=0$.

		$h=1/16$ $\tau=2^{-8}$	$h=1/32$ $\tau=2^{-10}$	$h=1/64$ $\tau=2^{-12}$	$h=1/128$ $\tau=2^{-14}$
$\beta=5$	$\ e\ _{l^2}$	3.465E-02	2.171E-03	1.352E-04	7.964E-06
	Rate	3.996	4.005	4.086	
	$\ \delta_x^+ e\ _{l^2}$	4.074E-02	2.558E-03	1.593E-04	9.380E-06
	Rate	3.994	4.005	4.086	
	$\ e\ _{l^\infty}$	2.235E-02	1.399E-03	8.716E-05	5.132E-06
	Rate	3.997	4.005	4.086	
$\beta=-5$	$\ e\ _{l^2}$	3.444E-02	2.159E-03	1.345E-04	7.919E-06
	Rate	3.996	4.005	4.086	
	$\ \delta_x^+ e\ _{l^2}$	2.965E-02	1.859E-03	1.158E-04	6.820E-06
	Rate	3.995	4.005	4.086	
	$\ e\ _{l^\infty}$	2.849E-02	1.786E-03	1.113E-04	6.552E-06
	Rate	3.995	4.005	4.086	

scheme, mathematical induction was used. Extensive numerical results were reported in the last section to confirm our error estimates. In practice, the CNCFD conserves the mass and energy quite well in the discretized level when $\tau = O(h^2)$ but at each step we have to solve a fully nonlinear difference equation which could be expensive in view of computation time cost, especially in 2- d and 3- d . The SICFD scheme is also unconditional stable and it conserves the mass and energy well and only a linear system is required to be solved each step. In addition, both scheme can be solved within at $O(M \log(M))$ time/operations with the help of Discrete Sine Transform (DST). The CNCFD and SICFD methods could be extended to Schrödinger-Poisson-Slater system directly.

Table 6: Error analysis of CNCFD at time $T=0.5$ for different Poisson constant β with external potential $V(x) = \frac{x^2}{2}$.

		$h=1/16$ $\tau=2^{-8}$	$h=1/32$ $\tau=2^{-10}$	$h=1/64$ $\tau=2^{-12}$	$h=1/128$ $\tau=2^{-14}$
$\beta=5$	$\ e\ _{l^2}$	3.558E-02	2.230E-03	1.389E-04	8.178E-06
	Rate	3.996	4.005	4.086	
	$\ \delta_x^+ e\ _{l^2}$	3.234E-02	2.029E-03	1.264E-04	7.440E-06
	Rate	3.995	4.005	4.086	
	$\ e\ _{l^\infty}$	2.382E-02	1.492E-03	9.292E-05	5.471E-06
	Rate	3.997	4.005	4.086	
$\beta=-5$	$\ e\ _{l^2}$	3.410E-02	2.138E-03	1.332E-04	7.840E-06
	Rate	3.996	4.005	4.086	
	$\ \delta_x^+ e\ _{l^2}$	3.547E-02	2.226E-03	1.386E-04	8.163E-06
	Rate	3.995	4.005	4.086	
	$\ e\ _{l^\infty}$	3.019E-02	1.893E-03	1.179E-04	6.945E-06
	Rate	3.995	4.005	4.086	

Table 7: Error analysis of SICFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x)=0$.

		$h=1/16$ $\tau=2^{-8}$	$h=1/32$ $\tau=2^{-10}$	$h=1/64$ $\tau=2^{-12}$	$h=1/128$ $\tau=2^{-14}$
$\beta=5$	$\ e\ _{l^2}$	6.682E-02	4.108E-03	2.551E-04	1.499E-05
	Rate	4.024	4.010	4.089	
	$\ \delta_x^+ e\ _{l^2}$	7.388E-02	4.532E-03	2.814E-04	1.653E-05
	Rate	4.027	4.010	4.089	
	$\ e\ _{l^\infty}$	4.371E-02	2.688E-03	1.669E-04	9.806E-06
	Rate	4.023	4.009	4.089	
$\beta=-5$	$\ e\ _{l^2}$	7.328E-02	4.501E-03	2.796E-04	1.643E-05
	Rate	4.025	4.009	4.089	
	$\ \delta_x^+ e\ _{l^2}$	1.657E-01	4.023E-03	2.498E-04	1.468E-05
	Rate	4.029	4.009	4.089	
	$\ e\ _{l^\infty}$	6.209E-02	3.810E-03	2.366E-04	1.391E-05
	Rate	4.027	4.009	4.089	

Acknowledgements

The author acknowledges Professor Weizhu Bao for suggesting this topic to me and thanks him for stimulating discussions. The author also thanks Prof Huaiyu Jian for continuous encouragement and support. The author is also thankful to the anonymous referees for their constructive remarks. This work was supported by Ministry of Education of Singapore grant R-146-000-120-112 and the National Natural Science Foundation of China Grant 11131005. This work was partially done while the author was visiting the Department of Mathematics, National University of Singapore.

Table 8: Error analysis of SICFD at time $T=0.5$ for different Poisson constant β without external potential, i.e., $V(x) = \frac{x^2}{2}$.

		$h=1/16$ $\tau=2^{-8}$	$h=1/32$ $\tau=2^{-10}$	$h=1/64$ $\tau=2^{-12}$	$h=1/128$ $\tau=2^{-14}$
$\beta=5$	$\ e\ _{l^2}$	6.817E-02	4.189E-03	2.601E-04	1.528E-05
	Rate	4.024	4.010	4.089	
	$\ \delta_x^+ e\ _{l^2}$	5.991E-02	3.674E-03	2.281E-04	1.340E-05
	Rate	4.027	4.010	4.089	
	$\ e\ _{l^\infty}$	4.663E-02	2.867E-03	1.780E-04	1.046E-05
	Rate	4.024	4.009	4.089	
$\beta=-5$	$\ e\ _{l^2}$	7.453E-02	4.578E-03	2.843E-04	1.671E-05
	Rate	4.025	4.009	4.089	
	$\ \delta_x^+ e\ _{l^2}$	8.045E-02	4.930E-03	3.062E-04	1.799E-05
	Rate	4.028	4.010	4.089	
	$\ e\ _{l^\infty}$	6.706E-02	4.115E-03	2.556E-04	1.502E-05
	Rate	4.027	4.010	4.089	

Table 9: Computation time (in seconds) comparison of CNCFD and SICFD scheme at time $T=1.0$ for Poisson constant $\beta=5$ with/without external potential $V(x)$.

		$h=1/16$ $\tau=2^{-8}$	$h=1/32$ $\tau=2^{-10}$	$h=1/64$ $\tau=2^{-12}$	$h=1/128$ $\tau=2^{-14}$
$V(x) = \frac{x^2}{2}$	CNCFD	3.837	15.039	83.029	577.870
	SICFD	1.289	6.140	37.554	284.854
$V(x) = 0$	CNCFD	3.831	15.233	82.103	595.886
	SICFD	0.376	2.228	15.106	110.759

References

- [1] W. Bao and Y.Y. Cai, Optimal error estimate of finite difference methods for the Gross-Pitauskii equation with angular momentum rotation, Math. Comp., to appear.




Figure 1: Evolution of the errors of the discrete mass and energy for SICFD scheme with mesh size $h=1/16$ when external potential $V(x)=0$ and Poisson constant $\beta=5$.

- [2] W. Bao, N.J. Mauser and H.P. Stimming, Effective one particle quantum dynamics of electrons: A numerical study of the Schrödinger-Poisson- $X\alpha$ model, *Comm. Math. Sci.*, 1 (2003), 809-831.
- [3] W. Bao, S. Jin and P. A. Markowich, Time-splitting spectral approximations for the Schrödinger equation in the semiclassical regime, *J. Comput. Phys.*, 175 (2002), 487-524.
- [4] T. Cazenave, *Semilinear Schrödinger equations*, (Courant Lecture Notes in Mathematics vol. 10), New York University, Courant Institute of Mathematical Sciences, AMS, 2003.
- [5] Q. Chang, B. Guo and H. Jiang, Finite Difference Method for Generalized Zakharov Equations, *Math. Comp.*, 64 (1995), 537-553.
- [6] Q. Chang, E. Jia and W. Sun, Difference Schemes for Solving the Generalized Nonlinear Schrödinger Equation, *J. Comput. Phys.*, 148 (1999), 397-415.
- [7] M. De Leo and D. Rial, Well posedness and smoothing effect of Schrödinger-Poisson equation, *J. Math. Phys.*, 48 (2007), 093509.
- [8] C.M. Dion and E. Cancés, Spectral method for the time-dependent Gross-Pitaevskii equation with a harmonic trap, *Phys. Rev. E*, 67 (2003), article 046706.
- [9] X.C. Dong, A short note on simplified pseudospectral methods for computing ground state and dynamics of spherically symmetric Schrödinger-Poisson-Slater system, *J. Comput. Phys.*, 230 (2011), 7917-7922.
- [10] R.T. Galssey, Convergence of an Energy-Preserving Scheme for the Zakharov Equations in one Space Dimension, *Math. Comp.*, 97 (1992), 83-102.
- [11] R. Harrison, I.M. Moroz and K.P. Tod, A numerical study of Schrödinger-Newton equations, *Nonlinearity*, 16 (2003), 101-122.
- [12] R. Landes, On Galerkin's method in the existence theory of quasilinear elliptic equations, *J. Funct. Anal.*, 39 (1980), 123-148.
- [13] S. Larsson and V. Thomée, *Partial differential equations with numerical methods*, Springer, 2009.
- [14] S. Masaki, Energy solution to Schrödinger-Poisson system in the two-dimensional whole space, *SIAM J. Math. Anal.*, 43 (2011), 2719-2731.
- [15] C. Ringhofer and J. Soler, Discrete Schrödinger-Poisson systems preserving energy and mass, *Appl. Math. Lett.*, 13 (2000), 27-32.
- [16] J.M. Sanz-Serna, Methods for the Numerical Solution of the Nonlinear Schroedinger Equation, *Math. Comp.*, 43 (1984), 21-27.
- [17] J. Shen and T. Tang, *Spectral and High-Order Methods with Applications*, Science Press, Beijing, 2006.
- [18] R.K. Shukla, M. Tatineni and X.L. Zhong, Very high-order compact finite difference schemes on non-uniform grids for incompressible Navier-Stokes equations, *J. Comput. Phys.*, 224 (2007), 1064-1094.
- [19] A. Soba, A finite element method solver for time-dependent and stationary Schrödinger equations with a generic potential, *Commun. Comput. Phys.*, 5 (2009), 914-927.
- [20] H. Steinrück, The one-dimensional Wigner-Poisson problem and its relation to the Schrödinger-Poisson problem, 22 (1991), 957-972.
- [21] H.P. Stimming, The IVP for the Schrödinger-Poisson- $X\alpha$ equation in one dimension, *Math. Models Methods Appl. Sci.*, 15 (2005), 1169-1180.
- [22] C. Sutmann, Compact finite difference schemes of sixth order for the Helmholtz equation, *J. Comput. Appl. Math.*, 203 (2007), 15-31.
- [23] I.H. Tan, G.L. Snider, L.D. Chang and E.L. Hu, A self-consistent solution of Schrödinger-Poisson equations using a nonuniform mesh, *J. Appl. Phys.*, 68 (1990), 4071-4076.
- [24] T. Wang, Maximum norm error bound of a linearized difference scheme for a coupled nonlinear Schrödinger equations, *J. Comput. Appl. Math.*, 235 (2011), 4237-4250.
- [25] J. Wang, W. Zhong and J. Zhang, A general meshsize fourth-order compact difference discretization scheme for 3D Poisson equation, *Appl. Math. Comput.*, 183 (2006), 804-812.
- [26] Y. Wang and J. Zhang, Sixth order compact scheme combined with multigrid method and extrapolation technique for 2D poisson equation, *J. Comput. Phys.*, 228 (2009), 137-146.
- [27] S.S. Xie, G.X. Li and S. Yi, Compact finite difference schemes with high accuracy for one-dimensional nonlinear Schrödinger equation, *Comput. Meth. Appl. Mech. Eng.*, 198 (2009), 1052-1060.
- [28] J. Zhang, Multigrid Method and Fourth-Order Compact Scheme for 2D Poisson Equation with Unequal Mesh-Size Discretization, *J. Comput. Phys.*, 179 (2002), 170-179.

- [29] Y. Zhang, Efficient discretization of Schrödinger-Poisson system with artificial boundary condition for Poisson potential in 2D, preprint.
- [30] Y. Zhang and X.C. Dong, On the computation of ground state and dynamics of Schrödinger-Poisson-Slater system, *J. Comput. Phys.*, 230 (2011), 2660-2676.