

1 **BOUNDARY PROBLEMS FOR THE FRACTIONAL AND**
2 **TEMPERED FRACTIONAL OPERATORS***

3 WEIHUA DENG[†], BUYANG LI[‡], WENYI TIAN[§], AND PINGWEN ZHANG[¶]

4 **Abstract.** For characterizing the Brownian motion in a bounded domain: Ω , it is well-known
5 that the boundary conditions of the classical diffusion equation just rely on the given information
6 of the solution along the boundary of a domain; on the contrary, for the Lévy flights or tempered
7 Lévy flights in a bounded domain, it involves the information of a solution in the complementary
8 set of Ω , i.e., $\mathbb{R}^n \setminus \Omega$, with the potential reason that paths of the corresponding stochastic process
9 are discontinuous. Guided by probability intuitions and the stochastic perspectives of anomalous
10 diffusion, we show the reasonable ways, ensuring the clear physical meaning and well-posedness of
11 the partial differential equations (PDEs), of specifying ‘boundary’ conditions for space fractional
12 PDEs modeling the anomalous diffusion. Some properties of the operators are discussed, and the
13 well-posednesses of the PDEs with generalized boundary conditions are proved.

14 **Key words.** Lévy flight; Tempered Lévy flight; Well-posedness; Generalized boundary condi-
15 tions

16 **1. Introduction.** The phrase ‘anomalous is normal’ says that anomalous dif-
17 fusion phenomena are ubiquitous in the natural world. It was first used in the title
18 of [27], which reveals that the diffusion of classical particles on a solid surface has
19 rich anomalous behaviour controlled by the friction coefficient. In fact, anomalous
20 diffusion is no longer a young topic. In the review paper [5], the evolution of par-
21 ticles in disordered environments was investigated; the specific effects of a bias on
22 anomalous diffusion were considered; and the generalizations of Einstein’s relation in
23 the presence of disorder were discussed. With the rapid development of the study
24 of anomalous dynamics in diverse field, some deterministic equations are derived,
25 governing the macroscopic behaviour of anomalous diffusion. In 2000, Metzler and
26 Klafter published the survey paper [25] for the equations governing transport dy-
27 namics in complex system with anomalous diffusion and non-exponential relaxation
28 patterns, i.e., fractional kinetic equations of the diffusion, advection-diffusion, and
29 Fokker-Planck type, derived asymptotically from basic random walk models and a
30 generalized master equation. Many mathematicians have been involved in the re-
31 search of fractional partial differential equations (PDEs). For fractional PDEs in a
32 bounded domain Ω , an important question is how to introduce physically meaningful
33 and mathematically well-posed boundary conditions on $\partial\Omega$ or $\mathbb{R}^n \setminus \Omega$.

34 Microscopically, diffusion is the net movement of particles from a region of higher
35 concentration to a region of lower concentration; for the normal diffusion (Brownian
36 motion), the second moment of the particle trajectories is a linear function of the
37 time t ; naturally, if it is a nonlinear function of t , we call the corresponding diffu-

*This work was partially supported by NSFC under Grant Nos. 11421101, 11421110001, 11626250, 11671182, and the Fundamental Research Funds for the Central Universities under Grant No. lzujbky-2017-ot10. WD thanks Mark M. Meerschaert and Zhen-Qing Chen for the discussions. We thank the anonymous reviewer for Remark 2 of Sec. 2.3.

[†]School of Mathematics and Statistics, Gansu Key Laboratory of Applied Mathematics and Complex Systems, Lanzhou University, Lanzhou 730000, P.R. China. Email: dengwh@lzu.edu.cn

[‡]Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong. Email: buyang.li@polyu.edu.hk

[§]Center for Applied Mathematics, Tianjin University, Tianjin 300072, P.R. China. Email: wenyi.tian@tju.edu.cn

[¶]School of Mathematical Sciences, Laboratory of Mathematics and Applied Mathematics, Peking University, Beijing 100871, P.R. China. Email: pzhang@pku.edu.cn

38 sion process anomalous diffusion or non-Brownian diffusion [25]. The microscopic
39 (stochastic) models describing anomalous diffusion include continuous time random
40 walks (CTRWs), Langevin type equation, Lévy processes, subordinated Lévy pro-
41 cesses, and fractional Brownian motion, etc.. The CTRWs contain two important
42 random variables describing the motion of particles [26], i.e., the waiting time ξ and
43 jump length η . If both the first moment of ξ and the second moment of η are finite
44 the scaling limit, then the CTRWs approximate Brownian motion. On the contrary,
45 if one of them is divergent, then the CTRWs characterize anomalous diffusion. Two
46 of the most important CTRW models are Lévy flights and Lévy walks. For Lévy
47 flights, the ξ with finite first moment and η with infinite second moment are inde-
48 pendent, leading to infinite propagation speed and the divergent second moments of
49 the distribution of the particles. This causes much difficulty in relating the models to
50 experimental data, especially when analyzing the scaling of the measured moments
51 in time [33]. With coupled distribution of ξ and η (the infinite speed is penalized by
52 the corresponding waiting times), we get the so-called Lévy walks [33]. Another idea
53 to ensure that the processes have bounded moments is to truncate the long tailed
54 probability distribution of Lévy flights [22]; they still look like a Lévy flight in not
55 too long a time. Currently, the most popular way to do the truncation is to use the
56 exponential tempering, offering the technical advantage of still being an infinitely di-
57 visible Lévy process after the operation [24]. The Lévy process to describe anomalous
58 diffusion is the scaling limit of CTRWs with independent ξ and η . It is character-
59 ized by its characteristic function. Except Brownian motion with drift, the paths of
60 all other proper Lévy processes are discontinuous. Sometimes, the Lévy flights are
61 conveniently described by the Brownian motion subordinated to a Lévy process [7].
62 Fractional Brownian motions are often taken as the models to characterize subdiffu-
63 sion [21].

64 Macroscopically, fractional (nonlocal) PDEs are the most popular and effective
65 models for anomalous diffusion, derived from the microscopic models. The solution
66 of fractional PDEs is generally the probability density function (PDF) of the position
67 of the particles undergoing anomalous dynamics; with the deepening of research, the
68 fractional PDEs governing the functional distribution of particles' trajectories are also
69 developed [31, 32]. Two ways are usually used to derive the fractional PDEs. One
70 is based on the Montroll-Weiss equation [26], i.e., in Fourier-Laplace space, the PDF
71 $p(\mathbf{X}, t)$ obeys

$$72 \quad (1) \quad \hat{p}(\mathbf{k}, u) = \frac{1 - \phi(u)}{u} \cdot \frac{\hat{p}_0(\mathbf{k})}{1 - \Psi(u, \mathbf{k})},$$

73 where $\hat{p}_0(\mathbf{k})$ is the Fourier transform of the initial data; $\phi(u)$ is the Laplace transform
74 of the PDF of waiting times ξ and $\Psi(u, \mathbf{k})$ the Laplace and the Fourier transforms of
75 the joint PDF of waiting times ξ and jump length η . If ξ and η are independent, then
76 $\Psi(u, \mathbf{k}) = \phi(u)\psi(\mathbf{k})$, where $\psi(\mathbf{k})$ is the Fourier transform of the PDF of η . Another
77 way is based on the characteristic function of the α -stable Lévy motion, being the
78 scaling limit of the CTRW model with power law distribution of jump length η . In
79 the high dimensional case, it is more convenient to make the derivation by using the
80 characteristic function of the stochastic process. According to the Lévy-Khinchin
81 formula [2], the characteristic function of Lévy process has a specific form

$$82 \quad (2) \quad \int_{\mathbb{R}^n} e^{i\mathbf{k}\cdot\mathbf{X}} p(\mathbf{X}, t) d\mathbf{X} = \mathbf{E}(e^{i\mathbf{k}\cdot\mathbf{X}}) = e^{t\Phi(\mathbf{k})},$$

where

$$\Phi(\mathbf{k}) = i\mathbf{a} \cdot \mathbf{k} - \frac{1}{2}(\mathbf{k} \cdot \mathbf{b}\mathbf{k}) + \int_{\mathbb{R}^n \setminus \{0\}} [e^{i\mathbf{k} \cdot \mathbf{X}} - 1 - i(\mathbf{k} \cdot \mathbf{X})\chi_{\{|\mathbf{X}| < 1\}}] \nu(d\mathbf{X});$$

83 here χ_I is the indicator function of the set I , $\mathbf{a} \in \mathbb{R}^n$, \mathbf{b} is a positive definite symmetric
 84 $n \times n$ matrix and ν is a sigma-finite Lévy measure on $\mathbb{R}^n \setminus \{0\}$. When \mathbf{a} and \mathbf{b} are
 85 zero and

$$86 \quad (3) \quad \nu(d\mathbf{X}) = \frac{\beta\Gamma(\frac{n+\beta}{2})}{2^{1-\beta}\pi^{n/2}\Gamma(1-\beta/2)} |\mathbf{X}|^{-\beta-n} d\mathbf{X},$$

87 the process is a rotationally symmetric β -stable Lévy motion and its PDF solves

$$88 \quad (4) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = \Delta^{\beta/2} p(\mathbf{X}, t),$$

89 where $\mathcal{F}(\Delta^{\beta/2} p(\mathbf{X}, t)) = -|\mathbf{k}|^\beta \mathcal{F}(p(\mathbf{X}, t))$ [29]. If replacing (3) by the measure of
 90 isotropic tempered power law with the tempering exponent λ , then we get the corre-
 91 sponding PDF evolution equation

$$92 \quad (5) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t),$$

93 where $(\Delta + \lambda)^{\beta/2}$ is defined by (32) in physical space and by (34) in Fourier space.

94 In practice, the choice of $\nu(d\mathbf{X})$ depends strongly on the concrete physical envi-
 95 ronment. For example, Figure 1 clearly shows the horizontal and vertical structure.
 96 So, we need to take the measure as (if it is superdiffusion)

$$97 \quad (6) \quad \nu(d\mathbf{X}) = \nu(d\mathbf{x}_1 d\mathbf{x}_2) = \frac{\beta_1\Gamma(\frac{1+\beta_1}{2})}{2^{1-\beta_1}\pi^{1/2}\Gamma(1-\beta_1/2)} |\mathbf{x}_1|^{-\beta_1-1} \delta(\mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2 \\ + \frac{\beta_2\Gamma(\frac{1+\beta_2}{2})}{2^{1-\beta_2}\pi^{1/2}\Gamma(1-\beta_2/2)} \delta(\mathbf{x}_1) |\mathbf{x}_2|^{-\beta_2-1} d\mathbf{x}_1 d\mathbf{x}_2,$$

98 where β_1 and β_2 belong to $(0, 2)$. If \mathbf{a} and \mathbf{b} equal to zero, then it leads to diffusion
 99 equation

$$100 \quad (7) \quad \frac{\partial p(\mathbf{x}_1, \mathbf{x}_2, t)}{\partial t} = \frac{\partial^{\beta_1} p(\mathbf{x}_1, \mathbf{x}_2, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \mathbf{x}_2, t)}{\partial |\mathbf{x}_2|^{\beta_2}}.$$

101 Under the guidelines of probability intuitions and stochastic perspectives [17] of
 102 Lévy flights or tempered Lévy flights, we discuss the reasonable ways of defining
 103 fractional partial differential operators and specifying the ‘boundary’ conditions for
 104 their macroscopic descriptions, i.e., the PDEs of the types Eqs. (4), (5), (7), and
 105 their extensions, e.g., the fractional Feynman-Kac equations [31, 32]. For the related
 106 discussions on the nonlocal diffusion problems from a mathematical point of view,
 107 one can see the review paper [11]. The divergence of the second moment and the
 108 discontinuity of the paths of Lévy flights predicate that the corresponding diffusion
 109 operators should defined on \mathbb{R}^n , which further signify that if we are solving the equa-
 110 tions in a bounded domain Ω , the information in $\mathbb{R}^n \setminus \Omega$ should also be involved. We
 111 will show that the generalized Dirichlet type boundary conditions should be specified
 112 as $p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$. If the particles are killed after leaving the domain Ω , then



Fig. 1: Sketch map for the physical environment suitable for Eq. (7).

113 $g(\mathbf{X}, t) \equiv 0$, i.e., the so-called absorbing boundary conditions. Because of the dis-
 114 continuity of the jumps of Lévy flights, a particular concept ‘escape probability’ can
 115 be introduced, which means the probability that the particle jumps from the domain
 116 Ω into a domain $H \subset \mathbb{R}^n \setminus \Omega$; for solving the escape probability, one just needs to
 117 specify $g(\mathbf{X}) = 1$ for $\mathbf{X} \in H$ and 0 for $\mathbf{X} \in (\mathbb{R}^n \setminus \Omega) \setminus H$ for the corresponding time-
 118 independent PDEs. As for the generalized Neumann type boundary conditions, our
 119 ideas come from the fact that the continuity equation (conservation law) holds for
 120 any kinds of diffusion, since the particles can not be created or destroyed. Based on
 121 the continuity equation and the governing equation of the PDF of Lévy or tempered
 122 Lévy flights, the corresponding flux \mathbf{j} can be obtained. So the generalized reflecting
 123 boundary conditions should be $\mathbf{j}|_{\mathbb{R}^n \setminus \Omega} \equiv 0$, which implies $(\nabla \cdot \mathbf{j})|_{\mathbb{R}^n \setminus \Omega} \equiv 0$. Then, the
 124 generalized Neumann type boundary conditions are given as $(\nabla \cdot \mathbf{j})|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$,
 125 e.g., for (4), it should be taken as $(\Delta^{\beta/2} p(\mathbf{X}, t))|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$. The well-posednesses
 126 of the equations under our specified generalized Dirichlet or Neumann type boundary
 127 conditions are well established.

128 Overall, this paper focuses on introducing physically reasonable boundary con-
 129 straints for a large class of fractional PDEs, building a bridge between the physical and
 130 mathematical communities for studying anomalous diffusion and fractional PDEs. In
 131 the next section, we recall the derivation of fractional PDEs. Some new concepts are
 132 introduced, such as the tempered fractional Laplacian, and some properties of anoma-
 133 lous diffusion are found. In Sec. 3, we discuss the reasonable ways of specifying the
 134 generalized boundary conditions for the fractional PDEs governing the position or
 135 functional distributions of Lévy flights and tempered Lévy flights. In Sec. 4, we prove
 136 well-posedness of the fractional PDEs under the generalized Dirichlet and Neumann
 137 boundary conditions defined on the complement of the bounded domain. Conclusion
 138 and remarks are given in the last section.

139 **2. Preliminaries.** For well understanding and inspiring the ways of specifying
 140 the ‘boundary constrains’ to PDEs governing the PDF of Lévy flights or tempered
 141 Lévy flights, we will show the ideas of deriving the microscopic and macroscopic
 142 models.

143 **2.1. Microscopic models for anomalous diffusion.** For the microscopic de-
 144 scription of the anomalous diffusion, we consider the trajectory of a particle or a
 145 stochastic process, i.e., $\mathbf{X}(t)$. If $\langle |\mathbf{X}(t)|^2 \rangle \sim t$, the process is normal, otherwise it is
 146 abnormal. The anomalous diffusions of most often happening in natural world are
 147 the cases that $\langle |\mathbf{X}(t)|^2 \rangle \sim t^\gamma$ with $\gamma \in [0, 1) \cup (1, 2]$. A Lévy flight is a random walk
 148 in which the jump length has a heavy tailed (power law) probability distribution,




Fig. 2: Random trajectories (1000 steps) of Lévy flight ($\beta = 0.8$), tempered Lévy flight ($\beta = 0.8, \lambda = 0.2$), and Brownian motion.

149 i.e., the PDF of jump length r is like $r^{-\beta-n}$ with $\beta \in (0, 2)$, and the distribution
 150 in direction is uniform. With the wide applications of Lévy flights in characterizing
 151 long-range interactions [3] or a nontrivial “crumpled” topology of a phase (or con-
 152 figuration) space of polymer systems [30], etc, its second and higher moments are
 153 divergent, leading to the difficulty in relating models to experimental data. In fact,
 154 for Lévy flights $\langle |\mathbf{X}(t)|^\delta \rangle \sim t^{\delta/\beta}$ with $0 < \delta < \beta \leq 2$. Under the framework of CTRW,
 155 the model Lévy walk [28] can circumvent this obstacle by putting a larger time cost
 156 to a longer displacement, i.e., using the space-time coupled jump length and waiting
 157 time distribution $\Psi(r, t) = \frac{1}{2}\delta(r - vt)\phi(t)$. Another popular model is the so-called
 158 tempered Lévy flights [18], in which the extremely long jumps is exponentially cut
 159 by using the distribution of jump length $e^{-r\lambda}r^{-\beta-n}$ with λ being a small modulation
 160 parameter (a smooth exponential regression towards zero). In not too long a time,
 161 the tempered Lévy flights display the dynamical behaviors of Lévy flights, ultraslowly
 162 converging to the normal diffusion. Figure 2 shows the trajectories of 1000 steps of
 163 Lévy flights, tempered Lévy flights, and Brownian motion in two dimensions; note
 164 the presence of rare but large jumps compared to the Brownian motion, playing the
 165 dominant role in the dynamics.

Using Berry-Esséen theorem [13], first established in 1941, which applies to the
 convergence to a Gaussian for a symmetric random walk whose jump probabilities
 have a finite third moment, we have that for the one dimensional tempered Lévy
 flights with the distribution of jump length $Ce^{-r\lambda}r^{-\beta-1}$ the convergence speed is

$$\frac{5}{2\sqrt{2C}} \frac{\Gamma(3-\beta)}{\Gamma(2-\beta)^{3/2}} \lambda^{-\frac{1}{2}\beta} \frac{1}{\sqrt{m}},$$

166 which means that the scaling law for the number of steps needed for Gaussian behavior
 167 to emerge as

$$168 \quad (8) \quad m \sim \lambda^{-\beta}.$$

169 More concretely, letting $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_m$ be i.i.d. random variables with PDF
 170 $Ce^{-r\lambda}r^{-\beta-1}$ and $E(|\mathbf{X}_1|^2) = \sigma^2 > 0$, then the cumulative distribution function
 171 (CDF) Q_m of $\mathbf{Y}_m = (\mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_m)/(\sigma\sqrt{m})$ converges to the CDF $Q(\mathbf{X})$
 172 of the standard normal distribution as

$$173 \quad |Q_m(\mathbf{X}) - Q(\mathbf{X})| < \frac{5}{2} \frac{\langle |\mathbf{X}|^3 \rangle}{\langle |\mathbf{X}|^2 \rangle^{3/2}} \frac{1}{\sqrt{m}} = \frac{5}{2\sqrt{2C}} \frac{\Gamma(3-\beta)}{\Gamma(2-\beta)^{3/2}} \lambda^{-\frac{1}{2}\beta} \frac{1}{\sqrt{m}},$$

174 since

$$175 \quad \langle |\mathbf{X}|^3 \rangle = C \int_{-\infty}^{\infty} |\mathbf{X}|^3 e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{-\beta-1} d|\mathbf{X}| = 2C \int_0^{\infty} e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{3-\beta-1} d|\mathbf{X}| = 2C\lambda^{\beta-3}\Gamma(3-\beta)$$

176 and

$$177 \quad \langle |\mathbf{X}|^2 \rangle = C \int_{-\infty}^{\infty} |\mathbf{X}|^2 e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{-\beta-1} d|\mathbf{X}| = 2C \int_0^{\infty} e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{2-\beta-1} d|\mathbf{X}| = 2C\lambda^{\beta-2}\Gamma(2-\beta).$$

178 From Eq. (8), it can be seen that with the decrease of λ , the required m for the
 179 crossover between Lévy flight behavior and Gaussian behavior increase rapidly. A
 180 little bit counterintuitive observation is that the number of variables required to the
 181 crossover increases with the increase of β .

182 We have described the distributions of jump length for Lévy flights and tempered
 183 Lévy flights, in which Poisson process is taken as the renewal process. We denote the
 184 Poisson process with rate $\zeta > 0$ as $N(t)$ and its waiting time distribution between two
 185 events is $\zeta e^{-\zeta t}$. Then the Lévy flights or tempered Lévy flights are the compound

186 Poisson process defined as $\mathbf{X}(t) = \sum_{j=0}^{N(t)} \mathbf{X}_j$, where \mathbf{X}_j are i.i.d. random variables with
 187 the distribution of power law or tempered power law. The characteristic function of
 188 $\mathbf{X}(t)$ can be calculated as follows. For real \mathbf{k} , we have

$$\begin{aligned} \hat{p}(\mathbf{k}, t) &= \mathbf{E}(e^{i\mathbf{k}\cdot\mathbf{X}(t)}) \\ &= \sum_{j=0}^{\infty} \mathbf{E}(e^{i\mathbf{k}\cdot\mathbf{X}(t)} | N(t) = j) P(N(t) = j) \\ 189 \quad (9) \quad &= \sum_{j=0}^{\infty} \mathbf{E}(e^{i\mathbf{k}\cdot(\mathbf{X}_0+\mathbf{X}_1+\dots+\mathbf{X}_j)} | N(t) = j) P(N(t) = j) \\ &= \sum_{j=0}^{\infty} \Phi_0(\mathbf{k})^j \frac{(\zeta t)^j}{j!} e^{-\zeta t} \\ &= e^{\zeta t(\Phi_0(\mathbf{k})-1)}, \end{aligned}$$

190 where $\Phi_0(\mathbf{k}) = \mathbf{E}(e^{i\mathbf{k}\cdot\mathbf{X}_0})$, being also the characteristic function of $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_j$
 191 since they are i.i.d.

192 In the CTRW model describing one dimensional Lévy flights or tempered Lévy
 193 flights, the PDF of waiting times is taken as $\zeta e^{-\zeta t}$ with its Laplace transform $\zeta/(u+\zeta)$
 194 and the PDF of jumping length is $c^{-\beta} r^{-\beta-1}$ or $e^{-\lambda r} r^{-\beta-1}$ with its Fourier transform
 195 $1 - c^\beta |k|^\beta$ or $1 - c_{\beta,\lambda}[(\lambda + ik)^\beta - \lambda^\beta] - c_{\beta,\lambda}[(\lambda - ik)^\beta - \lambda^\beta]$. Substituting them into
 196 the Montroll-Weiss Eq. (1) with $\hat{p}_0(k) = 1$ (the initial position of particles is at zero),
 197 we get that $\hat{p}(k, u)$ of Lévy flights solves

$$198 \quad (10) \quad \hat{p}(k, u) = \frac{1}{u + \zeta c^\beta |k|^\beta};$$

199 and the $\hat{p}(k, u)$ of tempered Lévy flights obeys

$$200 \quad (11) \quad \hat{p}(k, u) = \frac{1}{u + \zeta C_{\beta,\lambda}[(\lambda + ik)^\beta - \lambda^\beta] + \zeta C_{\beta,\lambda}[(\lambda - ik)^\beta - \lambda^\beta]}.$$

201 If the subdiffusion is involved, we need to choose the PDF of waiting times as
 202 $\tilde{c}^{1+\alpha}t^{-\alpha-1}$ with $\alpha \in (0, 1)$ and its Laplace transform $1 - \tilde{c}^\alpha u^\alpha$. Then from (1),
 203 we get that

$$204 \quad (12) \quad \hat{p}(k, u) = \frac{\tilde{c}^\alpha}{u^{1-\alpha}(1 - (1 - \tilde{c}_\alpha u^\alpha)\psi(k))}.$$

205 For high dimensional case, the Lévy flights can also be characterized by Brow-
 206 nian motion subordinated to a Lévy process. Let $\mathbf{Y}(t)$ be a Brownian motion with
 207 Fourier exponent $-|\mathbf{k}|^2$ and $S(t)$ a subordinator with Laplace exponent $u^{\beta/2}$ that is
 208 independent of $\mathbf{Y}(t)$. The process $\mathbf{X}(t) = \mathbf{Y}(S(t))$ is describing Lévy flights with
 209 Fourier exponent $-|\mathbf{k}|^\beta$, being the subordinate process of $\mathbf{Y}(t)$. In effect, denote the
 210 characteristic function of $\mathbf{Y}(t)$ as $\Phi_y(\mathbf{k})$ and the one of $S(t)$ as $\Phi_s(u)$. Then the
 211 characteristic function of $\mathbf{X}(t)$ is as follows:

$$\begin{aligned} \hat{p}_x(\mathbf{k}, t) &= \int_{\mathbb{R}^n} e^{i\mathbf{k}\cdot\mathbf{X}} p_x(\mathbf{X}, t) d\mathbf{X} \\ &= \int_0^\infty \int_{\mathbb{R}^n} e^{i\mathbf{k}\cdot\mathbf{Y}} p_y(\mathbf{Y}, \tau) d\mathbf{Y} p_s(\tau, t) d\tau \\ &= \int_0^\infty e^{-\tau(-\Phi_y(\mathbf{k}))} p_s(\tau, t) d\tau \\ &= e^{-t\Phi_s(-\Phi_y(\mathbf{k}))}, \end{aligned}$$

212 (13)

213 where p_x , p_y , and p_s , are respectively the PDFs of the stochastic processes \mathbf{X} , \mathbf{Y} , and
 214 S . Similarly, in the following, we denote p with subscript (lowercase letter) as the
 215 PDF of the corresponding stochastic process (uppercase letter).

216 This paper mainly focuses on Lévy flights and tempered Lévy flights. If one is
 217 interested in subdiffusion, instead of Poisson process, the fractional Poisson process
 218 should be taken as the renewal process, in which the time interval between each pair
 219 of events follows the power law distribution. Let $\mathbf{Y}(t)$ be a general Lévy process
 220 with Fourier exponent $\Phi_y(\mathbf{k})$ and $S(t)$ a strictly increasing subordinator with Laplace
 221 exponent u^α ($\alpha \in (0, 1)$). Define the inverse subordinator $E(t) = \inf\{\tau > 0 : S(\tau) >$
 222 $t\}$. Since $t = S(\tau)$ and $\tau = E(t)$ are inverse processes, we have $P(E(t) \leq \tau) =$
 223 $P(S(\tau) \geq t)$. Hence

$$224 \quad (14) \quad p_e(\tau, t) = \frac{\partial P(E(t) \leq \tau)}{\partial \tau} = \frac{\partial}{\partial \tau} [1 - P(S(\tau) < t)] = -\frac{\partial}{\partial \tau} \int_0^t p_s(y, \tau) dy.$$

225 In the above equation, taking Laplace transform w.r.t t leads to

$$226 \quad (15) \quad p_e(\tau, u) = -\frac{\partial}{\partial \tau} u^{-1} e^{-\tau u^\alpha} = u^{\alpha-1} e^{-\tau u^\alpha}.$$

227 For the PDF $p_x(\mathbf{X}, t)$ of $\mathbf{X}(t) = \mathbf{Y}(E(t))$, there holds

$$228 \quad (16) \quad p_x(\mathbf{X}, t) = \int_0^\infty p_y(\mathbf{X}, \tau) p_e(\tau, t) d\tau.$$

229 Performing Fourier transform w.r.t. \mathbf{X} and Laplace transform w.r.t. t to the above

230 equation results in

$$\begin{aligned}
 \hat{p}_x(\mathbf{k}, u) &= \int_0^\infty \hat{p}_y(\mathbf{k}, \tau) p_e(\tau, u) d\tau \\
 &= \int_0^\infty e^{-\tau \Phi_y(\mathbf{k})} u^{\alpha-1} e^{-\tau u^\alpha} d\tau \\
 &= \frac{u^{\alpha-1}}{u^\alpha + \Phi_y(\mathbf{k})}.
 \end{aligned}
 \tag{17}$$

232 **Remark.** According to Fogedby [15], the stochastic trajectories of (scale limited)
 233 CTRW $\mathbf{X}(E_t)$ can also be expressed in terms of the coupled Langevin equation

$$\begin{cases} \dot{\mathbf{X}}(\tau) = F(\mathbf{X}(\tau)) + \eta(\tau), \\ \dot{S}(\tau) = \xi(\tau), \end{cases}
 \tag{18}$$

235 where $F(\mathbf{X})$ is a vector field; E_t is the inverse process of $S(t)$; the noises $\eta(\tau)$ and
 236 $\xi(\tau)$ are statistically independent, corresponding to the distributions of jump length
 237 and waiting times.

238 **2.2. Derivation of the macroscopic description from the microscopic**
 239 **models.** This section focuses on the derivation of the deterministic equations gov-
 240 erning the PDF of position of the particles undergoing anomalous diffusion. It shows
 241 that the operators related to (tempered) power law jump lengths should be defined
 242 on the whole unbounded domain \mathbb{R}^n , which can also be inspired by the rare but ex-
 243 tremely long jump lengths displayed in Figure 2; the fact that among all proper Lévy
 244 processes Brownian motion is the unique one with continuous paths further consol-
 245 idates the reasonable way of defining the operators. We derive the PDEs based on
 246 Eqs. (9), (13), and (16), since they apply for both one and higher dimensional cases.
 247 For one dimensional case, sometimes it is convenient to use (10), (11), and (12).

248 When the diffusion process is rotationally symmetric β -stable, i.e., it is isotropic
 249 with PDF of jump length $c_{\beta,n} r^{-\beta-n}$ and its Fourier transform $1 - |\mathbf{k}|^\beta$, where n is
 250 the space dimension. In Eq. (9), taking ζ equal to 1, we get the Cauchy equation

$$\frac{d\hat{p}(\mathbf{k}, t)}{dt} = -|\mathbf{k}|^\beta \hat{p}(\mathbf{k}, t).
 \tag{19}$$

252 Performing inverse Fourier transform to the above equation leads to

$$\frac{\partial p(\mathbf{X}, t)}{\partial t} = \Delta^{\beta/2} p(\mathbf{X}, t),
 \tag{20}$$

254 where

$$\begin{aligned}
 \Delta^{\beta/2} p(\mathbf{X}, t) &= -c_{n,\beta} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C} B_\varepsilon(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 &= \frac{1}{2} c_{n,\beta} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{|\mathbf{Y}|^{n+\beta}} d\mathbf{Y}
 \end{aligned}
 \tag{21}$$

256 with [9]

$$c_{n,\beta} = \frac{\beta \Gamma(\frac{n+\beta}{2})}{2^{1-\beta} \pi^{n/2} \Gamma(1 - \beta/2)}.
 \tag{22}$$

258 For the more general cases of Eq. (9), there is the Cauchy equation

$$259 \quad (23) \quad \frac{d\hat{p}(\mathbf{k}, t)}{dt} = (\Phi_0(\mathbf{k}) - 1)\hat{p}(\mathbf{k}, t),$$

260 so the PDF of the stochastic process \mathbf{X} solves (taking $\zeta = 1$)

$$261 \quad (24) \quad \begin{aligned} \frac{\partial p(\mathbf{X}, t)}{\partial t} &= \mathcal{F}^{-1}\{(\Phi_0(\mathbf{k}) - 1)\hat{p}(\mathbf{k}, t)\} \\ &= \int_{\mathbb{R}^n \setminus \{0\}} [p(\mathbf{X} + \mathbf{Y}, t) - p(\mathbf{X}, t)] \nu(d\mathbf{Y}), \end{aligned}$$

where $\nu(d\mathbf{Y})$ is the probability measure of the jump length. Sometimes, to overcome the possible divergence of the terms on the right hand side of Eq. (24) because of the possible strong singularity of $\nu(d\mathbf{Y})$ at zero, the term

$$\Phi_0(\mathbf{k}) - 1 = \int_{\mathbb{R}^n \setminus \{0\}} [e^{i\mathbf{k} \cdot \mathbf{Y}} - 1] \nu(d\mathbf{Y})$$

262 is approximately replaced by

$$263 \quad (25) \quad \int_{\mathbb{R}^n \setminus \{0\}} [e^{i\mathbf{k} \cdot \mathbf{Y}} - 1 - i(\mathbf{k} \cdot \mathbf{Y})_{\chi_{\{|\mathbf{Y}| < 1\}}}] \nu(d\mathbf{Y});$$

264 then the corresponding modification to Eq. (24) is

$$265 \quad (26) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = \int_{\mathbb{R}^n \setminus \{0\}} \left[p(\mathbf{X} + \mathbf{Y}, t) - p(\mathbf{X}, t) - \sum_{i=1}^n \mathbf{y}_i (\partial_i p(\mathbf{X}, t))_{\chi_{\{|\mathbf{Y}| < 1\}}} \right] \nu(d\mathbf{Y}),$$

266 where \mathbf{y}_i is the component of \mathbf{Y} , i.e., $\mathbf{Y} = \{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n\}^T$. If $\nu(-d\mathbf{Y}) = \nu(d\mathbf{Y})$,
267 the integration of the summation term of above equation equals to zero.

268 If the diffusion is in the environment having a structure like Figure 1, the proba-
269 bility measure should be taken as

$$(27) \quad \begin{aligned} \nu(d\mathbf{X}) &= \nu(dx_1 dx_2 dx_3 \cdots dx_n) \\ &= \frac{\beta_1 \Gamma(\frac{1+\beta_1}{2})}{2^{1-\beta_1} \pi^{1/2} \Gamma(1 - \beta_1/2)} |\mathbf{x}_1|^{-\beta_1-1} \delta(\mathbf{x}_2) \delta(\mathbf{x}_3) \cdots \delta(\mathbf{x}_n) dx_1 dx_2 dx_3 \cdots dx_n \\ 270 \quad &+ \frac{\beta_2 \Gamma(\frac{1+\beta_2}{2})}{2^{1-\beta_2} \pi^{1/2} \Gamma(1 - \beta_2/2)} |\mathbf{x}_2|^{-\beta_2-1} \delta(\mathbf{x}_1) \delta(\mathbf{x}_3) \cdots \delta(\mathbf{x}_n) dx_1 dx_2 dx_3 \cdots dx_n + \cdots \\ &+ \frac{\beta_n \Gamma(\frac{1+\beta_n}{2})}{2^{1-\beta_n} \pi^{1/2} \Gamma(1 - \beta_n/2)} |\mathbf{x}_n|^{-\beta_n-1} \delta(\mathbf{x}_1) \delta(\mathbf{x}_2) \cdots \delta(\mathbf{x}_{n-1}) dx_1 dx_2 dx_3 \cdots dx_n, \end{aligned}$$

271 where $\beta_1, \beta_2, \dots, \beta_n$ belong to $(0, 2)$. Plugging Eq. (27) into Eq. (24) leads to
(28)

$$272 \quad \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2}} + \cdots + \frac{\partial^{\beta_n} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n}},$$

where

$$\mathcal{F}_j \left(\frac{\partial^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|^{\beta_j}} \right) = -|\mathbf{k}_j|^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{k}_j, \mathbf{x}_{j+1}, \dots, \mathbf{x}_n, t)$$

273 and $\frac{\partial^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|^{\beta_j}}$ in physical space is defined by (21) with $n = 1$; in particular, when
 274 $\beta_j \in (1, 2)$, it can also be written as

(29)
 275
$$\frac{\partial^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|^{\beta_j}} = -\frac{1}{2 \cos(\beta_j \pi / 2) \Gamma(2 - \beta_j)} \frac{\partial^2}{\partial \mathbf{x}_j^2} \int_{-\infty}^{\infty} |\mathbf{x}_j - \mathbf{y}|^{1 - \beta_j} p(\mathbf{x}_1, \dots, \mathbf{y}, \dots, \mathbf{x}_n, t) d\mathbf{y}.$$

It should be emphasized here that when characterizing diffusion processes related with Lévy flights the operators should be defined in the whole space. Another issue that also should be stressed is that when $\beta_1 = \beta_2 = \dots = \beta_n = 1$, Eq. (28) is still describing the phenomena of anomalous diffusion, including the cases that they belong to $(0, 1)$; the corresponding ‘first’ order operator is nonlocal, being different from the classical first order operator, but they have the same energy in the sense that

$$\begin{aligned} & \mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|} \right) \overline{\mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|} \right)} \\ &= (k_j)^2 \hat{p}^2(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{k}_j, \mathbf{x}_{j+1}, \dots, \mathbf{x}_n, t) \\ &= \mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial \mathbf{x}_j} \right) \overline{\mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial \mathbf{x}_j} \right)}; \\ & \mathcal{F} \left(\Delta^{1/2} p(\mathbf{X}, t) \right) \overline{\mathcal{F} \left(\Delta^{1/2} p(\mathbf{X}, t) \right)} \\ &= |\mathbf{k}|^2 \hat{p}^2(\mathbf{k}, t) = \mathcal{F}(\nabla p(\mathbf{X}, t)) \cdot \overline{\mathcal{F}(\nabla p(\mathbf{X}, t))}, \end{aligned}$$

276 even though $\Delta^{1/2}$ and ∇ are completely different operators, where the notation \bar{v}
 277 stands for the complex conjugate of v .

278 If the subdiffusion is involved, the derivation of the macroscopic equation should
 279 be based on Eq. (17). For getting the term related to time derivative, the inverse
 280 Laplace transform should be performed on $u^\alpha \hat{p}(\mathbf{k}, u) - u^{\alpha-1}$. Since $\hat{p}(\mathbf{k}, t = 0)$ is taken
 281 as 1, there exists

282 (30)
$$\mathcal{L}^{-1}(u^\alpha \hat{p}(\mathbf{k}, u) - u^{\alpha-1}) = \frac{1}{\Gamma(1 - \alpha)} \int_0^t (t - \tau)^{-\alpha} \frac{\partial \hat{p}(\mathbf{k}, \tau)}{\partial \tau} d\tau,$$

283 which is usually denoted as ${}_0^C D_t^\alpha \hat{p}(\mathbf{k}, t)$, the so-called Caputo fractional derivative. So,
 284 if both the PDFs of the waiting time and jump lengths of the stochastic process \mathbf{X} were
 285 power law, the corresponding models can be obtained by replacing $\frac{\partial}{\partial t}$ with ${}_0^C D_t^\alpha$ in
 286 Eqs. (20), (24), (26), and (28). Furthermore, if there is an external force $F(\mathbf{X})$ in the
 287 considered stochastic process \mathbf{X} , we need to add an additional term $\nabla \cdot (F(\mathbf{X})p(\mathbf{X}, t))$
 288 on the right hand side of Eqs. (20), (24), (26), and (28).

289 Here we turn to another important and interesting topic: tempered Lévy flights.
 290 Practically it is not easy to collect the value of a function in the unbounded area
 291 $\mathbb{R}^n \setminus \Omega$. This is one of the achievements of using tempered fractional Laplacian. It is
 292 still isotropic but with PDF of jump length $c_{\beta, n, \lambda} e^{-\lambda r} r^{-\beta-n}$. The PDF of tempered
 293 Lévy flights solves

294 (31)
$$\frac{\partial p(\mathbf{X}, t)}{\partial t} = (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t),$$

295 where

$$\begin{aligned}
 (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t) &= -c_{n,\beta,\lambda} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\varepsilon}(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{e^{\lambda|\mathbf{X}-\mathbf{Y}|} |\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 (32) \quad &= \frac{1}{2} c_{n,\beta,\lambda} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y}
 \end{aligned}$$

297 with

$$(33) \quad c_{n,\beta,\lambda} = \frac{\Gamma(\frac{n}{2})}{2\pi^{\frac{n}{2}} |\Gamma(-\beta)|}.$$

299 The choice of the constant as the one given in (33) leads to

$$(34) \quad \mathcal{F}\left((\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t)\right) = (-1)^{\lfloor \beta \rfloor} (\lambda^\beta - (\lambda^2 + |\mathbf{k}|^2)^{\frac{\beta}{2}} + O(|\mathbf{k}|^2)) \hat{p}(\mathbf{k}, t),$$

301 where $\beta \in (0, 1) \cup (1, 2)$, and $\lfloor \beta \rfloor$ means the biggest integer, being smaller than or
 302 equal to β .

303 However, if $\lambda = 0$, one needs to choose the constant as the one given in (22) to
 304 make sure $\mathcal{F}\left(\Delta^{\beta/2} p(\mathbf{X}, t)\right) = -|\mathbf{k}|^\beta \hat{p}(\mathbf{k}, t)$. The reason is as follows.

$$\mathcal{F}\left((\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t)\right) = \frac{1}{2} c_{n,\beta,\lambda} \int_{\mathbb{R}^n} \frac{e^{i\mathbf{k}\cdot\mathbf{Y}} + e^{-i\mathbf{k}\cdot\mathbf{Y}} - 2}{|\mathbf{Y}|^{n+\beta}} e^{-\lambda|\mathbf{Y}|} d\mathbf{Y} \cdot \mathcal{F}(p(\mathbf{X}, t))$$

305

$$= -c_{n,\beta,\lambda} \int_{\mathbb{R}^n} \frac{1 - \cos(\mathbf{k} \cdot \mathbf{Y})}{|\mathbf{Y}|^{n+\beta}} e^{-\lambda|\mathbf{Y}|} d\mathbf{Y} \cdot \mathcal{F}(p(\mathbf{X}, t)).$$

306 For $\beta \in (0, 1) \cup (1, 2)$, then we have

$$\begin{aligned}
& \int_{\mathbb{R}^n} \frac{1 - \cos(\mathbf{k} \cdot \mathbf{Y})}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y} = \int_{\mathbb{R}^n} \frac{1 - \cos(|\mathbf{k}|\mathbf{y}_1)}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y} = |\mathbf{k}|^\beta \int_{\mathbb{R}^n} \frac{1 - \cos(\mathbf{x}_1)}{|\mathbf{X}|^{n+\beta}} e^{-\frac{\lambda}{|\mathbf{k}|}|\mathbf{X}|} d\mathbf{X} \\
& = C|\mathbf{k}|^\beta \int_0^\infty \frac{1}{r^{n+\beta}} e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{n-1} \left(\int_0^\pi (1 - \cos(r \cos \theta_1)) \sin^{n-2}(\theta_1) d\theta_1 \right) dr \\
& = \frac{1}{(-\beta)(-\beta+1)} C|\mathbf{k}|^{\beta-2} \lambda^2 \int_0^\infty e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{-\beta+1} \left(\int_0^\pi (1 - \cos(r \cos \theta_1)) \sin^{n-2}(\theta_1) d\theta_1 \right) dr \\
& \quad - \frac{1}{(-\beta)(-\beta+1)} C|\mathbf{k}|^{\beta-1} \lambda \int_0^\infty e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{-\beta+1} \left(\int_0^\pi \sin(r \cos \theta_1) \sin^{n-2}(\theta_1) \cos(\theta_1) d\theta_1 \right) dr \\
& \quad - \frac{1}{-\beta} C|\mathbf{k}|^\beta \int_0^\infty e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{-\beta} \left(\int_0^\pi \sin(r \cos \theta_1) \sin^{n-2}(\theta_1) \cos(\theta_1) d\theta_1 \right) dr \\
307 & = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \lambda^\beta \left[1 - {}_2F_1\left(\frac{2-\beta}{2}, \frac{3-\beta}{2}; \frac{n}{2}; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right. \\
& \quad \left. - \frac{2-\beta}{n} \frac{|\mathbf{k}|^2}{\lambda^2} {}_2F_1\left(\frac{3-\beta}{2}, 2-\frac{\beta}{2}; \frac{n}{2}+1; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right. \\
& \quad \left. - \frac{1-\beta}{n} \frac{|\mathbf{k}|^2}{\lambda^2} {}_2F_1\left(\frac{2-\beta}{2}, \frac{3-\beta}{2}; \frac{n}{2}+1; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right] \\
& = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \left[\lambda^\beta - \lambda^\beta {}_2F_1\left(-\frac{\beta}{2}, \frac{1-\beta}{2}; \frac{n}{2}; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right] \\
& = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \left[\lambda^\beta - \lambda^\beta \left(1 + \frac{|\mathbf{k}|^2}{\lambda^2}\right)^{\frac{\beta}{2}} {}_2F_1\left(-\frac{\beta}{2}, \frac{n+\beta-1}{2}; \frac{n}{2}; \frac{|\mathbf{k}|^2}{\lambda^2+|\mathbf{k}|^2}\right) \right] \\
& = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \left[\lambda^\beta - (\lambda^2 + |\mathbf{k}|^2)^{\frac{\beta}{2}} {}_2F_1\left(-\frac{\beta}{2}, \frac{n+\beta-1}{2}; \frac{n}{2}; \frac{|\mathbf{k}|^2}{\lambda^2+|\mathbf{k}|^2}\right) \right],
\end{aligned}$$

308 where ${}_2F_1$ is the Gaussian hypergeometric function and

$$309 \quad C = \left(\int_0^\pi \sin^{n-3}(\theta_2) d\theta_2 \right) \cdots \left(\int_0^\pi \sin(\theta_{n-2}) d\theta_{n-2} \right) \left(\int_0^{2\pi} d\theta_{n-1} \right) = \frac{2\pi^{\frac{n-1}{2}}}{\Gamma(\frac{n-1}{2})}.$$

310 So

$$311 \quad c_{n,\beta,\lambda} = \frac{\Gamma(\frac{n}{2})}{2\pi^{\frac{n}{2}} |\Gamma(-\beta)|}.$$

312 The PDEs for tempered Lévy flights or tempered Lévy flights combined with subdif-
313 fusion can be similarly derived, as those done in this section for Lévy flights or Lévy
314 flights combined with subdiffusion. Here, we present the counterpart of Eq. (28),

$$315 \quad \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1, \lambda}} + \frac{\partial^{\beta_2, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2, \lambda}} + \dots + \frac{\partial^{\beta_n, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n, \lambda}},$$

316 where the operator $\frac{\partial^{\beta_j, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_j, t)}{\partial |\mathbf{x}_j|^{\beta_j, \lambda}}$ is defined by taking $\beta = \beta_j$ and $n = 1$ in Eq.
317 (32). Again, even for the tempered Lévy flights, all the related operators should be
318 defined on the whole space, because of the very rare but still possible unbounded
319 jump lengths.

320 All the above derived PDEs are governing the PDF of the position of particles. If
321 one wants to dig out more deep informations of the corresponding stochastic processes,
322 analyzing the distribution of the functional defined by $A = \int_0^t U(\mathbf{X}(\tau)) d\tau$ is one of
323 the choices, where U is a prespecified function. Denote the PDF of the functional A

324 and position \mathbf{X} as $G(\mathbf{X}, A, t)$ and the counterpart of A in Fourier space as q . Then
 325 $\hat{G}(\mathbf{X}, q, t)$ solves [31]

$$326 \quad (36) \quad \frac{\partial \hat{G}(\mathbf{X}, q, t)}{\partial t} = K_{\alpha, \beta} \Delta^{\beta/2} D_t^{1-\alpha} \hat{G}(\mathbf{X}, q, t) + iqU(\mathbf{X}) \hat{G}(\mathbf{X}, q, t)$$

327 for Lévy flights combined with subdiffusion; and [32]

$$328 \quad (37) \quad \frac{\partial \hat{G}(\mathbf{X}, q, t)}{\partial t} = K_{\alpha, \beta} (\Delta + \lambda)^{\beta/2} D_t^{1-\alpha} \hat{G}(\mathbf{X}, q, t) + iqU(\mathbf{X}) \hat{G}(\mathbf{X}, q, t)$$

for tempered Lévy flights combined with subdiffusion, where

$$D_t^{1-\alpha} \hat{G}(\mathbf{X}, q, t) = \frac{1}{\Gamma(\alpha)} \left[\frac{\partial}{\partial t} - iqU(\mathbf{X}) \right] \int_0^t \frac{e^{i(t-\tau)qU(\mathbf{X})}}{(t-\tau)^{1-\alpha}} \hat{G}(\mathbf{X}, q, \tau) d\tau.$$

329 If one is only interested in the functional A (not caring position \mathbf{X}), then $\hat{G}_{\mathbf{X}_0}(q, t)$
 330 is, respectively, governed by [31]

$$331 \quad (38) \quad \frac{\partial \hat{G}_{\mathbf{X}_0}(q, t)}{\partial t} = K_{\alpha, \beta} D_t^{1-\alpha} \Delta^{\beta/2} \hat{G}_{\mathbf{X}_0}(q, t) + iqU(\mathbf{X}) \hat{G}_{\mathbf{X}_0}(q, t)$$

332 and [32]

$$333 \quad (39) \quad \frac{\partial \hat{G}_{\mathbf{X}_0}(q, t)}{\partial t} = K_{\alpha, \beta} D_t^{1-\alpha} (\Delta + \lambda)^{\beta/2} \hat{G}_{\mathbf{X}_0}(q, t) + iqU(\mathbf{X}) \hat{G}_{\mathbf{X}_0}(q, t)$$

334 for Lévy flights and tempered Lévy flights, combined with subdiffusion; the \mathbf{X}_0 in
 335 $\hat{G}_{\mathbf{X}_0}(q, t)$ means the initial position of particles, being a parameter.

336 **3. Specifying the generalized boundary conditions for the fractional**
 337 **PDEs.** After introducing the microscopic models and deriving the macroscopic ones,
 338 we have insight into anomalous diffusions, especially Lévy flights and tempered Lévy
 339 flights. In Section 2, all the derived equations are time dependent. From the process
 340 of derivation, one can see that the issue of initial condition can be easily/reasonably
 341 fixed, as classical ones, just specifying the value of $p(\mathbf{X}, 0)$ in the domain Ω . For Lévy
 342 processes, except Brownian motion, all others have discontinuous paths. As a result,
 343 the boundary $\partial\Omega$ itself (see Figure 3) can not be hit by the majority of discontinuous
 344 sample trajectories. This implies that when solving the PDEs derived in Section 2, the
 345 generalized boundary conditions must be introduced, i.e., the information of $p(\mathbf{X}, t)$
 346 on the domain $\mathbb{R}^n \setminus \Omega$ must be properly accounted for. In the following, we focus on
 347 Eqs. (20), (28), (31), (35) to discuss the boundary issues.

348 **3.1. Generalized Dirichlet type boundary conditions.** The appropriate
 349 initial and boundary value problems for Eq. (20) should be

$$350 \quad (40) \quad \begin{cases} \frac{\partial p(\mathbf{X}, t)}{\partial t} = \Delta^{\beta/2} p(\mathbf{X}, t) = \frac{-\beta \Gamma(\frac{n+\beta}{2})}{2^{1-\beta} \pi^{n/2} \Gamma(1-\beta/2)} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C} B_\varepsilon(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} & \text{in } \Omega, \\ p(\mathbf{X}, 0)|_\Omega = p_0(\mathbf{X}), \\ p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t). \end{cases}$$




Fig. 3: Domain of solving equations given in Section 2.

351 In Eq. (40), the term

$$\begin{aligned}
 & \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\varepsilon}(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\mathcal{C}_{B_\varepsilon}(\mathbf{X}) \cap \Omega)} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} + \int_{\mathbb{R}^n \setminus \Omega} \frac{p(\mathbf{X}, t) - g(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 352 \quad (41) \quad &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\mathcal{C}_{B_\varepsilon}(\mathbf{X}) \cap \Omega)} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} + p(\mathbf{X}, t) \int_{\mathbb{R}^n \setminus \Omega} |\mathbf{X} - \mathbf{Y}|^{-n-\beta} d\mathbf{Y} \\
 & \quad + \int_{\mathbb{R}^n \setminus \Omega} \frac{-g(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y}.
 \end{aligned}$$

353 According to Eq. (41), $g(\mathbf{X}, t)$ should satisfy that there exist positive M and C such
 354 that when $|\mathbf{X}| > M$,

$$355 \quad (42) \quad \frac{|g(\mathbf{X}, t)|}{|\mathbf{X}|^{\beta-\varepsilon}} < C \quad \text{for positive small } \varepsilon.$$

356 In particular, when Eq. (42) holds, the function $\int_{\mathbb{R}^n \setminus \Omega} \frac{-g(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y}$ of \mathbf{X} has any
 357 order of derivative if $g(\mathbf{X}, t)$ is integrable in any bounded domain. One of the most
 358 popular cases is $g(\mathbf{X}, t) \equiv 0$, which is the so-called absorbing boundary condition,
 359 implying that the particle is killed whenever it leaves the domain Ω . For example,
 360 consider the steady state fractional diffusion equation

$$361 \quad (43) \quad \begin{cases} \Delta^{\beta/2} p(\mathbf{X}) = -1 & \text{in } \Omega, \\ p(\mathbf{X})|_{\mathbb{R}^n \setminus \Omega} = 0. \end{cases}$$

362 The meaning of the solution $p(\mathbf{X})$ of (43) is the mean first exit time [6, 19] of particles
 363 performing Lévy flights [8]; if taking $\Omega = \{\mathbf{X} : |\mathbf{X}| < r\}$, then [16]

$$364 \quad (44) \quad p(\mathbf{X}) = \frac{\Gamma(n/2)(r^2 - |\mathbf{x}|^2)^{\beta/2}}{2^\beta \Gamma(1 + \beta/2) \Gamma(n/2 + \beta/2)}.$$

365 We generate the sample trajectories of Lévy flights to plot one and two dimensional
 366 mean first exit times, respectively, in Figures 4 and 5, which confirm the effectiveness
 367 of the model (43).




Fig. 4: Behaviors of $p(\mathbf{x})$ generated by at least 1.5×10^6 trajectories with $\beta = 0.5$, $n = 1$, $r = 100$. The (red) solid line is theoretical result (eq. (44)). The symbols are for the simulation result.




Fig. 5: Behaviors of $p(\mathbf{x})$ generated by at least 1.5×10^6 trajectories with $\beta = 0.5$, $n = 2$, $r = 100$. The (red) solid line is theoretical result (eq. (44)). The symbols are for the simulation result.

368 Another interesting case is for the steady state fractional diffusion equation

369 (45)
$$\begin{cases} \Delta^{\beta/2} p(\mathbf{X}) = 0 & \text{in } \Omega, \\ p(\mathbf{X})|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}). \end{cases}$$

370 Given a domain $H \subset \mathbb{R}^n \setminus \Omega$, if taking $g(\mathbf{X}) = 1$ for $\mathbf{X} \in H$ and 0 for $\mathbf{X} \in (\mathbb{R}^n \setminus \Omega) \setminus H$,
 371 then the solution of (45) means the probability that the particles undergoing Lévy
 372 flights lands in H after first escaping the domain Ω [8]. If $g(\mathbf{X}) \equiv 1$ in $\mathbb{R}^n \setminus \Omega$, then
 373 $p(\mathbf{X})$ equals to 1 in Ω because of the probability interpretation. This can also be
 374 analytically checked.

375 For the initial and boundary value problem Eq. (28), it should be written as

$$376 \quad (46) \quad \begin{cases} \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2}} \\ \quad \quad \quad + \dots + \frac{\partial^{\beta_n} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n}} \quad \text{in } \Omega, \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, 0)|_{\Omega} = p_0(\mathbf{x}_1, \dots, \mathbf{x}_n), \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{x}_1, \dots, \mathbf{x}_n, t). \end{cases}$$

377 Similar to (41), in (46) the term

$$378 \quad (47) \quad \begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\varepsilon}(\mathbf{x}_j)} \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t) - p(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\mathcal{C}_{B_\varepsilon}(\mathbf{x}_j) \cap \Omega)} \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t) - p(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j \\ & \quad + p(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t) \int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} |\mathbf{x}_j - \mathbf{y}_j|^{-1-\beta_j} d\mathbf{y}_j \\ & \quad + \int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} \frac{-g(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j. \end{aligned}$$

379 From Eq. (47), for $j = 1, \dots, n$, $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$ should satisfies that
380 there exist positive M and C such that when $|\mathbf{x}_j| > M$,

$$381 \quad (48) \quad \frac{|g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)|}{|\mathbf{x}_j|^{\beta_j - \varepsilon}} < C \quad \text{for positive small } \varepsilon.$$

382 The discussions below Eq. (45) still makes sense for Eq. (46). If $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$
383 satisfies Eq. (48), and it is integrable w.r.t. \mathbf{x}_j in any bounded interval. Then
384 $\int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} \frac{-g(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j$ has any order of partial derivative w.r.t. \mathbf{x}_j .

385 The initial and boundary value problem for Eq. (31) is

$$386 \quad (49) \quad \begin{cases} \frac{\partial p(\mathbf{X}, t)}{\partial t} = (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t) \quad \text{in } \Omega, \\ p(\mathbf{X}, 0)|_{\Omega} = p_0(\mathbf{X}), \\ p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t). \end{cases}$$

387 Like the discussions for Eq. (40), $g(\mathbf{X}, t)$ should satisfies that there exist positive M
388 and C such that when $|\mathbf{X}| > M$,

$$389 \quad (50) \quad \frac{|g(\mathbf{X}, t)|}{e^{(\lambda - \varepsilon)|\mathbf{X}|}} < C \quad \text{for positive small } \varepsilon.$$

390 If Eq. (50) holds and $g(\mathbf{X}, t)$ is integrable in any bounded domain, the function
391 $\int_{\mathbb{R}^n \setminus \Omega} \frac{-g(\mathbf{Y}, t)}{e^{\lambda|\mathbf{X} - \mathbf{Y}|} |\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y}$ of \mathbf{X} has any order of derivative.

392 Again, the corresponding tempered steady state fraction diffusion equation is

$$393 \quad (51) \quad \begin{cases} (\Delta + \lambda)^{\beta/2} p(\mathbf{X}) = 0 \quad \text{in } \Omega, \\ p(\mathbf{X})|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}). \end{cases}$$

394 For $H \subset \mathbb{R}^n \setminus \Omega$, if taking $g(\mathbf{X}) = 1$ for $\mathbf{X} \in H$ and 0 for $\mathbf{X} \in (\mathbb{R}^n \setminus \Omega) \setminus H$, then the
 395 solution of (51) means the probability that the particles undergoing tempered Lévy
 396 flights lands in H after first escaping the domain Ω . If $g(\mathbf{X}) \equiv 1$ in $\mathbb{R}^n \setminus \Omega$, then $p(\mathbf{X})$
 397 equals to 1 in Ω .

398 The initial and boundary value problem (35) should be written as

$$399 \quad (52) \quad \begin{cases} \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1, \lambda}} + \frac{\partial^{\beta_2, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2, \lambda}} \\ \quad \quad \quad + \dots + \frac{\partial^{\beta_n, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n, \lambda}} \quad \text{in } \Omega, \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, 0)|_{\Omega} = p_0(\mathbf{x}_1, \dots, \mathbf{x}_n), \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{x}_1, \dots, \mathbf{x}_n, t). \end{cases}$$

400 For $j = 1, \dots, n$, $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$ should satisfy that there exist positive
 401 M and C such that when $|\mathbf{x}_j| > M$,

$$402 \quad (53) \quad \frac{|g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)|}{e^{(\lambda - \varepsilon)|\mathbf{x}_j|}} < C \quad \text{for positive small } \varepsilon.$$

403 If $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$ is integrable w.r.t. \mathbf{x}_j in any bounded interval and satisfies
 404 Eq. (53), then $\int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} \frac{-g(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{e^{\lambda|\mathbf{x}_j - \mathbf{y}_j|} |\mathbf{x}_j - \mathbf{y}_j|^{1 + \beta_j}} d\mathbf{y}_j$ has any order of partial derivative
 405 w.r.t. \mathbf{x}_j .

406 The ways of specifying the initial and boundary conditions for Eqs. (36) and (38)
 407 are the same as Eq. (40). But for Eq. (36), the corresponding (42) should be changed
 408 as

$$409 \quad (54) \quad \frac{|U(\mathbf{X})g(\mathbf{X}, t)|}{|\mathbf{X}|^{\beta - \varepsilon}} < C \quad \text{for positive small } \varepsilon.$$

410 Similarly, the initial and boundary conditions of Eqs. (37) and (39) should be specified
 411 as the ones of Eq. (49). But for Eq. (37), the corresponding (50) needs to be changed
 412 as

$$413 \quad (55) \quad \frac{|U(\mathbf{X})g(\mathbf{X}, t)|}{e^{(\lambda - \varepsilon)|\mathbf{X}|}} < C \quad \text{for positive small } \varepsilon.$$

414 For the existence and uniqueness of the corresponding time-independent equations,
 415 one may refer to [14].

416 **3.2. Generalized Neumann type boundary conditions.** Because of the in-
 417herent discontinuity of the trajectories of Lévy flights or tempered Lévy flights, the
 418 traditional Neumann type boundary conditions can not be simply extended to the
 419 fractional PDEs. For the related discussions, see, e.g., [4, 10]. Based on the mod-
 420 els built in Sec. 2 and the law of mass conservation, we derive the reasonable ways
 421 of specifying the Neumann type boundary conditions, especially the reflecting ones.
 422 Let us first recall the derivation of classical diffusion equation. For normal diffusion
 423 (Brownian motion), microscopically the first moment of the distribution of waiting
 424 times and the second moment of the distribution of jump length are bounded, i.e., in
 425 Laplace and Fourier spaces, they are respectively like $1 - c_1 u$ and $1 - c_2 |\mathbf{k}|^2$; plugging
 426 them into Eq. (1) or Eq. (9) and performing integral transformations lead to the
 427 classical diffusion equation

$$428 \quad (56) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = (c_2/c_1) \Delta p(\mathbf{X}, t).$$




Fig. 6: Sketch map of particles jumping into, or jumping out of, or passing through the domain: Ω .

429 On the other hand, because of mass conservation, the continuity equation states that
 430 a change in density in any part of a system is due to inflow and outflow of particles
 431 into and out of that part of system, i.e., no particles are created or destroyed:

432 (57)
$$\frac{\partial p(\mathbf{X}, t)}{\partial t} = -\nabla \cdot \mathbf{j},$$

433 where \mathbf{j} is the flux of diffusing particles. Combining (56) with (57), one may take

434 (58)
$$\mathbf{j} = -(c_2/c_1)\nabla p(\mathbf{X}, t),$$

435 which is exactly Fick's law, a phenomenological postulation, saying that the flux goes
 436 from regions of high concentration to regions of low concentration with a magnitude
 437 proportional to the concentration gradient. In fact, for a long history, even up to
 438 now, most of the people are more familiar with the process: using the continuity
 439 equation (57) and Fick's law (58) derives the diffusion equation (56). The so-called
 440 reflecting boundary condition for (56) is to let the flux \mathbf{j} be zero along the boundary
 441 of considered domain.

442 Here we want to stress that Eq. (57) holds for any kind of diffusions, including
 443 the normal and anomalous ones. For Eqs. (40,46,49,52) governing the PDF of Lévy
 444 flights or tempered Lévy flights, using the continuity equation (57), one can get the
 445 corresponding fluxes and the counterparts of Fick's law; may we call it fractional
 446 Fick's law. Combining (40) with (57), one may let

447 (59)
$$\mathbf{j}_\Delta = \left\{ -\frac{1}{2n} c_{n,\beta} \int_{-\infty}^{\mathbf{x}_i} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{|\mathbf{Y}|^{n+\beta}} d\mathbf{Y} d\mathbf{x}_i \right\}_{n \times 1}$$

448 being the flux for the diffusion operator $\Delta^{\beta/2}$ with $\beta \in (0, 2)$, or calling it fractional
 449 Fick's law corresponding to $\Delta^{\beta/2}$. From (46) and (57), one may choose

(60)
 450
$$\mathbf{j}_{hv} = \left\{ -\frac{1}{2} c_{1,\beta_i} \int_{-\infty}^{\mathbf{x}_i} \int_{-\infty}^{+\infty} \frac{p(\mathbf{X} + \tilde{\mathbf{Y}}_i, t) + p(\mathbf{X} - \tilde{\mathbf{Y}}_i, t) - 2 \cdot p(\mathbf{X}, t)}{|\mathbf{y}_i|^{1+\beta_i}} d\mathbf{y}_i d\mathbf{x}_i \right\}_{n \times 1},$$

451 where $\tilde{\mathbf{Y}}_i = \{\mathbf{x}_1, \dots, \mathbf{y}_i, \dots, \mathbf{x}_n\}^T$, being the flux (fractional Fick's law) corresponding
 452 to the horizontal and vertical type fractional operators. Similarly, we can also get the

453 flux (fractional Fick's law) corresponding to the tempered fractional Laplacian and
 454 tempered horizontal and vertical type fractional operators, being respectively taken
 455 as

(61)

$$456 \quad \mathbf{j}_{\Delta,\lambda} = \left\{ -\frac{1}{2n} c_{n,\beta,\lambda} \int_{-\infty}^{\mathbf{x}_i} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y} d\mathbf{x}_i \right\}_{n \times 1}$$

457 and

(62)

$$458 \quad \mathbf{j}_{hv,\lambda} = \left\{ -\frac{1}{2} c_{1,\beta_i,\lambda} \int_{-\infty}^{\mathbf{x}_i} \int_{-\infty}^{+\infty} \frac{p(\mathbf{X} + \tilde{\mathbf{Y}}_i, t) + p(\mathbf{X} - \tilde{\mathbf{Y}}_i, t) - 2 \cdot p(\mathbf{X}, t)}{e^{\lambda|\mathbf{y}_i|} |\mathbf{y}_i|^{1+\beta_i}} d\mathbf{y}_i d\mathbf{x}_i \right\}_{n \times 1}$$

459 with $\tilde{\mathbf{Y}}_i = \{\mathbf{x}_1, \dots, \mathbf{y}_i, \dots, \mathbf{x}_n\}^T$.

460 Naturally, the Neumann type boundary conditions of (40,46,49,52) should be
 461 closely related to the values of the fluxes in the domain: $\mathbb{R}^n \setminus \Omega$; if the fluxes are
 462 zero in it, then one gets the so-called reflecting boundary conditions of the equations.
 463 Microscopically the motion of particles undergoing Lévy flights or tempered Lévy
 464 flights are much different from the Brownian motion; very rare but extremely long
 465 jumps dominate the dynamics, making the trajectories of the particles discontinuous.
 466 As shown in Figure 6, the particles may jump into, or jump out of, or even pass
 467 through the domain: Ω . But the number of particles inside Ω is conservative, which
 468 can be easily verified by making the integration of (57) in the domain Ω , i.e.,

$$469 \quad (63) \quad \frac{\partial}{\partial t} \int_{\Omega} p(\mathbf{X}, t) d\mathbf{X} = - \int_{\Omega} \nabla \cdot \mathbf{j} d\mathbf{X} = - \int_{\partial\Omega} \mathbf{j} \cdot \mathbf{n} ds = 0,$$

470 where \mathbf{n} is the outward-pointing unit normal vector on the boundary. If $\mathbf{j}|_{\mathbb{R}^n \setminus \Omega} = 0$,
 471 then for (40) $\Delta^{\frac{\beta}{2}} p(\mathbf{X}, t) = \nabla \cdot \mathbf{j} = 0$ in $\mathbb{R}^n \setminus \Omega$. So, the Neumann type boundary
 472 conditions for (40), (46), (49), and (52) can be, heuristically, defined as

$$473 \quad (64) \quad \Delta^{\frac{\beta}{2}} p(\mathbf{X}, t) = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

474 (65)

$$475 \quad \frac{\partial^{\beta_1} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2}} + \dots + \frac{\partial^{\beta_n} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n}} = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

476

$$477 \quad (66) \quad (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t) = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

478 and

(67)

$$479 \quad \frac{\partial^{\beta_1, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1, \lambda}} + \frac{\partial^{\beta_2, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2, \lambda}} + \dots + \frac{\partial^{\beta_n, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n, \lambda}} = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

480 respectively. The corresponding reflecting boundary conditions are with $g(\mathbf{X}) \equiv 0$.

481 **Remark:** 1. The Neumann type boundary conditions (64)-(67) derived in this
 482 section are independent of the choice of the flux \mathbf{j} , provided that it satisfies the con-
 483 dition (57). 2. Sending $\beta \rightarrow 2$, the usual Neumann boundary condition is recovered;
 484 for example, consider the model

$$485 \quad (68) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ \Delta^{\frac{\beta}{2}} p = g & \text{in } \mathbb{R}^n \setminus \Omega. \end{cases}$$

486

The weak solution $p : [0, +\infty) \rightarrow H^{\beta/2}(\mathbb{R}^n)$ of (68) satisfies

$$\int_{\Omega} p_t q d\mathbf{X} + \int_{\mathbb{R}^n} \Delta^{\beta/4} p \Delta^{\beta/4} q d\mathbf{X} = \int_{\Omega} f q d\mathbf{X} - \int_{\mathbb{R}^n \setminus \Omega} g q d\mathbf{X} \quad \forall q \in H^{\beta/2}(\mathbb{R}^n).$$

Taking $\beta \rightarrow 2$ leads to the usual Neumann boundary condition since

$$\int_{\mathbb{R}^n \setminus \Omega} g q d\mathbf{X} = \int_{\mathbb{R}^n \setminus \Omega} (\Delta p) q d\mathbf{X} = - \int_{\partial\Omega} \frac{\partial p}{\partial n} q ds - \int_{\mathbb{R}^n \setminus \Omega} \nabla p \cdot \nabla q d\mathbf{X}$$

and

$$\int_{\mathbb{R}^n} \Delta^{1/2} p \Delta^{1/2} q d\mathbf{X} = \int_{\mathbb{R}^n} \nabla p \cdot \nabla q d\mathbf{X}.$$

4. Well-posedness and regularity of the fractional PDEs with generalized BCs. Here, we show the well-posedness of the models discussed in the above sections, taking the models with the operator $\Delta^{\frac{\beta}{2}}$ as examples; the other ones can be similarly proved. For any real number $s \in \mathbb{R}$, we denote by $H^s(\mathbb{R}^n)$ the conventional Sobolev space of functions (see [1, 23]), equipped with the norm

$$\|u\|_{H^s(\mathbb{R}^n)} := \left(\int_{\mathbb{R}^n} (1 + |\mathbf{k}|^{2s}) |\hat{u}(\mathbf{k})|^2 d\mathbf{k} \right)^{\frac{1}{2}},$$

where \hat{u} denotes the Fourier transform of u . The notation $H^s(\Omega)$ denotes the space of functions on Ω that admit extensions to $H^s(\mathbb{R}^n)$, equipped with the quotient norm

$$\|u\|_{H^s(\Omega)} := \inf_{\tilde{u}} \|\tilde{u}\|_{H^s(\mathbb{R}^n)},$$

487 where the infimum extends over all possible $\tilde{u} \in H^s(\mathbb{R}^n)$ such that $\tilde{u} = u$ on Ω (in
488 the sense of distributions). The dual space of $H^s(\Omega)$ will be denoted by $H^s(\Omega)'$. The
489 following inequality will be used below:

$$490 \quad (69) \quad C^{-1} (\|\Delta^{\frac{\beta}{4}} u\|_{L^2(\mathbb{R}^n)} + \|u\|_{L^2(\Omega)}) \leq \|u\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \leq C (\|\Delta^{\frac{\beta}{4}} u\|_{L^2(\mathbb{R}^n)} + \|u\|_{L^2(\Omega)}).$$

492 Let $H_0^s(\Omega)$ be the subspace of $H^s(\mathbb{R}^n)$ consisting of functions which are zero in
493 $\mathbb{R}^n \setminus \Omega$. It is isomorphic to the completion of $C_0^\infty(\Omega)$ in $H^s(\Omega)$. The dual space of
494 $H_0^s(\Omega)$ will be denoted by $H^{-s}(\Omega)$.

495 For any Banach space B , the space $L^2(0, T; B)$ consists of functions $u : (0, T) \rightarrow B$
496 such that

$$497 \quad (70) \quad \|u\|_{L^2(0, T; B)} := \left(\int_0^T \|u(\cdot, t)\|_B^2 dt \right)^{\frac{1}{2}} < \infty,$$

499 and $H^1(0, T; B) = \{u \in L^2(0, T; B) : \partial_t u \in L^2(0, T; B)\}$; see [12].

500 **4.1. Dirichlet problem.** For any given $g \in \mathbb{R} \cup (L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \cap H^1(0, T;$
501 $H^{-\frac{\beta}{2}}(\mathbb{R}^n))) \hookrightarrow C([0, T]; L^2(\mathbb{R}^n))$, consider the time-dependent Dirichlet problem

$$502 \quad (71) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ p = g & \text{in } \mathbb{R}^n \setminus \Omega, \\ p(\cdot, 0) = p_0 & \text{in } \Omega. \end{cases}$$

503

504 The weak formulation of (71) is to find $p = g + \phi$ such that

$$505 \quad (72) \quad \phi \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)) \cap H^1(0, T; H^{-\frac{\beta}{2}}(\Omega)) \hookrightarrow C([0, T]; L^2(\Omega))$$

507 and

$$508 \quad (73) \quad \int_0^T \int_{\Omega} \partial_t \phi q \, d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} \phi \Delta^{\frac{\beta}{4}} q \, d\mathbf{X} dt = \int_0^T \int_{\Omega} (f + \Delta^{\frac{\beta}{2}} g - \partial_t g) q \, d\mathbf{X} dt$$

$$509 \quad \forall q \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)).$$

511 It is easy to see that $a(\phi, q) := \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} \phi \Delta^{\frac{\beta}{4}} q \, d\mathbf{X}$ is a coercive bilinear form
512 on $H_0^{\frac{\beta}{2}}(\Omega) \times H_0^{\frac{\beta}{2}}(\Omega)$ (cf. [34, section 30.2]) and $\ell(q) := \int_{\Omega} (f + \Delta^{\frac{\beta}{2}} g - \partial_t g) q \, d\mathbf{X}$ is a
513 continuous linear functional on $L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega))$. Such a problem as (73) has a unique
514 weak solution (cf. [34, Theorem 30.A]).

515 The weak solution actually depends only on the values of g in $\mathbb{R}^n \setminus \Omega$, independent
516 of the values of g in Ω . To see this, suppose that $g, \tilde{g} \in \mathbb{R} \cup (L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \cap$
517 $H^1(0, T; H^{-\frac{\beta}{2}}(\mathbb{R}^n))) \hookrightarrow C([0, T]; L^2(\mathbb{R}^n))$ are two functions such that $g = \tilde{g}$ in $\mathbb{R}^n \setminus \Omega$,
518 and p and \tilde{p} are the weak solutions of

$$519 \quad (74) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ p = g & \text{in } \mathbb{R}^n \setminus \Omega, \\ p(\cdot, 0) = p_0 & \text{in } \Omega, \end{cases} \quad \text{and} \quad \begin{cases} \frac{\partial \tilde{p}}{\partial t} - \Delta^{\frac{\beta}{2}} \tilde{p} = f & \text{in } \Omega, \\ \tilde{p} = \tilde{g} & \text{in } \mathbb{R}^n \setminus \Omega, \\ \tilde{p}(\cdot, 0) = p_0 & \text{in } \Omega, \end{cases}$$

521 respectively. Then the function $p - \tilde{p} \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)) \cap H^1(0, T; H^{-\frac{\beta}{2}}(\Omega))$ satisfies
522 (75)

$$522 \quad \int_0^T \int_{\Omega} \partial_t (p - \tilde{p}) q \, d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} (p - \tilde{p}) \Delta^{\frac{\beta}{4}} q \, d\mathbf{X} dt = 0 \quad \forall q \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)).$$

524 Substituting $q = p - \tilde{p}$ into the equation above immediately yields $p - \tilde{p} = 0$ a.e. in
525 $\mathbb{R}^n \times (0, T)$.

526 **4.2. Neumann problem.** Consider the Neumann problem

$$527 \quad (76) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ \Delta^{\frac{\beta}{2}} p = g & \text{in } \mathbb{R}^n \setminus \Omega, \\ p(\cdot, 0) = p_0 & \text{in } \Omega. \end{cases}$$

529 **DEFINITION 1 (Weak solutions).** *The weak formulation of (76) is to find $p \in$*
530 $L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \cap C([0, T]; L^2(\Omega))$ *such that*

$$531 \quad (77) \quad \partial_t p \in L^2(0, T; H^{\frac{\beta}{2}}(\Omega)') \text{ and } p(\cdot, 0) = p_0,$$

533 *satisfying the following equation:*

$$534 \quad (78) \quad \int_0^T \int_{\Omega} \partial_t p(\mathbf{X}, t) q(\mathbf{X}, t) \, d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p(\mathbf{X}, t) \Delta^{\frac{\beta}{4}} q(\mathbf{X}, t) \, d\mathbf{X} dt$$

$$= \int_0^T \int_{\Omega} f(\mathbf{X}, t) q(\mathbf{X}, t) \, d\mathbf{X} dt - \int_0^T \int_{\mathbb{R}^n \setminus \Omega} g(\mathbf{X}, t) q(\mathbf{X}, t) \, d\mathbf{X} dt$$

$$\forall q \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)).$$

535 THEOREM 2 (Existence and uniqueness of weak solutions). *If $p_0 \in L^2(\Omega)$, $f \in$
536 $L^2(0, T; H^{\frac{\beta}{2}}(\Omega)')$ and $g \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)')$, then there exists a unique weak so-
537 lution of (76) in the sense of Definition 1.*

538 *Proof* Let $t_k = k\tau$, $k = 0, 1, \dots, N$, be a partition of the time interval $[0, T]$, with
539 step size $\tau = T/N$, and define

$$540 \quad (79) \quad f_k(\mathbf{X}) := \frac{1}{\tau} \int_{t_{k-1}}^{t_k} f(\mathbf{X}, t) dt, \quad k = 0, 1, \dots, N,$$

$$541 \quad (80) \quad g_k(\mathbf{X}) := \frac{1}{\tau} \int_{t_{k-1}}^{t_k} g(\mathbf{X}, t) dt, \quad k = 0, 1, \dots, N.$$

543 Consider the time-discrete problem: for a given $p_{k-1} \in L^2(\mathbb{R}^n)$, find $p_k \in H^{\frac{\beta}{2}}(\mathbb{R}^n)$
544 such that the following equation holds:

$$545 \quad \frac{1}{\tau} \int_{\Omega} p_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} + \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p_k(\mathbf{X}) \Delta^{\frac{\beta}{4}} q(\mathbf{X}) d\mathbf{X}$$

$$546 \quad (81) \quad = \frac{1}{\tau} \int_{\Omega} p_{k-1}(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} + \int_{\Omega} f_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} - \int_{\mathbb{R}^n \setminus \Omega} g_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} \quad \forall q \in H^{\frac{\beta}{2}}(\mathbb{R}^n).$$

548 In view of (69), the left-hand side of the equation above is a coercive bilinear form
549 on $H^{\frac{\beta}{2}}(\mathbb{R}^n) \times H^{\frac{\beta}{2}}(\mathbb{R}^n)$, while the right-hand side is a continuous linear functional on
550 $H^{\frac{\beta}{2}}(\mathbb{R}^n)$. Consequently, the Lax–Milgram Lemma implies that there exists a unique
551 solution $p_k \in H^{\frac{\beta}{2}}(\mathbb{R}^n)$ for (81).

552 Substituting $q = p_k$ into (81) yields

$$553 \quad \frac{\|p_k\|_{L^2(\Omega)}^2 - \|p_{k-1}\|_{L^2(\Omega)}^2}{2\tau} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}^2$$

$$554 \quad \leq \|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} \|p_k\|_{H^{\frac{\beta}{2}}(\Omega)} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)}$$

$$555 \quad \leq (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}) \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)}$$

$$556 \quad (82) \quad \leq (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}) (\|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}^2 + \|p_k\|_{L^2(\Omega)}^2).$$

558 Then, summing up the inequality above for $k = 1, 2, \dots, N$, we have

$$559 \quad \max_{1 \leq k \leq N} \|p_k\|_{L^2(\Omega)}^2 + \tau \sum_{k=1}^N \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}^2$$

$$560 \quad (83) \quad \leq \|p_0\|_{L^2(\Omega)}^2 + C\tau \sum_{k=1}^N (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)' }^2 + \|p_k\|_{L^2(\Omega)}^2).$$

562 By applying Grönwall's inequality to the last estimate, there exists a positive constant
563 τ_0 such that when $\tau < \tau_0$ we have

$$564 \quad \max_{1 \leq k \leq N} \|p_k\|_{L^2(\Omega)}^2 + \tau \sum_{k=1}^N \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)}^2$$

$$565 \quad (84) \quad \leq C \|p_0\|_{L^2(\Omega)}^2 + C\tau \sum_{k=1}^N (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)' }^2).$$

567 Since any $q \in H^{\frac{\beta}{2}}(\Omega)$ can be extended to $q \in H^{\frac{\beta}{2}}(\mathbb{R}^n)$ with $\|q\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \leq 2\|q\|_{H^{\frac{\beta}{2}}(\Omega)}$,
 568 choosing such a q in (81) yields

$$\begin{aligned}
 569 \quad & \left| \int_{\Omega} \frac{p_k(\mathbf{X}) - p_{k-1}(\mathbf{X})}{\tau} q(\mathbf{X}) d\mathbf{X} \right| \\
 570 \quad & = \left| \int_{\Omega} f_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} - \int_{\mathbb{R}^n \setminus \Omega} g_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} - \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p_k(\mathbf{X}) \Delta^{\frac{\beta}{4}} q(\mathbf{X}) d\mathbf{X} \right| \\
 571 \quad & \leq C(\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}) \|q\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \\
 572 \quad & \leq C(\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}) \|q\|_{H^{\frac{\beta}{2}}(\Omega)}. \\
 573
 \end{aligned}$$

574 which implies (via duality)

$$\begin{aligned}
 575 \quad (85) \quad & \left\| \frac{p_k - p_{k-1}}{\tau} \right\|_{H^{\frac{\beta}{2}}(\Omega)'} \leq C(\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}). \\
 576
 \end{aligned}$$

577 The last inequality and (84) can be combined and written as

$$\begin{aligned}
 578 \quad & \max_{1 \leq k \leq N} \|p_k\|_{L^2(\Omega)}^2 + \tau \sum_{k=1}^N \left(\left\| \frac{p_k - p_{k-1}}{\tau} \right\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)}^2 \right) \\
 579 \quad (86) \quad & \leq C\|p_0\|_{L^2(\Omega)}^2 + C\tau \sum_{k=1}^N (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}^2). \\
 580
 \end{aligned}$$

581 If we define the piecewise constant functions

$$\begin{aligned}
 582 \quad (87) \quad & f^{(\tau)}(\mathbf{X}, t) := f_k(\mathbf{X}) = \frac{1}{\tau} \int_{t_{k-1}}^{t_k} f(\mathbf{X}, t) dt \quad \text{for } t \in (t_{k-1}, t_k], \quad k = 0, 1, \dots, N,
 \end{aligned}$$

$$\begin{aligned}
 583 \quad (88) \quad & g^{(\tau)}(\mathbf{X}, t) := g_k(\mathbf{X}) = \frac{1}{\tau} \int_{t_{k-1}}^{t_k} g(\mathbf{X}, t) dt \quad \text{for } t \in (t_{k-1}, t_k], \quad k = 0, 1, \dots, N,
 \end{aligned}$$

$$\begin{aligned}
 584 \quad (89) \quad & p_+^{(\tau)}(\mathbf{X}, t) := p_k(\mathbf{X}) \quad \text{for } t \in (t_{k-1}, t_k], \quad k = 0, 1, \dots, N,
 \end{aligned}$$

586 and the piecewise linear function

$$\begin{aligned}
 (90) \quad & \\
 587 \quad & p^{(\tau)}(\mathbf{X}, t) := \frac{t_k - t}{\tau} p_{k-1}(\mathbf{X}) + \frac{t - t_{k-1}}{\tau} p_k(\mathbf{X}) \quad \text{for } t \in [t_{k-1}, t_k], \quad k = 0, 1, \dots, N, \\
 588
 \end{aligned}$$

589 then (81) and (86) imply

$$\begin{aligned}
 & \int_0^T \int_{\Omega} \partial_t p^{(\tau)}(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p_+^{(\tau)}(\mathbf{X}, t) \Delta^{\frac{\beta}{4}} q(\mathbf{X}, t) d\mathbf{X} dt \\
 590 \quad & = \int_0^T \int_{\Omega} f^{(\tau)}(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt - \int_0^T \int_{\mathbb{R}^n \setminus \Omega} g^{(\tau)}(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt \\
 & \forall q \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)),
 \end{aligned}$$

591 and

$$\begin{aligned}
& \|p^{(\tau)}\|_{C([0,T];L^2(\Omega))} + \|\partial_t p^{(\tau)}\|_{L^2(0,T;H^{\frac{\beta}{2}}(\Omega)')} \\
& + \|p^{(\tau)}\|_{L^\infty(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n))} + \|p_+^{(\tau)}\|_{L^\infty(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n))} \\
592 & \leq C \left(\|f^{(\tau)}\|_{L^2(0,T;H^{\frac{\beta}{2}}(\Omega)')} + \|g^{(\tau)}\|_{L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)')} \right) \\
& \leq C \left(\|f\|_{L^2(0,T;H^{\frac{\beta}{2}}(\Omega)')} + \|g\|_{L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)')} \right),
\end{aligned}$$

593 respectively, where the constant C is independent of the step size τ . The last in-
594 equality implies that $p^{(\tau)}$ is bounded in $H^1(0, T; H^{\frac{\beta}{2}}(\Omega)') \cap L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \hookrightarrow$
595 $C([0, T]; L^2(\Omega))$. Consequently, there exists $p \in H^1(0, T; H^{\frac{\beta}{2}}(\Omega)') \cap L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \hookrightarrow$
596 $C([0, T]; L^2(\Omega))$ and a subsequence $\tau_j \rightarrow 0$ such that

597 (91) $p^{(\tau_j)}$ converges to p weakly in $L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n))$,

598 (92) $p_+^{(\tau_j)}$ converges to p weakly in $L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n))$,

599 (93) $\partial_t p^{(\tau_j)}$ converges to $\partial_t p$ weakly in $L^2(0, T; H^{\frac{\beta}{2}}(\Omega)')$,

600 (94) $p^{(\tau_j)}$ converges to p weakly in $C([0, T]; H^{\frac{\beta}{2}}(\Omega)')$ (see [20, Appendix C]).

602 By taking $\tau = \tau_j \rightarrow 0$ in (91), we obtain (78). This proves the existence of a weak
603 solution p satisfying (77).

604 If there are two weak solutions p and \tilde{p} , then their difference $\eta = p - \tilde{p}$ satisfies
605 the equation

$$(95) \quad \int_0^T \int_{\Omega} \partial_t(p - \tilde{p})q \, d\mathbf{X}dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}}(p - \tilde{p})\Delta^{\frac{\beta}{4}}q \, d\mathbf{X}dt = 0 \quad \forall q \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)).$$

608 Substituting $q = p - \tilde{p}$ into the equation yields

$$(96) \quad \|p(\cdot, t) - \tilde{p}(\cdot, t)\|_{L^2(\Omega)}^2 + \|\Delta^{\frac{\beta}{4}}(p - \tilde{p})\|_{L^2(0,T;L^2(\mathbb{R}^n))}^2 = \|p(\cdot, 0) - \tilde{p}(\cdot, 0)\|_{L^2(\Omega)}^2 = 0,$$

611 which implies $p = \tilde{p}$ a.e. in $\mathbb{R}^n \times (0, T)$. The uniqueness is proved.

612 **Remark:** From the analysis of this section we see that, although the initial data
613 $p_0(\mathbf{X})$ physically exists in the whole space \mathbb{R}^n , one only needs to know its values in Ω
614 to solve the PDEs (under both Dirichlet and Neumann boundary conditions).

615 **5. Conclusion.** In the past decades, fractional PDEs become popular as the
616 effective models of characterizing Lévy flights or tempered Lévy flights. This paper
617 is trying to answer the question: What are the physically meaningful and mathe-
618 matically reasonable boundary constraints for the models? We physically introduce
619 the process of the derivation of the fractional PDEs based on the microscopic mod-
620 els describing Lévy flights or tempered Lévy flights, and demonstrate that from a
621 physical point of view when solving the fractional PDEs in a bounded domain Ω , the
622 informations of the models in $\mathbb{R}^n \setminus \Omega$ should be involved. Inspired by the deriva-
623 tion process, we specify the Dirichlet type boundary constraint of the fractional

624 PDEs as $p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$ and Neumann type boundary constraints as, e.g.,
 625 $(\Delta^{\beta/2} p(\mathbf{X}, t))|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$ for the fractional Laplacian operator, which is the usual
 626 Neumann boundary condition when $\beta \rightarrow 2$.

627 The tempered fractional Laplacian operator $(\Delta + \lambda)^{\beta/2}$ is physically introduced
 628 and mathematically defined. For the four specific fractional PDEs given in this paper,
 629 we prove their well-posedness with the specified Dirichlet or Neumann type boundary
 630 constraints. In fact, it can be easily checked that these fractional PDEs are not
 631 well-posed if their boundary constraints are (locally) given in the traditional way;
 632 the potential reason is that locally dealing with the boundary contradicts with the
 633 principles that the Lévy or tempered Lévy flights follow.

634

REFERENCES

- 635 [1] R. ADAMS AND A. J. J. F. FOURNIER, *Sobolev Spaces*, Elsevier/Academic Press, Amsterdam,
 636 second ed., 2003.
- 637 [2] D. APPLEBAUM, *Lévy processes and stochastic calculus*, Cambridge University Press, Cam-
 638 bridge, second ed., 2009, <https://doi.org/10.1017/CBO9780511809781>.
- 639 [3] E. BARKAI, A. V. NAUMOV, Y. G. VAINER, M. BAUER, AND L. KADOR, *Lévy statistics for*
 640 *random single-molecule line shapes in a glass*, Phys. Rev. Lett., 91 (2003), p. 075502,
 641 <https://doi.org/10.1103/physrevlett.91.075502>.
- 642 [4] G. BARLES, C. GEORGELIN, AND E. R. JAKOBSEN, *On Neumann and oblique derivatives*
 643 *boundary conditions for nonlocal elliptic equations*, J. Differential Equations, 256 (2014),
 644 pp. 1368–1394, <https://doi.org/10.1016/j.jde.2013.11.001>.
- 645 [5] J.-P. BOUCHAUD AND A. GEORGES, *Anomalous diffusion in disordered media: Statistical*
 646 *mechanisms, models and physical applications*, Phys. Rep., 195 (1990), pp. 127–293,
 647 [https://doi.org/10.1016/0370-1573\(90\)90099-n](https://doi.org/10.1016/0370-1573(90)90099-n).
- 648 [6] A.V. CHECHKIN, R. METZLER, V.Y. GONCHAR, J. KLAFTER, AND L.V. TANATAROV, *First*
 649 *passage and arrival time densities for Lévy flights and the failure of the method of images*,
 650 J. Phys. A: Math. Gen., 36 (2003), pp. L537–L544, [https://stacks.iop.org/JPhysA/36/](https://stacks.iop.org/JPhysA/36/L537)
 651 [L537](https://stacks.iop.org/JPhysA/36/L537).
- 652 [7] Z.-Q. CHEN AND R. SONG, *Two-sided eigenvalue estimates for subordinate processes in do-*
 653 *main*, J. Funct. Anal., 226 (2005), pp. 90–113, <https://doi.org/10.1016/j.jfa.2005.05.004>.
- 654 [8] W. H. DENG, X. C. WU, AND W. L. WANG, *Mean exit time and escape probability for the*
 655 *anomalous processes with the tempered power-law waiting times*, EPL, 117 (2017), p. 10009,
 656 <https://doi.org/10.1209/0295-5075/117/10009>.
- 657 [9] E. DI NEZZA, G. PALATUCCI, AND E. VALDINOCI, *Hitchhiker’s guide to the fractional Sobolev*
 658 *spaces*, Bull. Sci. math., 136 (2012), pp. 521–573, [https://doi.org/10.1016/j.bulsci.2011.12.](https://doi.org/10.1016/j.bulsci.2011.12.004)
 659 [004](https://doi.org/10.1016/j.bulsci.2011.12.004).
- 660 [10] S. DIPIERRO, X. ROS-OTON, AND E. VALDINOCI, *Nonlocal problems with Neumann boundary*
 661 *conditions*, Rev. Mat. Iberoam., (2017).
- 662 [11] Q. DU, M. GUNZBURGER, R. B. LEHOUCQ, AND K. ZHOU, *Analysis and approximation of*
 663 *nonlocal diffusion problems with volume constraints*, SIAM Rev., 54 (2012), pp. 667–696,
 664 <https://doi.org/10.1137/110833294>.
- 665 [12] L. C. EVANS, *Partial differential equations*, American Mathematical Society, Providence, RI,
 666 second ed., 2010.
- 667 [13] W. FELLER, *An introduction to probability theory and its applications*, John Wiley & Sons,
 668 Inc., New York, 1971. Vol. 2, Chap. XVI. 8, p. 525.
- 669 [14] M. FELSINGER, M. KASSMANN, AND P. VOIGT, *The Dirichlet problem for nonlocal operators*,
 670 Math. Z., 279 (2015), pp. 779–809, <https://doi.org/10.1007/s00209-014-1394-3>.
- 671 [15] H. C. FOGEDBY, *Langevin equations for continuous time Lévy flights*, Phys. Rev. E, 50 (1994),
 672 pp. 1657–1660, <https://doi.org/10.1103/physreve.50.1657>.
- 673 [16] R. K. GETTOOR, *First passage times for symmetric stable processes in space*, Trans. Amer.
 674 Math. Soc., 101 (1961), pp. 75–90, <https://doi.org/10.1090/S0002-9947-1961-0137148-5>.
- 675 [17] Q.-Y. GUAN AND Z.-M. MA, *Boundary problems for fractional Laplacians*, Stoch. Dyn., 5
 676 (2005), pp. 385–424, <https://doi.org/10.1142/s021949370500150x>.
- 677 [18] I. KOPONEN, *Analytic approach to the problem of convergence of truncated Lévy flights towards*
 678 *the Gaussian stochastic process*, Phys. Rev. E, 52 (1995), pp. 1197–1199, [https://doi.org/](https://doi.org/10.1103/physreve.52.1197)
 679 [10.1103/physreve.52.1197](https://doi.org/10.1103/physreve.52.1197).
- 680 [19] T. KOREN, M.A. LOMHOLT, A.V. CHECHKIN, J. KLAFTER, AND R. METZLER, *Leapover lengths*

- 681 and first passage time statistics for Lévy flights, Phys. Rev. Lett., 99 (2007), p. 160602,
682 <https://doi.org/10.1103/PhysRevLett.99.160602>.
- 683 [20] P.-L. LIONS, *Mathematical Topics in Fluid Mechanics: Volume 1: Incompressible Models*,
684 Clarendon Press, Oxford, USA., 1996.
- 685 [21] B. B. MANDELBROT AND J. W. V. NESS, *Fractional Brownian motions, fractional noises and*
686 *applications*, SIAM Rev., 10 (1968), pp. 422–437, <https://doi.org/10.1137/1010093>.
- 687 [22] R. N. MANTEGNA AND H. E. STANLEY, *Stochastic process with ultraslow convergence to a*
688 *Gaussian: The truncated Lévy flight*, Phys. Rev. Lett., 73 (1994), pp. 2946–2949, <https://doi.org/10.1103/physrevlett.73.2946>.
- 689 [23] W. MCLEAN, *Strongly elliptic systems and boundary integral equations*, Cambridge university
690 press, 2000.
- 691 [24] M. M. MEERSCHAERT AND A. SIKORSKII, *Stochastic Models for Fractional Calculus*, Walter de
692 Gruyter, Berlin, 2012.
- 693 [25] R. METZLER AND J. KLAFTER, *The random walk's guide to anomalous diffusion: a frac-*
694 *tional dynamics approach*, Phys. Rep., 339 (2000), pp. 1–77, [https://doi.org/10.1016/](https://doi.org/10.1016/S0370-1573(00)00070-3)
695 [S0370-1573\(00\)00070-3](https://doi.org/10.1016/S0370-1573(00)00070-3).
- 696 [26] E. W. MONTROLL AND G. H. WEISS, *Random walks on lattices. II*, J. Math. Phys., 6 (1965),
697 pp. 167–181, <https://doi.org/10.1063/1.1704269>.
- 698 [27] J. M. SANCHO, A. M. LACASTA, K. LINDENBERG, I. M. SOKOLOV, AND A. H. ROMERO, *Diffusion*
699 *on a solid surface: Anomalous is normal*, Phys. Rev. Lett., 92 (2004), p. 250601, <https://doi.org/10.1103/physrevlett.92.250601>.
- 700 [28] M. F. SHLESINGER, J. KLAFTER, AND B. J. WEST, *Levy walks with applications to turbu-*
701 *lence and chaos*, Phys. A, 140 (1986), pp. 212–218, [https://doi.org/10.1016/0378-4371\(86\)](https://doi.org/10.1016/0378-4371(86)90224-4)
702 [90224-4](https://doi.org/10.1016/0378-4371(86)90224-4).
- 703 [29] L. SILVESTRE, *Regularity of the obstacle problem for a fractional power of the Laplace operator*,
704 Commun. Pure Appl. Math., 60 (2007), pp. 67–112, <https://doi.org/10.1002/cpa.20153>.
- 705 [30] I. M. SOKOLOV, J. MAI, AND A. BLUMEN, *Paradoxal diffusion in chemical space for nearest-*
706 *neighbor walks over polymer chains*, Phys. Rev. Lett., 79 (1997), pp. 857–860, <https://doi.org/10.1103/physrevlett.79.857>.
- 707 [31] L. TURGEMAN, S. CARMÍ, AND E. BARKAI, *Fractional Feynman-Kac equation for non-Brownian*
708 *functionals*, Phys. Rev. Lett., 103 (2009), p. 190201, [https://doi.org/10.1103/physrevlett.](https://doi.org/10.1103/physrevlett.103.190201)
709 [103.190201](https://doi.org/10.1103/physrevlett.103.190201).
- 710 [32] X. C. WU, W. H. DENG, AND E. BARKAI, *Tempered fractional Feynman-Kac equation: Theory*
711 *and examples*, Phys. Rev. E, 93 (2016), p. 032151, [https://doi.org/10.1103/physreve.93.](https://doi.org/10.1103/physreve.93.032151)
712 [032151](https://doi.org/10.1103/physreve.93.032151).
- 713 [33] V. ZABURDAEV, S. DENISOV, AND J. KLAFTER, *Lévy walks*, Rev. Mod. Phys., 87 (2015), pp. 483–
714 530, <https://doi.org/10.1103/revmodphys.87.483>.
- 715 [34] E. ZEIDLER, *Nonlinear Functional Analysis and its Applications II/B: Nonlinear Mono-*
716 *tone Operators*, Springer Science+Business Media, LLC, 1990, [https://doi.org/10.1007/](https://doi.org/10.1007/978-1-4612-0981-2)
717 [978-1-4612-0981-2](https://doi.org/10.1007/978-1-4612-0981-2). Translated by the author and by Leo F. Boron.
- 718
- 719
- 720