

BOUNDARY PROBLEMS FOR THE FRACTIONAL AND TEMPERED FRACTIONAL OPERATORS*

WEIHUA DENG[†], BUYANG LI[‡], WENYI TIAN[§], AND PINGWEN ZHANG[¶]

Abstract. For characterizing the Brownian motion in a bounded domain: Ω , it is well-known that the boundary conditions of the classical diffusion equation just rely on the given information of the solution along the boundary of a domain; on the contrary, for the Lévy flights or tempered Lévy flights in a bounded domain, it involves the information of a solution in the complementary set of Ω , i.e., $\mathbb{R}^n \setminus \Omega$, with the potential reason that paths of the corresponding stochastic process are discontinuous. Guided by probability intuitions and the stochastic perspectives of anomalous diffusion, we show the reasonable ways, ensuring the clear physical meaning and well-posedness of the partial differential equations (PDEs), of specifying ‘boundary’ conditions for space fractional PDEs modeling the anomalous diffusion. Some properties of the operators are discussed, and the well-posednesses of the PDEs with generalized boundary conditions are proved.

Key words. Lévy flight; Tempered Lévy flight; Well-posedness; Generalized boundary conditions

1. Introduction. The phrase ‘anomalous is normal’ says that anomalous diffusion phenomena are ubiquitous in the natural world. It was first used in the title of [27], which reveals that the diffusion of classical particles on a solid surface has rich anomalous behaviour controlled by the friction coefficient. In fact, anomalous diffusion is no longer a young topic. In the review paper [5], the evolution of particles in disordered environments was investigated; the specific effects of a bias on anomalous diffusion were considered; and the generalizations of Einstein’s relation in the presence of disorder were discussed. With the rapid development of the study of anomalous dynamics in diverse field, some deterministic equations are derived, governing the macroscopic behaviour of anomalous diffusion. In 2000, Metzler and Klafter published the survey paper [25] for the equations governing transport dynamics in complex system with anomalous diffusion and non-exponential relaxation patterns, i.e., fractional kinetic equations of the diffusion, advection-diffusion, and Fokker-Planck type, derived asymptotically from basic random walk models and a generalized master equation. Many mathematicians have been involved in the research of fractional partial differential equations (PDEs). For fractional PDEs in a bounded domain Ω , an important question is how to introduce physically meaningful and mathematically well-posed boundary conditions on $\partial\Omega$ or $\mathbb{R}^n \setminus \Omega$.

Microscopically, diffusion is the net movement of particles from a region of higher concentration to a region of lower concentration; for the normal diffusion (Brownian motion), the second moment of the particle trajectories is a linear function of the time t ; naturally, if it is a nonlinear function of t , we call the corresponding diffu-

*This work was partially supported by NSFC under Grant Nos. 11421101, 1142110001, 11626250, 11671182, and the Fundamental Research Funds for the Central Universities under Grant No. lzujbky-2017-ot10. WD thanks Mark M. Meerschaert and Zhen-Qing Chen for the discussions. We thank the anonymous reviewer for Remark 2 of Sec. 2.3.

[†]School of Mathematics and Statistics, Gansu Key Laboratory of Applied Mathematics and Complex Systems, Lanzhou University, Lanzhou 730000, P.R. China. Email: dengwh@lzu.edu.cn

[‡]Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong. Email: buyang.li@polyu.edu.hk

[§]Center for Applied Mathematics, Tianjin University, Tianjin 300072, P.R. China. Email: wenyi.tian@tju.edu.cn

[¶]School of Mathematical Sciences, Laboratory of Mathematics and Applied Mathematics, Peking University, Beijing 100871, P.R. China. Email: pzhang@pku.edu.cn

sion process anomalous diffusion or non-Brownian diffusion [25]. The microscopic (stochastic) models describing anomalous diffusion include continuous time random walks (CTRWs), Langevin type equation, Lévy processes, subordinated Lévy processes, and fractional Brownian motion, etc.. The CTRWs contain two important random variables describing the motion of particles [26], i.e., the waiting time ξ and jump length η . If both the first moment of ξ and the second moment of η are finite in the scaling limit, then the CTRWs approximate Brownian motion. On the contrary, if one of them is divergent, then the CTRWs characterize anomalous diffusion. Two of the most important CTRW models are Lévy flights and Lévy walks. For Lévy flights, the ξ with finite first moment and η with infinite second moment are independent, leading to infinite propagation speed and the divergent second moments of the distribution of the particles. This causes much difficulty in relating the models to experimental data, especially when analyzing the scaling of the measured moments in time [33]. With coupled distribution of ξ and η (the infinite speed is penalized by the corresponding waiting times), we get the so-called Lévy walks [33]. Another idea to ensure that the processes have bounded moments is to truncate the long tailed probability distribution of Lévy flights [22]; they still look like a Lévy flight in not too long a time. Currently, the most popular way to do the truncation is to use the exponential tempering, offering the technical advantage of still being an infinitely divisible Lévy process after the operation [24]. The Lévy process to describe anomalous diffusion is the scaling limit of CTRWs with independent ξ and η . It is characterized by its characteristic function. Except Brownian motion with drift, the paths of all other proper Lévy processes are discontinuous. Sometimes, the Lévy flights are conveniently described by the Brownian motion subordinated to a Lévy process [7]. Fractional Brownian motions are often taken as the models to characterize subdiffusion [21].

Macroscopically, fractional (nonlocal) PDEs are the most popular and effective models for anomalous diffusion, derived from the microscopic models. The solution of fractional PDEs is generally the probability density function (PDF) of the position of the particles undergoing anomalous dynamics; with the deepening of research, the fractional PDEs governing the functional distribution of particles' trajectories are also developed [31, 32]. Two ways are usually used to derive the fractional PDEs. One is based on the Montroll-Weiss equation [26], i.e., in Fourier-Laplace space, the PDF $p(\mathbf{X}, t)$ obeys

$$(1) \quad \hat{p}(\mathbf{k}, u) = \frac{1 - \phi(u)}{u} \cdot \frac{\hat{p}_0(\mathbf{k})}{1 - \Psi(u, \mathbf{k})},$$

where $\hat{p}_0(\mathbf{k})$ is the Fourier transform of the initial data; $\phi(u)$ is the Laplace transform of the PDF of waiting times ξ and $\Psi(u, \mathbf{k})$ the Laplace and the Fourier transforms of the joint PDF of waiting times ξ and jump length η . If ξ and η are independent, then $\Psi(u, \mathbf{k}) = \phi(u)\psi(\mathbf{k})$, where $\psi(\mathbf{k})$ is the Fourier transform of the PDF of η . Another way is based on the characteristic function of the α -stable Lévy motion, being the scaling limit of the CTRW model with power law distribution of jump length η . In the high dimensional case, it is more convenient to make the derivation by using the characteristic function of the stochastic process. According to the Lévy-Khinchin formula [2], the characteristic function of Lévy process has a specific form

$$(2) \quad \int_{\mathbb{R}^n} e^{i\mathbf{k} \cdot \mathbf{X}} p(\mathbf{X}, t) d\mathbf{X} = \mathbf{E}(e^{i\mathbf{k} \cdot \mathbf{X}}) = e^{t\Phi(\mathbf{k})},$$

where

$$\Phi(\mathbf{k}) = i\mathbf{a} \cdot \mathbf{k} - \frac{1}{2}(\mathbf{k} \cdot \mathbf{b}\mathbf{k}) + \int_{\mathbb{R}^n \setminus \{0\}} [e^{i\mathbf{k} \cdot \mathbf{X}} - 1 - i(\mathbf{k} \cdot \mathbf{X})\chi_{\{|\mathbf{X}| < 1\}}] \nu(d\mathbf{X});$$

here χ_I is the indicator function of the set I , $\mathbf{a} \in \mathbb{R}^n$, $\mathbf{b}$ is a positive definite symmetric $n \times n$ matrix and ν is a sigma-finite Lévy measure on $\mathbb{R}^n \setminus \{0\}$. When $\mathbf{a}$ and $\mathbf{b}$ are zero and

$$(3) \quad \nu(d\mathbf{X}) = \frac{\beta\Gamma(\frac{n+\beta}{2})}{2^{1-\beta}\pi^{n/2}\Gamma(1-\beta/2)} |\mathbf{X}|^{-\beta-n} d\mathbf{X},$$

the process is a rotationally symmetric β -stable Lévy motion and its PDF solves

$$(4) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = \Delta^{\beta/2} p(\mathbf{X}, t),$$

where $\mathcal{F}(\Delta^{\beta/2} p(\mathbf{X}, t)) = -|\mathbf{k}|^\beta \mathcal{F}(p(\mathbf{X}, t))$ [29]. If replacing (3) by the measure of isotropic tempered power law with the tempering exponent λ , then we get the corresponding PDF evolution equation

$$(5) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t),$$

where $(\Delta + \lambda)^{\beta/2}$ is defined by (32) in physical space and by (34) in Fourier space.

In practice, the choice of $\nu(d\mathbf{X})$ depends strongly on the concrete physical environment. For example, Figure 1 clearly shows the horizontal and vertical structure. So, we need to take the measure as (if it is superdiffusion)

$$(6) \quad \begin{aligned} \nu(d\mathbf{X}) = \nu(d\mathbf{x}_1 d\mathbf{x}_2) &= \frac{\beta_1 \Gamma(\frac{1+\beta_1}{2})}{2^{1-\beta_1} \pi^{1/2} \Gamma(1-\beta_1/2)} |\mathbf{x}_1|^{-\beta_1-1} \delta(\mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2 \\ &+ \frac{\beta_2 \Gamma(\frac{1+\beta_2}{2})}{2^{1-\beta_2} \pi^{1/2} \Gamma(1-\beta_2/2)} \delta(\mathbf{x}_1) |\mathbf{x}_2|^{-\beta_2-1} d\mathbf{x}_1 d\mathbf{x}_2, \end{aligned}$$

where β_1 and β_2 belong to $(0, 2)$. If $\mathbf{a}$ and $\mathbf{b}$ equal to zero, then it leads to diffusion equation

$$(7) \quad \frac{\partial p(\mathbf{x}_1, \mathbf{x}_2, t)}{\partial t} = \frac{\partial^{\beta_1} p(\mathbf{x}_1, \mathbf{x}_2, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \mathbf{x}_2, t)}{\partial |\mathbf{x}_2|^{\beta_2}}.$$

Under the guidelines of probability intuitions and stochastic perspectives [17] of Lévy flights or tempered Lévy flights, we discuss the reasonable ways of defining fractional partial differential operators and specifying the ‘boundary’ conditions for their macroscopic descriptions, i.e., the PDEs of the types Eqs. (4), (5), (7), and their extensions, e.g., the fractional Feynman-Kac equations [31, 32]. For the related discussions on the nonlocal diffusion problems from a mathematical point of view, one can see the review paper [11]. The divergence of the second moment and the discontinuity of the paths of Lévy flights predicate that the corresponding diffusion operators should be defined on $\mathbb{R}^n$, which further signify that if we are solving the equations in a bounded domain Ω , the information in $\mathbb{R}^n \setminus \Omega$ should also be involved. We will show that the generalized Dirichlet type boundary conditions should be specified as $p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$. If the particles are killed after leaving the domain Ω , then



Fig. 1: Sketch map for the physical environment suitable for Eq. (7).

$g(\mathbf{X}, t) \equiv 0$, i.e., the so-called absorbing boundary conditions. Because of the discontinuity of the jumps of Lévy flights, a particular concept ‘escape probability’ can be introduced, which means the probability that the particle jumps from the domain Ω into a domain $H \subset \mathbb{R}^n \setminus \Omega$; for solving the escape probability, one just needs to specify $g(\mathbf{X}) = 1$ for $\mathbf{X} \in H$ and 0 for $\mathbf{X} \in (\mathbb{R}^n \setminus \Omega) \setminus H$ for the corresponding time-independent PDEs. As for the generalized Neumann type boundary conditions, our ideas come from the fact that the continuity equation (conservation law) holds for any kinds of diffusion, since the particles can not be created or destroyed. Based on the continuity equation and the governing equation of the PDF of Lévy or tempered Lévy flights, the corresponding flux $\mathbf{j}$ can be obtained. So the generalized reflecting boundary conditions should be $\mathbf{j}|_{\mathbb{R}^n \setminus \Omega} \equiv 0$, which implies $(\nabla \cdot \mathbf{j})|_{\mathbb{R}^n \setminus \Omega} \equiv 0$. Then, the generalized Neumann type boundary conditions are given as $(\nabla \cdot \mathbf{j})|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$, e.g., for (4), it should be taken as $(\Delta^{\beta/2} p(\mathbf{X}, t))|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$. The well-posednesses of the equations under our specified generalized Dirichlet or Neumann type boundary conditions are well established.

Overall, this paper focuses on introducing physically reasonable boundary constraints for a large class of fractional PDEs, building a bridge between the physical and mathematical communities for studying anomalous diffusion and fractional PDEs. In the next section, we recall the derivation of fractional PDEs. Some new concepts are introduced, such as the tempered fractional Laplacian, and some properties of anomalous diffusion are found. In Sec. 3, we discuss the reasonable ways of specifying the generalized boundary conditions for the fractional PDEs governing the position or functional distributions of Lévy flights and tempered Lévy flights. In Sec. 4, we prove well-posedness of the fractional PDEs under the generalized Dirichlet and Neumann boundary conditions defined on the complement of the bounded domain. Conclusion and remarks are given in the last section.

2. Preliminaries. For well understanding and inspiring the ways of specifying the ‘boundary constraints’ to PDEs governing the PDF of Lévy flights or tempered Lévy flights, we will show the ideas of deriving the microscopic and macroscopic models.

2.1. Microscopic models for anomalous diffusion. For the microscopic description of the anomalous diffusion, we consider the trajectory of a particle or a stochastic process, i.e., $\mathbf{X}(t)$. If $\langle |\mathbf{X}(t)|^2 \rangle \sim t$, the process is normal, otherwise it is abnormal. The anomalous diffusions of most often happening in natural world are the cases that $\langle |\mathbf{X}(t)|^2 \rangle \sim t^\gamma$ with $\gamma \in [0, 1) \cup (1, 2]$. A Lévy flight is a random walk in which the jump length has a heavy tailed (power law) probability distribution,




Fig. 2: Random trajectories (1000 steps) of Lévy flight ($\beta = 0.8$), tempered Lévy flight ($\beta = 0.8$, $\lambda = 0.2$), and Brownian motion.

i.e., the PDF of jump length r is like $r^{-\beta-n}$ with $\beta \in (0, 2)$, and the distribution in direction is uniform. With the wide applications of Lévy flights in characterizing long-range interactions [3] or a nontrivial “crumpled” topology of a phase (or configuration) space of polymer systems [30], etc, its second and higher moments are divergent, leading to the difficulty in relating models to experimental data. In fact, for Lévy flights $\langle |\mathbf{X}(t)|^\delta \rangle \sim t^{\delta/\beta}$ with $0 < \delta < \beta \leq 2$. Under the framework of CTRW, the model Lévy walk [28] can circumvent this obstacle by putting a larger time cost to a longer displacement, i.e., using the space-time coupled jump length and waiting time distribution $\Psi(r, t) = \frac{1}{2}\delta(r - vt)\phi(t)$. Another popular model is the so-called tempered Lévy flights [18], in which the extremely long jumps is exponentially cut by using the distribution of jump length $e^{-r\lambda}r^{-\beta-n}$ with λ being a small modulation parameter (a smooth exponential regression towards zero). In not too long a time, the tempered Lévy flights display the dynamical behaviors of Lévy flights, ultraslowly converging to the normal diffusion. Figure 2 shows the trajectories of 1000 steps of Lévy flights, tempered Lévy flights, and Brownian motion in two dimensions; note the presence of rare but large jumps compared to the Brownian motion, playing the dominant role in the dynamics.

Using Berry-Esséen theorem [13], first established in 1941, which applies to the convergence to a Gaussian for a symmetric random walk whose jump probabilities have a finite third moment, we have that for the one dimensional tempered Lévy flights with the distribution of jump length $Ce^{-r\lambda}r^{-\beta-1}$ the convergence speed is

$$\frac{5}{2\sqrt{2C}} \frac{\Gamma(3-\beta)}{\Gamma(2-\beta)^{3/2}} \lambda^{-\frac{1}{2}\beta} \frac{1}{\sqrt{m}},$$

which means that the scaling law for the number of steps needed for Gaussian behavior to emerge as

$$(8) \quad m \sim \lambda^{-\beta}.$$

More concretely, letting $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_m$ be i.i.d. random variables with PDF $Ce^{-r\lambda}r^{-\beta-1}$ and $E(|\mathbf{X}_1|^2) = \sigma^2 > 0$, then the cumulative distribution function (CDF) Q_m of $\mathbf{Y}_m = (\mathbf{X}_1 + \mathbf{X}_2 + \dots + \mathbf{X}_m)/(\sigma\sqrt{m})$ converges to the CDF $Q(\mathbf{X})$ of the standard normal distribution as

$$|Q_m(\mathbf{X}) - Q(\mathbf{X})| < \frac{5}{2} \frac{\langle |\mathbf{X}|^3 \rangle}{\langle |\mathbf{X}|^2 \rangle^{3/2}} \frac{1}{\sqrt{m}} = \frac{5}{2\sqrt{2C}} \frac{\Gamma(3-\beta)}{\Gamma(2-\beta)^{3/2}} \lambda^{-\frac{1}{2}\beta} \frac{1}{\sqrt{m}},$$

174 since

$$175 \quad \langle |\mathbf{X}|^3 \rangle = C \int_{-\infty}^{\infty} |\mathbf{X}|^3 e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{-\beta-1} d|\mathbf{X}| = 2C \int_0^{\infty} e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{3-\beta-1} d|\mathbf{X}| = 2C\lambda^{\beta-3}\Gamma(3-\beta)$$

176 and

$$177 \quad \langle |\mathbf{X}|^2 \rangle = C \int_{-\infty}^{\infty} |\mathbf{X}|^2 e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{-\beta-1} d|\mathbf{X}| = 2C \int_0^{\infty} e^{-\lambda|\mathbf{X}|} |\mathbf{X}|^{2-\beta-1} d|\mathbf{X}| = 2C\lambda^{\beta-2}\Gamma(2-\beta).$$

178 From Eq. (8), it can be seen that with the decrease of λ , the required m for the
179 crossover between Lévy flight behavior and Gaussian behavior increase rapidly. A
180 little bit counterintuitive observation is that the number of variables required to the
181 crossover increases with the increase of β .

182 We have described the distributions of jump length for Lévy flights and tempered
183 Lévy flights, in which Poisson process is taken as the renewal process. We denote the
184 Poisson process with rate $\zeta > 0$ as $N(t)$ and its waiting time distribution between two
185 events is $\zeta e^{-\zeta t}$. Then the Lévy flights or tempered Lévy flights are the compound
186 Poisson process defined as $\mathbf{X}(t) = \sum_{j=0}^{N(t)} \mathbf{X}_j$, where $\mathbf{X}_j$ are i.i.d. random variables with
187 the distribution of power law or tempered power law. The characteristic function of
188 $\mathbf{X}(t)$ can be calculated as follows. For real $\mathbf{k}$, we have

$$\begin{aligned} \hat{p}(\mathbf{k}, t) &= \mathbf{E}(e^{i\mathbf{k} \cdot \mathbf{X}(t)}) \\ &= \sum_{j=0}^{\infty} \mathbf{E}(e^{i\mathbf{k} \cdot \mathbf{X}(t)} | N(t) = j) P(N(t) = j) \\ 189 \quad (9) \quad &= \sum_{j=0}^{\infty} \mathbf{E}(e^{i\mathbf{k} \cdot (\mathbf{X}_0 + \mathbf{X}_1 + \dots + \mathbf{X}_j)} | N(t) = j) P(N(t) = j) \\ &= \sum_{j=0}^{\infty} \Phi_0(\mathbf{k})^j \frac{(\zeta t)^j}{j!} e^{-\zeta t} \\ &= e^{\zeta t(\Phi_0(\mathbf{k}) - 1)}, \end{aligned}$$

190 where $\Phi_0(\mathbf{k}) = \mathbf{E}(e^{i\mathbf{k} \cdot \mathbf{X}_0})$, being also the characteristic function of $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_j$
191 since they are i.i.d.

192 In the CTRW model describing one dimensional Lévy flights or tempered Lévy
193 flights, the PDF of waiting times is taken as $\zeta e^{-\zeta t}$ with its Laplace transform $\zeta/(u+\zeta)$
194 and the PDF of jumping length is $c^{-\beta} r^{-\beta-1}$ or $e^{-\lambda r} r^{-\beta-1}$ with its Fourier transform
195 $1 - c^\beta |k|^\beta$ or $1 - c_{\beta,\lambda}[(\lambda + ik)^\beta - \lambda^\beta] - c_{\beta,\lambda}[(\lambda - ik)^\beta - \lambda^\beta]$. Substituting them into
196 the Montroll-Weiss Eq. (1) with $\hat{p}_0(k) = 1$ (the initial position of particles is at zero),
197 we get that $\hat{p}(k, u)$ of Lévy flights solves

$$198 \quad (10) \quad \hat{p}(k, u) = \frac{1}{u + \zeta c^\beta |k|^\beta};$$

199 and the $\hat{p}(k, u)$ of tempered Lévy flights obeys

$$200 \quad (11) \quad \hat{p}(k, u) = \frac{1}{u + \zeta C_{\beta,\lambda}[(\lambda + ik)^\beta - \lambda^\beta] + \zeta C_{\beta,\lambda}[(\lambda - ik)^\beta - \lambda^\beta]}.$$

201 If the subdiffusion is involved, we need to choose the PDF of waiting times as
 202 $\tilde{c}^{1+\alpha}t^{-\alpha-1}$ with $\alpha \in (0, 1)$ and its Laplace transform $1 - \tilde{c}^\alpha u^\alpha$. Then from (1),
 203 we get that

$$204 \quad (12) \quad \hat{p}(k, u) = \frac{\tilde{c}^\alpha}{u^{1-\alpha}(1 - (1 - \tilde{c}^\alpha u^\alpha)\psi(k))}.$$

205 For high dimensional case, the Lévy flights can also be characterized by Brow-
 206 nian motion subordinated to a Lévy process. Let $\mathbf{Y}(t)$ be a Brownian motion with
 207 Fourier exponent $-|\mathbf{k}|^2$ and $S(t)$ a subordinator with Laplace exponent $u^{\beta/2}$ that is
 208 independent of $\mathbf{Y}(t)$. The process $\mathbf{X}(t) = \mathbf{Y}(S(t))$ is describing Lévy flights with
 209 Fourier exponent $-|\mathbf{k}|^\beta$, being the subordinate process of $\mathbf{Y}(t)$. In effect, denote the
 210 characteristic function of $\mathbf{Y}(t)$ as $\Phi_y(\mathbf{k})$ and the one of $S(t)$ as $\Phi_s(u)$. Then the
 211 characteristic function of $\mathbf{X}(t)$ is as follows:

$$\begin{aligned} \hat{p}_x(\mathbf{k}, t) &= \int_{\mathbb{R}^n} e^{i\mathbf{k} \cdot \mathbf{X}} p_x(\mathbf{X}, t) d\mathbf{X} \\ &= \int_0^\infty \int_{\mathbb{R}^n} e^{i\mathbf{k} \cdot \mathbf{Y}} p_y(\mathbf{Y}, \tau) d\mathbf{Y} p_s(\tau, t) d\tau \\ &= \int_0^\infty e^{-\tau(-\Phi_y(\mathbf{k}))} p_s(\tau, t) d\tau \\ &= e^{-t\Phi_s(-\Phi_y(\mathbf{k}))}, \end{aligned} \quad 212 \quad (13)$$

213 where p_x , p_y , and p_s , are respectively the PDFs of the stochastic processes $\mathbf{X}$, $\mathbf{Y}$, and
 214 S . Similarly, in the following, we denote p with subscript (lowercase letter) as the
 215 PDF of the corresponding stochastic process (uppercase letter).

216 This paper mainly focuses on Lévy flights and tempered Lévy flights. If one is
 217 interested in subdiffusion, instead of Poisson process, the fractional Poisson process
 218 should be taken as the renewal process, in which the time interval between each pair
 219 of events follows the power law distribution. Let $\mathbf{Y}(t)$ be a general Lévy process
 220 with Fourier exponent $\Phi_y(\mathbf{k})$ and $S(t)$ a strictly increasing subordinator with Laplace
 221 exponent u^α ($\alpha \in (0, 1)$). Define the inverse subordinator $E(t) = \inf\{\tau > 0 : S(\tau) >$
 222 $t\}$. Since $t = S(\tau)$ and $\tau = E(t)$ are inverse processes, we have $P(E(t) \leq \tau) =$
 223 $P(S(\tau) \geq t)$. Hence

$$224 \quad (14) \quad p_e(\tau, t) = \frac{\partial P(E(t) \leq \tau)}{\partial \tau} = \frac{\partial}{\partial \tau} [1 - P(S(\tau) < t)] = -\frac{\partial}{\partial \tau} \int_0^t p_s(y, \tau) dy.$$

225 In the above equation, taking Laplace transform w.r.t t leads to

$$226 \quad (15) \quad p_e(\tau, u) = -\frac{\partial}{\partial \tau} u^{-1} e^{-\tau u^\alpha} = u^{\alpha-1} e^{-\tau u^\alpha}.$$

227 For the PDF $p_x(\mathbf{X}, t)$ of $\mathbf{X}(t) = \mathbf{Y}(E(t))$, there holds

$$228 \quad (16) \quad p_x(\mathbf{X}, t) = \int_0^\infty p_y(\mathbf{X}, \tau) p_e(\tau, t) d\tau.$$

229 Performing Fourier transform w.r.t. $\mathbf{X}$ and Laplace transform w.r.t. t to the above

equation results in

$$\begin{aligned}
\hat{p}_x(\mathbf{k}, u) &= \int_0^\infty \hat{p}_y(\mathbf{k}, \tau) p_e(\tau, u) d\tau \\
&= \int_0^\infty e^{-\tau \Phi_y(\mathbf{k})} u^{\alpha-1} e^{-\tau u^\alpha} d\tau \\
&= \frac{u^{\alpha-1}}{u^\alpha + \Phi_y(\mathbf{k})}.
\end{aligned}
\tag{17}$$

Remark. According to Fogedby [15], the stochastic trajectories of (scale limited) CTRW $\mathbf{X}(E_t)$ can also be expressed in terms of the coupled Langevin equation

$$\begin{cases} \dot{\mathbf{X}}(\tau) = F(\mathbf{X}(\tau)) + \eta(\tau), \\ \dot{S}(\tau) = \xi(\tau), \end{cases}
\tag{18}$$

where $F(\mathbf{X})$ is a vector field; E_t is the inverse process of $S(t)$; the noises $\eta(\tau)$ and $\xi(\tau)$ are statistically independent, corresponding to the distributions of jump length and waiting times.

2.2. Derivation of the macroscopic description from the microscopic models. This section focuses on the derivation of the deterministic equations governing the PDF of position of the particles undergoing anomalous diffusion. It shows that the operators related to (tempered) power law jump lengths should be defined on the whole unbounded domain $\mathbb{R}^n$, which can also be inspired by the rare but extremely long jump lengths displayed in Figure 2; the fact that among all proper Lévy processes Brownian motion is the unique one with continuous paths further consolidates the reasonable way of defining the operators. We derive the PDEs based on Eqs. (9), (13), and (16), since they apply for both one and higher dimensional cases. For one dimensional case, sometimes it is convenient to use (10), (11), and (12).

When the diffusion process is rotationally symmetric β -stable, i.e., it is isotropic with PDF of jump length $c_{\beta,n} r^{-\beta-n}$ and its Fourier transform $1 - |\mathbf{k}|^\beta$, where n is the space dimension. In Eq. (9), taking ζ equal to 1, we get the Cauchy equation

$$\frac{d\hat{p}(\mathbf{k}, t)}{dt} = -|\mathbf{k}|^\beta \hat{p}(\mathbf{k}, t).
\tag{19}$$

Performing inverse Fourier transform to the above equation leads to

$$\frac{\partial p(\mathbf{X}, t)}{\partial t} = \Delta^{\beta/2} p(\mathbf{X}, t),
\tag{20}$$

where

$$\begin{aligned}
\Delta^{\beta/2} p(\mathbf{X}, t) &= -c_{n,\beta} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C} B_\varepsilon(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
&= \frac{1}{2} c_{n,\beta} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{|\mathbf{Y}|^{n+\beta}} d\mathbf{Y}
\end{aligned}
\tag{21}$$

with [9]

$$c_{n,\beta} = \frac{\beta \Gamma(\frac{n+\beta}{2})}{2^{1-\beta} \pi^{n/2} \Gamma(1 - \beta/2)}.
\tag{22}$$

258 For the more general cases of Eq. (9), there is the Cauchy equation

$$259 \quad (23) \quad \frac{d\hat{p}(\mathbf{k}, t)}{dt} = (\Phi_0(\mathbf{k}) - 1)\hat{p}(\mathbf{k}, t),$$

260 so the PDF of the stochastic process $\mathbf{X}$ solves (taking $\zeta = 1$)

$$261 \quad (24) \quad \begin{aligned} \frac{\partial p(\mathbf{X}, t)}{\partial t} &= \mathcal{F}^{-1}\{(\Phi_0(\mathbf{k}) - 1)\hat{p}(\mathbf{k}, t)\} \\ &= \int_{\mathbb{R}^n \setminus \{0\}} [p(\mathbf{X} + \mathbf{Y}, t) - p(\mathbf{X}, t)] \nu(d\mathbf{Y}), \end{aligned}$$

where $\nu(d\mathbf{Y})$ is the probability measure of the jump length. Sometimes, to overcome the possible divergence of the terms on the right hand side of Eq. (24) because of the possible strong singularity of $\nu(d\mathbf{Y})$ at zero, the term

$$\Phi_0(\mathbf{k}) - 1 = \int_{\mathbb{R}^n \setminus \{0\}} [e^{i\mathbf{k} \cdot \mathbf{Y}} - 1] \nu(d\mathbf{Y})$$

262 is approximately replaced by

$$263 \quad (25) \quad \int_{\mathbb{R}^n \setminus \{0\}} [e^{i\mathbf{k} \cdot \mathbf{Y}} - 1 - i(\mathbf{k} \cdot \mathbf{Y})_{\chi_{\{|\mathbf{Y}| < 1\}}}] \nu(d\mathbf{Y});$$

264 then the corresponding modification to Eq. (24) is

$$265 \quad (26) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = \int_{\mathbb{R}^n \setminus \{0\}} \left[p(\mathbf{X} + \mathbf{Y}, t) - p(\mathbf{X}, t) - \sum_{i=1}^n \mathbf{y}_i (\partial_i p(\mathbf{X}, t))_{\chi_{\{|\mathbf{Y}| < 1\}}} \right] \nu(d\mathbf{Y}),$$

266 where $\mathbf{y}_i$ is the component of $\mathbf{Y}$, i.e., $\mathbf{Y} = \{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n\}^T$. If $\nu(-d\mathbf{Y}) = \nu(d\mathbf{Y})$,
267 the integration of the summation term of above equation equals to zero.

268 If the diffusion is in the environment having a structure like Figure 1, the proba-
269 bility measure should be taken as

$$\begin{aligned} (27) \quad \nu(d\mathbf{X}) &= \nu(d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \cdots d\mathbf{x}_n) \\ &= \frac{\beta_1 \Gamma(\frac{1+\beta_1}{2})}{2^{1-\beta_1} \pi^{1/2} \Gamma(1 - \beta_1/2)} |\mathbf{x}_1|^{-\beta_1-1} \delta(\mathbf{x}_2) \delta(\mathbf{x}_3) \cdots \delta(\mathbf{x}_n) d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \cdots d\mathbf{x}_n \\ 270 \quad &+ \frac{\beta_2 \Gamma(\frac{1+\beta_2}{2})}{2^{1-\beta_2} \pi^{1/2} \Gamma(1 - \beta_2/2)} |\mathbf{x}_2|^{-\beta_2-1} \delta(\mathbf{x}_1) \delta(\mathbf{x}_3) \cdots \delta(\mathbf{x}_n) d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \cdots d\mathbf{x}_n + \cdots \\ &+ \frac{\beta_n \Gamma(\frac{1+\beta_n}{2})}{2^{1-\beta_n} \pi^{1/2} \Gamma(1 - \beta_n/2)} |\mathbf{x}_n|^{-\beta_n-1} \delta(\mathbf{x}_1) \delta(\mathbf{x}_2) \cdots \delta(\mathbf{x}_{n-1}) d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}_3 \cdots d\mathbf{x}_n, \end{aligned}$$

271 where $\beta_1, \beta_2, \dots, \beta_n$ belong to $(0, 2)$. Plugging Eq. (27) into Eq. (24) leads to

$$272 \quad (28) \quad \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2}} + \cdots + \frac{\partial^{\beta_n} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n}},$$

where

$$\mathcal{F}_j \left(\frac{\partial^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|^{\beta_j}} \right) = -|\mathbf{k}_j|^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{k}_j, \mathbf{x}_{j+1}, \dots, \mathbf{x}_n, t)$$

273 and $\frac{\partial^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|^{\beta_j}}$ in physical space is defined by (21) with $n = 1$; in particular, when
 274 $\beta_j \in (1, 2)$, it can also be written as
 (29)
 275 $\frac{\partial^{\beta_j} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|^{\beta_j}} = -\frac{1}{2 \cos(\beta_j \pi / 2) \Gamma(2 - \beta_j)} \frac{\partial^2}{\partial \mathbf{x}_j^2} \int_{-\infty}^{\infty} |\mathbf{x}_j - \mathbf{y}|^{1-\beta_j} p(\mathbf{x}_1, \dots, \mathbf{y}, \dots, \mathbf{x}_n, t) d\mathbf{y}.$

It should be emphasized here that when characterizing diffusion processes related with Lévy flights the operators should be defined in the whole space. Another issue that also should be stressed is that when $\beta_1 = \beta_2 = \dots = \beta_n = 1$, Eq. (28) is still describing the phenomena of anomalous diffusion, including the cases that they belong to $(0, 1)$; the corresponding ‘first’ order operator is nonlocal, being different from the classical first order operator, but they have the same energy in the sense that

$$\begin{aligned} & \mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|} \right) \overline{\mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_j|} \right)} \\ &= (k_j)^2 \hat{p}^2(\mathbf{x}_1, \dots, \mathbf{x}_{j-1}, \mathbf{k}_j, \mathbf{x}_{j+1}, \dots, \mathbf{x}_n, t) \\ &= \mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial \mathbf{x}_j} \right) \overline{\mathcal{F} \left(\frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial \mathbf{x}_j} \right)}; \\ & \mathcal{F} \left(\Delta^{1/2} p(\mathbf{X}, t) \right) \overline{\mathcal{F} \left(\Delta^{1/2} p(\mathbf{X}, t) \right)} \\ &= |\mathbf{k}|^2 \hat{p}^2(\mathbf{k}, t) = \mathcal{F}(\nabla p(\mathbf{X}, t)) \cdot \overline{\mathcal{F}(\nabla p(\mathbf{X}, t))}, \end{aligned}$$

276 even though $\Delta^{1/2}$ and ∇ are completely different operators, where the notation $\bar{v}$
 277 stands for the complex conjugate of v .

278 If the subdiffusion is involved, the derivation of the macroscopic equation should
 279 be based on Eq. (17). For getting the term related to time derivative, the inverse
 280 Laplace transform should be performed on $u^\alpha \hat{p}(\mathbf{k}, u) - u^{\alpha-1}$. Since $\hat{p}(\mathbf{k}, t = 0)$ is taken
 281 as 1, there exists

282 (30)
$$\mathcal{L}^{-1}(u^\alpha \hat{p}(\mathbf{k}, u) - u^{\alpha-1}) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-\tau)^{-\alpha} \frac{\partial \hat{p}(\mathbf{k}, \tau)}{\partial \tau} d\tau,$$

283 which is usually denoted as ${}_0^C D_t^\alpha \hat{p}(\mathbf{k}, t)$, the so-called Caputo fractional derivative. So,
 284 if both the PDFs of the waiting time and jump lengths of the stochastic process $\mathbf{X}$ are
 285 power law, the corresponding models can be obtained by replacing $\frac{\partial}{\partial t}$ with ${}_0^C D_t^\alpha$ in
 286 Eqs. (20), (24), (26), and (28). Furthermore, if there is an external force $F(\mathbf{X})$ in the
 287 considered stochastic process $\mathbf{X}$, we need to add an additional term $\nabla \cdot (F(\mathbf{X})p(\mathbf{X}, t))$
 288 on the right hand side of Eqs. (20), (24), (26), and (28).

289 Here we turn to another important and interesting topic: tempered Lévy flights.
 290 Practically it is not easy to collect the value of a function in the unbounded area
 291 $\mathbb{R}^n \setminus \Omega$. This is one of the achievements of using tempered fractional Laplacian. It is
 292 still isotropic but with PDF of jump length $c_{\beta, n, \lambda} e^{-\lambda r} r^{-\beta-n}$. The PDF of tempered
 293 Lévy flights solves

294 (31)
$$\frac{\partial p(\mathbf{X}, t)}{\partial t} = (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t),$$

295 where

$$\begin{aligned}
 (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t) &= -c_{n,\beta,\lambda} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\varepsilon}(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{e^{\lambda|\mathbf{X}-\mathbf{Y}|} |\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 (32) \quad &= \frac{1}{2} c_{n,\beta,\lambda} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y}
 \end{aligned}$$

297 with

$$(33) \quad c_{n,\beta,\lambda} = \frac{\Gamma(\frac{n}{2})}{2\pi^{\frac{n}{2}} |\Gamma(-\beta)|}.$$

299 The choice of the constant as the one given in (33) leads to

$$(34) \quad \mathcal{F}\left((\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t)\right) = (-1)^{\lfloor \beta \rfloor} (\lambda^\beta - (\lambda^2 + |\mathbf{k}|^2)^{\frac{\beta}{2}} + O(|\mathbf{k}|^2)) \hat{p}(\mathbf{k}, t),$$

301 where $\beta \in (0, 1) \cup (1, 2)$, and $\lfloor \beta \rfloor$ means the biggest integer, being smaller than or
 302 equal to β .

303 However, if $\lambda = 0$, one needs to choose the constant as the one given in (22) to
 304 make sure $\mathcal{F}\left(\Delta^{\beta/2} p(\mathbf{X}, t)\right) = -|\mathbf{k}|^\beta \hat{p}(\mathbf{k}, t)$. The reason is as follows.

$$\begin{aligned}
 \mathcal{F}\left((\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t)\right) &= \frac{1}{2} c_{n,\beta,\lambda} \int_{\mathbb{R}^n} \frac{e^{i\mathbf{k} \cdot \mathbf{Y}} + e^{-i\mathbf{k} \cdot \mathbf{Y}} - 2}{|\mathbf{Y}|^{n+\beta}} e^{-\lambda|\mathbf{Y}|} d\mathbf{Y} \cdot \mathcal{F}(p(\mathbf{X}, t)) \\
 &= -c_{n,\beta,\lambda} \int_{\mathbb{R}^n} \frac{1 - \cos(\mathbf{k} \cdot \mathbf{Y})}{|\mathbf{Y}|^{n+\beta}} e^{-\lambda|\mathbf{Y}|} d\mathbf{Y} \cdot \mathcal{F}(p(\mathbf{X}, t)).
 \end{aligned}$$

306 For $\beta \in (0, 1) \cup (1, 2)$, then we have

$$\begin{aligned}
& \int_{\mathbb{R}^n} \frac{1 - \cos(\mathbf{k} \cdot \mathbf{Y})}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y} = \int_{\mathbb{R}^n} \frac{1 - \cos(|\mathbf{k}|\mathbf{y}_1)}{e^{\lambda|\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y} = |\mathbf{k}|^\beta \int_{\mathbb{R}^n} \frac{1 - \cos(\mathbf{x}_1)}{|\mathbf{X}|^{n+\beta}} e^{-\frac{\lambda}{|\mathbf{k}|}|\mathbf{x}|} d\mathbf{X} \\
& = C|\mathbf{k}|^\beta \int_0^\infty \frac{1}{r^{n+\beta}} e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{n-1} \left(\int_0^\pi (1 - \cos(r \cos \theta_1)) \sin^{n-2}(\theta_1) d\theta_1 \right) dr \\
& = \frac{1}{(-\beta)(-\beta+1)} C|\mathbf{k}|^{\beta-2} \lambda^2 \int_0^\infty e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{-\beta+1} \left(\int_0^\pi (1 - \cos(r \cos \theta_1)) \sin^{n-2}(\theta_1) d\theta_1 \right) dr \\
& \quad - \frac{1}{(-\beta)(-\beta+1)} C|\mathbf{k}|^{\beta-1} \lambda \int_0^\infty e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{-\beta+1} \left(\int_0^\pi \sin(r \cos \theta_1) \sin^{n-2}(\theta_1) \cos(\theta_1) d\theta_1 \right) dr \\
& \quad - \frac{1}{-\beta} C|\mathbf{k}|^\beta \int_0^\infty e^{-\frac{\lambda}{|\mathbf{k}|}r} r^{-\beta} \left(\int_0^\pi \sin(r \cos \theta_1) \sin^{n-2}(\theta_1) \cos(\theta_1) d\theta_1 \right) dr \\
307 & = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \lambda^\beta \left[1 - {}_2F_1\left(\frac{2-\beta}{2}, \frac{3-\beta}{2}; \frac{n}{2}; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right. \\
& \quad \left. - \frac{2-\beta}{n} \frac{|\mathbf{k}|^2}{\lambda^2} {}_2F_1\left(\frac{3-\beta}{2}, 2-\frac{\beta}{2}; \frac{n}{2}+1; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right. \\
& \quad \left. - \frac{1-\beta}{n} \frac{|\mathbf{k}|^2}{\lambda^2} {}_2F_1\left(\frac{2-\beta}{2}, \frac{3-\beta}{2}; \frac{n}{2}+1; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right] \\
& = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \left[\lambda^\beta - \lambda^\beta {}_2F_1\left(-\frac{\beta}{2}, \frac{1-\beta}{2}; \frac{n}{2}; -\frac{|\mathbf{k}|^2}{\lambda^2}\right) \right] \\
& = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \left[\lambda^\beta - \lambda^\beta \left(1 + \frac{|\mathbf{k}|^2}{\lambda^2}\right)^{\frac{\beta}{2}} {}_2F_1\left(-\frac{\beta}{2}, \frac{n+\beta-1}{2}; \frac{n}{2}; \frac{|\mathbf{k}|^2}{\lambda^2+|\mathbf{k}|^2}\right) \right] \\
& = C|\Gamma(-\beta)| \frac{\sqrt{\pi}\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})} \left[\lambda^\beta - (\lambda^2 + |\mathbf{k}|^2)^{\frac{\beta}{2}} {}_2F_1\left(-\frac{\beta}{2}, \frac{n+\beta-1}{2}; \frac{n}{2}; \frac{|\mathbf{k}|^2}{\lambda^2+|\mathbf{k}|^2}\right) \right],
\end{aligned}$$

308 where ${}_2F_1$ is the Gaussian hypergeometric function and

$$309 \quad C = \left(\int_0^\pi \sin^{n-3}(\theta_2) d\theta_2 \right) \cdots \left(\int_0^\pi \sin(\theta_{n-2}) d\theta_{n-2} \right) \left(\int_0^{2\pi} d\theta_{n-1} \right) = \frac{2\pi^{\frac{n-1}{2}}}{\Gamma(\frac{n-1}{2})}.$$

310 So

$$311 \quad c_{n,\beta,\lambda} = \frac{\Gamma(\frac{n}{2})}{2\pi^{\frac{n}{2}} |\Gamma(-\beta)|}.$$

312 The PDEs for tempered Lévy flights or tempered Lévy flights combined with subdif-
313 fusion can be similarly derived, as those done in this section for Lévy flights or Lévy
314 flights combined with subdiffusion. Here, we present the counterpart of Eq. (28),
(35)

$$315 \quad \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1, \lambda}} + \frac{\partial^{\beta_2, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2, \lambda}} + \cdots + \frac{\partial^{\beta_n, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n, \lambda}},$$

316 where the operator $\frac{\partial^{\beta_j, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_j, t)}{\partial |\mathbf{x}_j|^{\beta_j, \lambda}}$ is defined by taking $\beta = \beta_j$ and $n = 1$ in Eq.
317 (32). Again, even for the tempered Lévy flights, all the related operators should be
318 defined on the whole space, because of the very rare but still possible unbounded
319 jump lengths.

320 All the above derived PDEs are governing the PDF of the position of particles. If
321 one wants to dig out more deep informations of the corresponding stochastic processes,
322 analyzing the distribution of the functional defined by $A = \int_0^t U(\mathbf{X}(\tau)) d\tau$ is one of
323 the choices, where U is a prespecified function. Denote the PDF of the functional A

and position $\mathbf{X}$ as $G(\mathbf{X}, A, t)$ and the counterpart of A in Fourier space as q . Then $\hat{G}(\mathbf{X}, q, t)$ solves [31]

$$(36) \quad \frac{\partial \hat{G}(\mathbf{X}, q, t)}{\partial t} = K_{\alpha, \beta} \Delta^{\beta/2} D_t^{1-\alpha} \hat{G}(\mathbf{X}, q, t) + iqU(\mathbf{X}) \hat{G}(\mathbf{X}, q, t)$$

for Lévy flights combined with subdiffusion; and [32]

$$(37) \quad \frac{\partial \hat{G}(\mathbf{X}, q, t)}{\partial t} = K_{\alpha, \beta} (\Delta + \lambda)^{\beta/2} D_t^{1-\alpha} \hat{G}(\mathbf{X}, q, t) + iqU(\mathbf{X}) \hat{G}(\mathbf{X}, q, t)$$

for tempered Lévy flights combined with subdiffusion, where

$$D_t^{1-\alpha} \hat{G}(\mathbf{X}, q, t) = \frac{1}{\Gamma(\alpha)} \left[\frac{\partial}{\partial t} - iqU(\mathbf{X}) \right] \int_0^t \frac{e^{i(t-\tau)qU(\mathbf{X})}}{(t-\tau)^{1-\alpha}} \hat{G}(\mathbf{X}, q, \tau) d\tau.$$

If one is only interested in the functional A (not caring position $\mathbf{X}$), then $\hat{G}_{\mathbf{X}_0}(q, t)$ is, respectively, governed by [31]

$$(38) \quad \frac{\partial \hat{G}_{\mathbf{X}_0}(q, t)}{\partial t} = K_{\alpha, \beta} D_t^{1-\alpha} \Delta^{\beta/2} \hat{G}_{\mathbf{X}_0}(q, t) + iqU(\mathbf{X}) \hat{G}_{\mathbf{X}_0}(q, t)$$

and [32]

$$(39) \quad \frac{\partial \hat{G}_{\mathbf{X}_0}(q, t)}{\partial t} = K_{\alpha, \beta} D_t^{1-\alpha} (\Delta + \lambda)^{\beta/2} \hat{G}_{\mathbf{X}_0}(q, t) + iqU(\mathbf{X}) \hat{G}_{\mathbf{X}_0}(q, t)$$

for Lévy flights and tempered Lévy flights, combined with subdiffusion; the $\mathbf{X}_0$ in $\hat{G}_{\mathbf{X}_0}(q, t)$ means the initial position of particles, being a parameter.

3. Specifying the generalized boundary conditions for the fractional PDEs. After introducing the microscopic models and deriving the macroscopic ones, we have insight into anomalous diffusions, especially Lévy flights and tempered Lévy flights. In Section 2, all the derived equations are time dependent. From the process of derivation, one can see that the issue of initial condition can be easily/reasonably fixed, as classical ones, just specifying the value of $p(\mathbf{X}, 0)$ in the domain Ω . For Lévy processes, except Brownian motion, all others have discontinuous paths. As a result, the boundary $\partial\Omega$ itself (see Figure 3) can not be hit by the majority of discontinuous sample trajectories. This implies that when solving the PDEs derived in Section 2, the generalized boundary conditions must be introduced, i.e., the information of $p(\mathbf{X}, t)$ on the domain $\mathbb{R}^n \setminus \Omega$ must be properly accounted for. In the following, we focus on Eqs. (20), (28), (31), (35) to discuss the boundary issues.

3.1. Generalized Dirichlet type boundary conditions. The appropriate initial and boundary value problems for Eq. (20) should be

$$(40) \quad \begin{cases} \frac{\partial p(\mathbf{X}, t)}{\partial t} = \Delta^{\beta/2} p(\mathbf{X}, t) = \frac{-\beta \Gamma(\frac{n+\beta}{2})}{2^{1-\beta} \pi^{n/2} \Gamma(1-\beta/2)} \lim_{\epsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\epsilon}(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} & \text{in } \Omega, \\ p(\mathbf{X}, 0)|_{\Omega} = p_0(\mathbf{X}), \\ p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t). \end{cases}$$




Fig. 3: Domain of solving equations given in Section 2.

In Eq. (40), the term

$$\begin{aligned}
 & \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\varepsilon}(\mathbf{X})} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\mathcal{C}_{B_\varepsilon}(\mathbf{X}) \cap \Omega)} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} + \int_{\mathbb{R}^n \setminus \Omega} \frac{p(\mathbf{X}, t) - g(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} \\
 &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\mathcal{C}_{B_\varepsilon}(\mathbf{X}) \cap \Omega)} \frac{p(\mathbf{X}, t) - p(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y} + p(\mathbf{X}, t) \int_{\mathbb{R}^n \setminus \Omega} |\mathbf{X} - \mathbf{Y}|^{-n-\beta} d\mathbf{Y} \\
 &\quad + \int_{\mathbb{R}^n \setminus \Omega} \frac{-g(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y}.
 \end{aligned}
 \tag{41}$$

According to Eq. (41), $g(\mathbf{X}, t)$ should satisfy that there exist positive M and C such that when $|\mathbf{X}| > M$,

$$\frac{|g(\mathbf{X}, t)|}{|\mathbf{X}|^{\beta-\varepsilon}} < C \quad \text{for positive small } \varepsilon.
 \tag{42}$$

In particular, when Eq. (42) holds, the function $\int_{\mathbb{R}^n \setminus \Omega} \frac{-g(\mathbf{Y}, t)}{|\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y}$ of $\mathbf{X}$ has any order of derivative if $g(\mathbf{X}, t)$ is integrable in any bounded domain. One of the most popular cases is $g(\mathbf{X}, t) \equiv 0$, which is the so-called absorbing boundary condition, implying that the particle is killed whenever it leaves the domain Ω . For example, consider the steady state fractional diffusion equation

$$\begin{cases} \Delta^{\beta/2} p(\mathbf{X}) = -1 & \text{in } \Omega, \\ p(\mathbf{X})|_{\mathbb{R}^n \setminus \Omega} = 0. \end{cases}
 \tag{43}$$

The meaning of the solution $p(\mathbf{X})$ of (43) is the mean first exit time [6, 19] of particles performing Lévy flights [8]; if taking $\Omega = \{\mathbf{X} : |\mathbf{X}| < r\}$, then [16]

$$p(\mathbf{X}) = \frac{\Gamma(n/2)(r^2 - |\mathbf{x}|^2)^{\beta/2}}{2^\beta \Gamma(1 + \beta/2) \Gamma(n/2 + \beta/2)}.
 \tag{44}$$

We generate the sample trajectories of Lévy flights to plot one and two dimensional mean first exit times, respectively, in Figures 4 and 5, which confirm the effectiveness of the model (43).




Fig. 4: Behaviors of $p(\mathbf{x})$ generated by at least 1.5×10^6 trajectories with $\beta = 0.5$, $n = 1$, $r = 100$. The (red) solid line is theoretical result (eq. (44)). The symbols are for the simulation result.




Fig. 5: Behaviors of $p(\mathbf{x})$ generated by at least 1.5×10^6 trajectories with $\beta = 0.5$, $n = 2$, $r = 100$. The (red) solid line is theoretical result (eq. (44)). The symbols are for the simulation result.

Another interesting case is for the steady state fractional diffusion equation

$$(45) \quad \begin{cases} \Delta^{\beta/2} p(\mathbf{X}) = 0 & \text{in } \Omega, \\ p(\mathbf{X})|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}). \end{cases}$$

Given a domain $H \subset \mathbb{R}^n \setminus \Omega$, if taking $g(\mathbf{X}) = 1$ for $\mathbf{X} \in H$ and 0 for $\mathbf{X} \in (\mathbb{R}^n \setminus \Omega) \setminus H$, then the solution of (45) means the probability that the particles undergoing Lévy flights lands in H after first escaping the domain Ω [8]. If $g(\mathbf{X}) \equiv 1$ in $\mathbb{R}^n \setminus \Omega$, then $p(\mathbf{X})$ equals to 1 in Ω because of the probability interpretation. This can also be analytically checked.

375 For the initial and boundary value problem Eq. (28), it should be written as

$$376 \quad (46) \quad \begin{cases} \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2}} \\ \quad + \dots + \frac{\partial^{\beta_n} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n}} \quad \text{in } \Omega, \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, 0)|_{\Omega} = p_0(\mathbf{x}_1, \dots, \mathbf{x}_n), \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{x}_1, \dots, \mathbf{x}_n, t). \end{cases}$$

377 Similar to (41), in (46) the term

$$378 \quad (47) \quad \begin{aligned} & \lim_{\varepsilon \rightarrow 0^+} \int_{\mathcal{C}_{B_\varepsilon}(\mathbf{x}_j)} \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t) - p(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{(\mathcal{C}_{B_\varepsilon}(\mathbf{x}_j) \cap \Omega)} \frac{p(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t) - p(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j \\ & \quad + p(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t) \int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} |\mathbf{x}_j - \mathbf{y}_j|^{-1-\beta_j} d\mathbf{y}_j \\ & \quad + \int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} \frac{-g(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j. \end{aligned}$$

379 From Eq. (47), for $j = 1, \dots, n$, $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$ should satisfies that
380 there exist positive M and C such that when $|\mathbf{x}_j| > M$,

$$381 \quad (48) \quad \frac{|g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)|}{|\mathbf{x}_j|^{\beta_j - \varepsilon}} < C \quad \text{for positive small } \varepsilon.$$

382 The discussions below Eq. (45) still makes sense for Eq. (46). If $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$
383 satisfies Eq. (48), and it is integrable w.r.t. $\mathbf{x}_j$ in any bounded interval. Then
384 $\int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} \frac{-g(\mathbf{x}_1, \dots, \mathbf{y}_j, \dots, \mathbf{x}_n, t)}{|\mathbf{x}_j - \mathbf{y}_j|^{1+\beta_j}} d\mathbf{y}_j$ has any order of partial derivative w.r.t. $\mathbf{x}_j$.

385 The initial and boundary value problem for Eq. (31) is

$$386 \quad (49) \quad \begin{cases} \frac{\partial p(\mathbf{X}, t)}{\partial t} = (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t) \quad \text{in } \Omega, \\ p(\mathbf{X}, 0)|_{\Omega} = p_0(\mathbf{X}), \\ p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t). \end{cases}$$

387 Like the discussions for Eq. (40), $g(\mathbf{X}, t)$ should satisfies that there exist positive M
388 and C such that when $|\mathbf{X}| > M$,

$$389 \quad (50) \quad \frac{|g(\mathbf{X}, t)|}{e^{(\lambda - \varepsilon)|\mathbf{X}|}} < C \quad \text{for positive small } \varepsilon.$$

390 If Eq. (50) holds and $g(\mathbf{X}, t)$ is integrable in any bounded domain, the function
391 $\int_{\mathbb{R}^n \setminus \Omega} \frac{-g(\mathbf{Y}, t)}{e^{\lambda|\mathbf{X} - \mathbf{Y}|} |\mathbf{X} - \mathbf{Y}|^{n+\beta}} d\mathbf{Y}$ of $\mathbf{X}$ has any order of derivative.

392 Again, the corresponding tempered steady state fraction diffusion equation is

$$393 \quad (51) \quad \begin{cases} (\Delta + \lambda)^{\beta/2} p(\mathbf{X}) = 0 \quad \text{in } \Omega, \\ p(\mathbf{X})|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}). \end{cases}$$

For $H \subset \mathbb{R}^n \setminus \Omega$, if taking $g(\mathbf{X}) = 1$ for $\mathbf{X} \in H$ and 0 for $\mathbf{X} \in (\mathbb{R}^n \setminus \Omega) \setminus H$, then the solution of (51) means the probability that the particles undergoing tempered Lévy flights lands in H after first escaping the domain Ω . If $g(\mathbf{X}) \equiv 1$ in $\mathbb{R}^n \setminus \Omega$, then $p(\mathbf{X})$ equals to 1 in Ω .

The initial and boundary value problem (35) should be written as

$$(52) \quad \begin{cases} \frac{\partial p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial t} = \frac{\partial^{\beta_1, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1, \lambda}} + \frac{\partial^{\beta_2, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2, \lambda}} \\ \quad + \dots + \frac{\partial^{\beta_n, \lambda} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n, \lambda}} \quad \text{in } \Omega, \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, 0)|_{\Omega} = p_0(\mathbf{x}_1, \dots, \mathbf{x}_n), \\ p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{x}_1, \dots, \mathbf{x}_n, t). \end{cases}$$

For $j = 1, \dots, n$, $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$ should satisfy that there exist positive M and C such that when $|\mathbf{x}_j| > M$,

$$(53) \quad \frac{|g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)|}{e^{(\lambda - \varepsilon)|\mathbf{x}_j|}} < C \quad \text{for positive small } \varepsilon.$$

If $g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)$ is integrable w.r.t. $\mathbf{x}_j$ in any bounded interval and satisfies Eq. (53), then $\int_{\mathbb{R} \setminus (\Omega \cap \mathbb{R}_j)} \frac{-g(\mathbf{x}_1, \dots, \mathbf{x}_j, \dots, \mathbf{x}_n, t)}{e^{\lambda|\mathbf{x}_j - \mathbf{y}_j|} |\mathbf{x}_j - \mathbf{y}_j|^{1 + \beta_j}} d\mathbf{y}_j$ has any order of partial derivative w.r.t. $\mathbf{x}_j$.

The ways of specifying the initial and boundary conditions for Eqs. (36) and (38) are the same as Eq. (40). But for Eq. (36), the corresponding (42) should be changed as

$$(54) \quad \frac{|U(\mathbf{X})g(\mathbf{X}, t)|}{|\mathbf{X}|^{\beta - \varepsilon}} < C \quad \text{for positive small } \varepsilon.$$

Similarly, the initial and boundary conditions of Eqs. (37) and (39) should be specified as the ones of Eq. (49). But for Eq. (37), the corresponding (50) needs to be changed as

$$(55) \quad \frac{|U(\mathbf{X})g(\mathbf{X}, t)|}{e^{(\lambda - \varepsilon)|\mathbf{X}|}} < C \quad \text{for positive small } \varepsilon.$$

For the existence and uniqueness of the corresponding time-independent equations, one may refer to [14].

3.2. Generalized Neumann type boundary conditions. Because of the inherent discontinuity of the trajectories of Lévy flights or tempered Lévy flights, the traditional Neumann type boundary conditions can not be simply extended to the fractional PDEs. For the related discussions, see, e.g., [4, 10]. Based on the models built in Sec. 2 and the law of mass conservation, we derive the reasonable ways of specifying the Neumann type boundary conditions, especially the reflecting ones. Let us first recall the derivation of classical diffusion equation. For normal diffusion (Brownian motion), microscopically the first moment of the distribution of waiting times and the second moment of the distribution of jump length are bounded, i.e., in Laplace and Fourier spaces, they are respectively like $1 - c_1 u$ and $1 - c_2 |\mathbf{k}|^2$; plugging them into Eq. (1) or Eq. (9) and performing integral transformations lead to the classical diffusion equation

$$(56) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = (c_2/c_1) \Delta p(\mathbf{X}, t).$$




Fig. 6: Sketch map of particles jumping into, or jumping out of, or passing through the domain: Ω .

On the other hand, because of mass conservation, the continuity equation states that a change in density in any part of a system is due to inflow and outflow of particles into and out of that part of system, i.e., no particles are created or destroyed:

$$(57) \quad \frac{\partial p(\mathbf{X}, t)}{\partial t} = -\nabla \cdot \mathbf{j},$$

where $\mathbf{j}$ is the flux of diffusing particles. Combining (56) with (57), one may take

$$(58) \quad \mathbf{j} = -(c_2/c_1)\nabla p(\mathbf{X}, t),$$

which is exactly Fick's law, a phenomenological postulation, saying that the flux goes from regions of high concentration to regions of low concentration with a magnitude proportional to the concentration gradient. In fact, for a long history, even up to now, most of the people are more familiar with the process: using the continuity equation (57) and Fick's law (58) derives the diffusion equation (56). The so-called reflecting boundary condition for (56) is to let the flux $\mathbf{j}$ be zero along the boundary of considered domain.

Here we want to stress that Eq. (57) holds for any kind of diffusions, including the normal and anomalous ones. For Eqs. (40,46,49,52) governing the PDF of Lévy flights or tempered Lévy flights, using the continuity equation (57), one can get the corresponding fluxes and the counterparts of Fick's law; may we call it fractional Fick's law. Combining (40) with (57), one may let

$$(59) \quad \mathbf{j}_\Delta = \left\{ -\frac{1}{2n} c_{n,\beta} \int_{-\infty}^{\mathbf{x}_i} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{|\mathbf{Y}|^{n+\beta}} d\mathbf{Y} d\mathbf{x}_i \right\}_{n \times 1}$$

being the flux for the diffusion operator $\Delta^{\beta/2}$ with $\beta \in (0, 2)$, or calling it fractional Fick's law corresponding to $\Delta^{\beta/2}$. From (46) and (57), one may choose

$$(60) \quad \mathbf{j}_{hv} = \left\{ -\frac{1}{2} c_{1,\beta_i} \int_{-\infty}^{\mathbf{x}_i} \int_{-\infty}^{+\infty} \frac{p(\mathbf{X} + \tilde{\mathbf{Y}}_i, t) + p(\mathbf{X} - \tilde{\mathbf{Y}}_i, t) - 2 \cdot p(\mathbf{X}, t)}{|\mathbf{y}_i|^{1+\beta_i}} d\mathbf{y}_i d\mathbf{x}_i \right\}_{n \times 1},$$

where $\tilde{\mathbf{Y}}_i = \{\mathbf{x}_1, \dots, \mathbf{y}_i, \dots, \mathbf{x}_n\}^T$, being the flux (fractional Fick's law) corresponding to the horizontal and vertical type fractional operators. Similarly, we can also get the

flux (fractional Fick's law) corresponding to the tempered fractional Laplacian and tempered horizontal and vertical type fractional operators, being respectively taken as

$$(61) \quad \mathbf{j}_{\Delta, \lambda} = \left\{ -\frac{1}{2n} c_{n, \beta, \lambda} \int_{-\infty}^{\mathbf{x}_i} \int_{\mathbb{R}^n} \frac{p(\mathbf{X} + \mathbf{Y}, t) + p(\mathbf{X} - \mathbf{Y}, t) - 2 \cdot p(\mathbf{X}, t)}{e^{\lambda |\mathbf{Y}|} |\mathbf{Y}|^{n+\beta}} d\mathbf{Y} d\mathbf{x}_i \right\}_{n \times 1}$$

and

$$(62) \quad \mathbf{j}_{hv, \lambda} = \left\{ -\frac{1}{2} c_{1, \beta_i, \lambda} \int_{-\infty}^{\mathbf{x}_i} \int_{-\infty}^{+\infty} \frac{p(\mathbf{X} + \tilde{\mathbf{Y}}_i, t) + p(\mathbf{X} - \tilde{\mathbf{Y}}_i, t) - 2 \cdot p(\mathbf{X}, t)}{e^{\lambda |\mathbf{y}_i|} |\mathbf{y}_i|^{1+\beta_i}} d\mathbf{y}_i d\mathbf{x}_i \right\}_{n \times 1}$$

with $\tilde{\mathbf{Y}}_i = \{\mathbf{x}_1, \dots, \mathbf{y}_i, \dots, \mathbf{x}_n\}^T$.

Naturally, the Neumann type boundary conditions of (40,46,49,52) should be closely related to the values of the fluxes in the domain: $\mathbb{R}^n \setminus \Omega$; if the fluxes are zero in it, then one gets the so-called reflecting boundary conditions of the equations. Microscopically the motion of particles undergoing Lévy flights or tempered Lévy flights are much different from the Brownian motion; very rare but extremely long jumps dominate the dynamics, making the trajectories of the particles discontinuous. As shown in Figure 6, the particles may jump into, or jump out of, or even pass through the domain: Ω . But the number of particles inside Ω is conservative, which can be easily verified by making the integration of (57) in the domain Ω , i.e.,

$$(63) \quad \frac{\partial}{\partial t} \int_{\Omega} p(\mathbf{X}, t) d\mathbf{X} = - \int_{\Omega} \nabla \cdot \mathbf{j} d\mathbf{X} = - \int_{\partial\Omega} \mathbf{j} \cdot \mathbf{n} ds = 0,$$

where $\mathbf{n}$ is the outward-pointing unit normal vector on the boundary. If $\mathbf{j}|_{\mathbb{R}^n \setminus \Omega} = 0$, then for (40) $\Delta^{\frac{\beta}{2}} p(\mathbf{X}, t) = \nabla \cdot \mathbf{j} = 0$ in $\mathbb{R}^n \setminus \Omega$. So, the Neumann type boundary conditions for (40), (46), (49), and (52) can be, heuristically, defined as

$$(64) \quad \Delta^{\frac{\beta}{2}} p(\mathbf{X}, t) = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

$$(65) \quad \frac{\partial^{\beta_1} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_1}} + \frac{\partial^{\beta_2} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_2}} + \dots + \frac{\partial^{\beta_n} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_n}} = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

$$(66) \quad (\Delta + \lambda)^{\beta/2} p(\mathbf{X}, t) = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

and

$$(67) \quad \frac{\partial^{\beta_{1, \lambda}} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_1|^{\beta_{1, \lambda}}} + \frac{\partial^{\beta_{2, \lambda}} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_2|^{\beta_{2, \lambda}}} + \dots + \frac{\partial^{\beta_{n, \lambda}} p(\mathbf{x}_1, \dots, \mathbf{x}_n, t)}{\partial |\mathbf{x}_n|^{\beta_{n, \lambda}}} = g(\mathbf{X}) \quad \text{in } \mathbb{R}^n \setminus \Omega,$$

respectively. The corresponding reflecting boundary conditions are with $g(\mathbf{X}) \equiv 0$.

Remark: 1. The Neumann type boundary conditions (64)-(67) derived in this section are independent of the choice of the flux $\mathbf{j}$, provided that it satisfies the condition (57). 2. Sending $\beta \rightarrow 2$, the usual Neumann boundary condition is recovered; for example, consider the model

$$(68) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ \Delta^{\frac{\beta}{2}} p = g & \text{in } \mathbb{R}^n \setminus \Omega. \end{cases}$$

The weak solution $p : [0, +\infty) \rightarrow H^{\beta/2}(\mathbb{R}^n)$ of (68) satisfies

$$\int_{\Omega} p_t q d\mathbf{X} + \int_{\mathbb{R}^n} \Delta^{\beta/4} p \Delta^{\beta/4} q d\mathbf{X} = \int_{\Omega} f q d\mathbf{X} - \int_{\mathbb{R}^n \setminus \Omega} g q d\mathbf{X} \quad \forall q \in H^{\beta/2}(\mathbb{R}^n).$$

Taking $\beta \rightarrow 2$ leads to the usual Neumann boundary condition since

$$\int_{\mathbb{R}^n \setminus \Omega} g q d\mathbf{X} = \int_{\mathbb{R}^n \setminus \Omega} (\Delta p) q d\mathbf{X} = - \int_{\partial\Omega} \frac{\partial p}{\partial n} q ds - \int_{\mathbb{R}^n \setminus \Omega} \nabla p \cdot \nabla q d\mathbf{X}$$

and

$$\int_{\mathbb{R}^n} \Delta^{1/2} p \Delta^{1/2} q d\mathbf{X} = \int_{\mathbb{R}^n} \nabla p \cdot \nabla q d\mathbf{X}.$$

4. Well-posedness and regularity of the fractional PDEs with generalized BCs. Here, we show the well-posedness of the models discussed in the above sections, taking the models with the operator $\Delta^{\frac{\beta}{2}}$ as examples; the other ones can be similarly proved. For any real number $s \in \mathbb{R}$, we denote by $H^s(\mathbb{R}^n)$ the conventional Sobolev space of functions (see [1, 23]), equipped with the norm

$$\|u\|_{H^s(\mathbb{R}^n)} := \left(\int_{\mathbb{R}^n} (1 + |\mathbf{k}|^{2s}) |\hat{u}(\mathbf{k})|^2 d\mathbf{k} \right)^{\frac{1}{2}},$$

where $\hat{u}$ denotes the Fourier transform of u . The notation $H^s(\Omega)$ denotes the space of functions on Ω that admit extensions to $H^s(\mathbb{R}^n)$, equipped with the quotient norm

$$\|u\|_{H^s(\Omega)} := \inf_{\tilde{u}} \|\tilde{u}\|_{H^s(\mathbb{R}^n)},$$

where the infimum extends over all possible $\tilde{u} \in H^s(\mathbb{R}^n)$ such that $\tilde{u} = u$ on Ω (in the sense of distributions). The dual space of $H^s(\Omega)$ will be denoted by $H^s(\Omega)'$. The following inequality will be used below:

$$(69) \quad C^{-1}(\|\Delta^{\frac{\beta}{4}} u\|_{L^2(\mathbb{R}^n)} + \|u\|_{L^2(\Omega)}) \leq \|u\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \leq C(\|\Delta^{\frac{\beta}{4}} u\|_{L^2(\mathbb{R}^n)} + \|u\|_{L^2(\Omega)}).$$

Let $H_0^s(\Omega)$ be the subspace of $H^s(\mathbb{R}^n)$ consisting of functions which are zero in $\mathbb{R}^n \setminus \Omega$. It is isomorphic to the completion of $C_0^\infty(\Omega)$ in $H^s(\Omega)$. The dual space of $H_0^s(\Omega)$ will be denoted by $H^{-s}(\Omega)$.

For any Banach space B , the space $L^2(0, T; B)$ consists of functions $u : (0, T) \rightarrow B$ such that

$$(70) \quad \|u\|_{L^2(0, T; B)} := \left(\int_0^T \|u(\cdot, t)\|_B^2 dt \right)^{\frac{1}{2}} < \infty,$$

and $H^1(0, T; B) = \{u \in L^2(0, T; B) : \partial_t u \in L^2(0, T; B)\}$; see [12].

4.1. Dirichlet problem. For any given $g \in \mathbb{R} \cup (L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \cap H^1(0, T; H^{-\frac{\beta}{2}}(\mathbb{R}^n))) \hookrightarrow C([0, T]; L^2(\mathbb{R}^n))$, consider the time-dependent Dirichlet problem

$$(71) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ p = g & \text{in } \mathbb{R}^n \setminus \Omega, \\ p(\cdot, 0) = p_0 & \text{in } \Omega. \end{cases}$$

The weak formulation of (71) is to find $p = g + \phi$ such that

$$(72) \quad \phi \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)) \cap H^1(0, T; H^{-\frac{\beta}{2}}(\Omega)) \hookrightarrow C([0, T]; L^2(\Omega))$$

and

$$(73) \quad \int_0^T \int_{\Omega} \partial_t \phi q d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} \phi \Delta^{\frac{\beta}{4}} q d\mathbf{X} dt = \int_0^T \int_{\Omega} (f + \Delta^{\frac{\beta}{2}} g - \partial_t g) q d\mathbf{X} dt$$

$$\forall q \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)).$$

It is easy to see that $a(\phi, q) := \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} \phi \Delta^{\frac{\beta}{4}} q d\mathbf{X}$ is a coercive bilinear form on $H_0^{\frac{\beta}{2}}(\Omega) \times H_0^{\frac{\beta}{2}}(\Omega)$ (cf. [34, section 30.2]) and $\ell(q) := \int_{\Omega} (f + \Delta^{\frac{\beta}{2}} g - \partial_t g) q d\mathbf{X}$ is a continuous linear functional on $L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega))$. Such a problem as (73) has a unique weak solution (cf. [34, Theorem 30.A]).

The weak solution actually depends only on the values of g in $\mathbb{R}^n \setminus \Omega$, independent of the values of g in Ω . To see this, suppose that $g, \tilde{g} \in \mathbb{R} \cup (L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \cap H^1(0, T; H^{-\frac{\beta}{2}}(\mathbb{R}^n))) \hookrightarrow C([0, T]; L^2(\mathbb{R}^n))$ are two functions such that $g = \tilde{g}$ in $\mathbb{R}^n \setminus \Omega$, and p and $\tilde{p}$ are the weak solutions of

$$(74) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ p = g & \text{in } \mathbb{R}^n \setminus \Omega, \\ p(\cdot, 0) = p_0 & \text{in } \Omega, \end{cases} \quad \text{and} \quad \begin{cases} \frac{\partial \tilde{p}}{\partial t} - \Delta^{\frac{\beta}{2}} \tilde{p} = f & \text{in } \Omega, \\ \tilde{p} = \tilde{g} & \text{in } \mathbb{R}^n \setminus \Omega, \\ \tilde{p}(\cdot, 0) = p_0 & \text{in } \Omega, \end{cases}$$

respectively. Then the function $p - \tilde{p} \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)) \cap H^1(0, T; H^{-\frac{\beta}{2}}(\Omega))$ satisfies

$$(75) \quad \int_0^T \int_{\Omega} \partial_t (p - \tilde{p}) q d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} (p - \tilde{p}) \Delta^{\frac{\beta}{4}} q d\mathbf{X} dt = 0 \quad \forall q \in L^2(0, T; H_0^{\frac{\beta}{2}}(\Omega)).$$

Substituting $q = p - \tilde{p}$ into the equation above immediately yields $p - \tilde{p} = 0$ a.e. in $\mathbb{R}^n \times (0, T)$.

4.2. Neumann problem. Consider the Neumann problem

$$(76) \quad \begin{cases} \frac{\partial p}{\partial t} - \Delta^{\frac{\beta}{2}} p = f & \text{in } \Omega, \\ \Delta^{\frac{\beta}{2}} p = g & \text{in } \mathbb{R}^n \setminus \Omega, \\ p(\cdot, 0) = p_0 & \text{in } \Omega. \end{cases}$$

DEFINITION 1 (Weak solutions). The weak formulation of (76) is to find $p \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)) \cap C([0, T]; L^2(\Omega))$ such that

$$(77) \quad \partial_t p \in L^2(0, T; H^{\frac{\beta}{2}}(\Omega)') \text{ and } p(\cdot, 0) = p_0,$$

satisfying the following equation:

$$(78) \quad \begin{aligned} & \int_0^T \int_{\Omega} \partial_t p(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p(\mathbf{X}, t) \Delta^{\frac{\beta}{4}} q(\mathbf{X}, t) d\mathbf{X} dt \\ & = \int_0^T \int_{\Omega} f(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt - \int_0^T \int_{\mathbb{R}^n \setminus \Omega} g(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt \\ & \forall q \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)). \end{aligned}$$

THEOREM 2 (Existence and uniqueness of weak solutions). *If $p_0 \in L^2(\Omega)$, $f \in L^2(0, T; H^{\frac{\beta}{2}}(\Omega)')$ and $g \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)')$, then there exists a unique weak solution of (76) in the sense of Definition 1.*

Proof Let $t_k = k\tau$, $k = 0, 1, \dots, N$, be a partition of the time interval $[0, T]$, with step size $\tau = T/N$, and define

$$(79) \quad f_k(\mathbf{X}) := \frac{1}{\tau} \int_{t_{k-1}}^{t_k} f(\mathbf{X}, t) dt, \quad k = 0, 1, \dots, N,$$

$$(80) \quad g_k(\mathbf{X}) := \frac{1}{\tau} \int_{t_{k-1}}^{t_k} g(\mathbf{X}, t) dt, \quad k = 0, 1, \dots, N.$$

Consider the time-discrete problem: for a given $p_{k-1} \in L^2(\mathbb{R}^n)$, find $p_k \in H^{\frac{\beta}{2}}(\mathbb{R}^n)$ such that the following equation holds:

$$(81) \quad \begin{aligned} & \frac{1}{\tau} \int_{\Omega} p_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} + \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p_k(\mathbf{X}) \Delta^{\frac{\beta}{4}} q(\mathbf{X}) d\mathbf{X} \\ &= \frac{1}{\tau} \int_{\Omega} p_{k-1}(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} + \int_{\Omega} f_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} - \int_{\mathbb{R}^n \setminus \Omega} g_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} \quad \forall q \in H^{\frac{\beta}{2}}(\mathbb{R}^n). \end{aligned}$$

In view of (69), the left-hand side of the equation above is a coercive bilinear form on $H^{\frac{\beta}{2}}(\mathbb{R}^n) \times H^{\frac{\beta}{2}}(\mathbb{R}^n)$, while the right-hand side is a continuous linear functional on $H^{\frac{\beta}{2}}(\mathbb{R}^n)$. Consequently, the Lax–Milgram Lemma implies that there exists a unique solution $p_k \in H^{\frac{\beta}{2}}(\mathbb{R}^n)$ for (81).

Substituting $q = p_k$ into (81) yields

$$(82) \quad \begin{aligned} & \frac{\|p_k\|_{L^2(\Omega)}^2 - \|p_{k-1}\|_{L^2(\Omega)}^2}{2\tau} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}^2 \\ & \leq \|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} \|p_k\|_{H^{\frac{\beta}{2}}(\Omega)} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)} \\ & \leq (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}) \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \\ & \leq (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}) (\|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}^2 + \|p_k\|_{L^2(\Omega)}^2). \end{aligned}$$

Then, summing up the inequality above for $k = 1, 2, \dots, N$, we have

$$(83) \quad \begin{aligned} & \max_{1 \leq k \leq N} \|p_k\|_{L^2(\Omega)}^2 + \tau \sum_{k=1}^N \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}^2 \\ & \leq \|p_0\|_{L^2(\Omega)}^2 + C\tau \sum_{k=1}^N (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}^2 + \|p_k\|_{L^2(\Omega)}^2). \end{aligned}$$

By applying Grönwall's inequality to the last estimate, there exists a positive constant τ_0 such that when $\tau < \tau_0$ we have

$$(84) \quad \begin{aligned} & \max_{1 \leq k \leq N} \|p_k\|_{L^2(\Omega)}^2 + \tau \sum_{k=1}^N \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)}^2 \\ & \leq C\|p_0\|_{L^2(\Omega)}^2 + C\tau \sum_{k=1}^N (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}^2). \end{aligned}$$

567 Since any $q \in H^{\frac{\beta}{2}}(\Omega)$ can be extended to $q \in H^{\frac{\beta}{2}}(\mathbb{R}^n)$ with $\|q\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \leq 2\|q\|_{H^{\frac{\beta}{2}}(\Omega)}$,
 568 choosing such a q in (81) yields

$$\begin{aligned}
 569 \quad & \left| \int_{\Omega} \frac{p_k(\mathbf{X}) - p_{k-1}(\mathbf{X})}{\tau} q(\mathbf{X}) d\mathbf{X} \right| \\
 570 \quad &= \left| \int_{\Omega} f_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} - \int_{\mathbb{R}^n \setminus \Omega} g_k(\mathbf{X}) q(\mathbf{X}) d\mathbf{X} - \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p_k(\mathbf{X}) \Delta^{\frac{\beta}{4}} q(\mathbf{X}) d\mathbf{X} \right| \\
 571 \quad &\leq C(\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}) \|q\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)} \\
 572 \quad &\leq C(\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}) \|q\|_{H^{\frac{\beta}{2}}(\Omega)}. \\
 573
 \end{aligned}$$

574 which implies (via duality)

$$\begin{aligned}
 575 \quad (85) \quad & \left\| \frac{p_k - p_{k-1}}{\tau} \right\|_{H^{\frac{\beta}{2}}(\Omega)'} \leq C(\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'} + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'} + \|\Delta^{\frac{\beta}{4}} p_k\|_{L^2(\mathbb{R}^n)}). \\
 576
 \end{aligned}$$

577 The last inequality and (84) can be combined and written as

$$\begin{aligned}
 578 \quad & \max_{1 \leq k \leq N} \|p_k\|_{L^2(\Omega)}^2 + \tau \sum_{k=1}^N \left(\left\| \frac{p_k - p_{k-1}}{\tau} \right\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|p_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n)}^2 \right) \\
 579 \quad (86) \quad & \leq C\|p_0\|_{L^2(\Omega)}^2 + C\tau \sum_{k=1}^N (\|f_k\|_{H^{\frac{\beta}{2}}(\Omega)'}^2 + \|g_k\|_{H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)'}^2). \\
 580
 \end{aligned}$$

581 If we define the piecewise constant functions

$$582 \quad (87) \quad f^{(\tau)}(\mathbf{X}, t) := f_k(\mathbf{X}) = \frac{1}{\tau} \int_{t_{k-1}}^{t_k} f(\mathbf{X}, t) dt \quad \text{for } t \in (t_{k-1}, t_k], \quad k = 0, 1, \dots, N,$$

$$583 \quad (88) \quad g^{(\tau)}(\mathbf{X}, t) := g_k(\mathbf{X}) = \frac{1}{\tau} \int_{t_{k-1}}^{t_k} g(\mathbf{X}, t) dt \quad \text{for } t \in (t_{k-1}, t_k], \quad k = 0, 1, \dots, N,$$

$$584 \quad (89) \quad p_+^{(\tau)}(\mathbf{X}, t) := p_k(\mathbf{X}) \quad \text{for } t \in (t_{k-1}, t_k], \quad k = 0, 1, \dots, N,$$

586 and the piecewise linear function

$$\begin{aligned}
 587 \quad (90) \quad & p^{(\tau)}(\mathbf{X}, t) := \frac{t_k - t}{\tau} p_{k-1}(\mathbf{X}) + \frac{t - t_{k-1}}{\tau} p_k(\mathbf{X}) \quad \text{for } t \in [t_{k-1}, t_k], \quad k = 0, 1, \dots, N, \\
 588
 \end{aligned}$$

589 then (81) and (86) imply

$$\begin{aligned}
 & \int_0^T \int_{\Omega} \partial_t p^{(\tau)}(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}} p_+^{(\tau)}(\mathbf{X}, t) \Delta^{\frac{\beta}{4}} q(\mathbf{X}, t) d\mathbf{X} dt \\
 590 \quad &= \int_0^T \int_{\Omega} f^{(\tau)}(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt - \int_0^T \int_{\mathbb{R}^n \setminus \Omega} g^{(\tau)}(\mathbf{X}, t) q(\mathbf{X}, t) d\mathbf{X} dt \\
 & \quad \forall q \in L^2(0, T; H^{\frac{\beta}{2}}(\mathbb{R}^n)),
 \end{aligned}$$

591 and

$$\begin{aligned}
& \|p^{(\tau)}\|_{C([0,T];L^2(\Omega))} + \|\partial_t p^{(\tau)}\|_{L^2(0,T;H^{\frac{\beta}{2}}(\Omega)')} \\
& + \|p^{(\tau)}\|_{L^\infty(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n))} + \|p_+^{(\tau)}\|_{L^\infty(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n))} \\
592 & \leq C \left(\|f^{(\tau)}\|_{L^2(0,T;H^{\frac{\beta}{2}}(\Omega)')} + \|g^{(\tau)}\|_{L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)')} \right) \\
& \leq C \left(\|f\|_{L^2(0,T;H^{\frac{\beta}{2}}(\Omega)')} + \|g\|_{L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n \setminus \Omega)')} \right),
\end{aligned}$$

593 respectively, where the constant C is independent of the step size τ . The last in-
594 equality implies that $p^{(\tau)}$ is bounded in $H^1(0,T;H^{\frac{\beta}{2}}(\Omega)') \cap L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n)) \hookrightarrow$
595 $C([0,T];L^2(\Omega))$. Consequently, there exists $p \in H^1(0,T;H^{\frac{\beta}{2}}(\Omega)') \cap L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n)) \hookrightarrow$
596 $C([0,T];L^2(\Omega))$ and a subsequence $\tau_j \rightarrow 0$ such that

$$\begin{aligned}
597 \quad (91) \quad & p^{(\tau_j)} \text{ converges to } p \text{ weakly in } L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n)), \\
598 \quad (92) \quad & p_+^{(\tau_j)} \text{ converges to } p \text{ weakly in } L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n)), \\
599 \quad (93) \quad & \partial_t p^{(\tau_j)} \text{ converges to } \partial_t p \text{ weakly in } L^2(0,T;H^{\frac{\beta}{2}}(\Omega)'), \\
600 \quad (94) \quad & p^{(\tau_j)} \text{ converges to } p \text{ weakly in } C([0,T];H^{\frac{\beta}{2}}(\Omega)') \quad (\text{see [20, Appendix C]}).
\end{aligned}$$

602 By taking $\tau = \tau_j \rightarrow 0$ in (91), we obtain (78). This proves the existence of a weak
603 solution p satisfying (77).

604 If there are two weak solutions p and $\tilde{p}$, then their difference $\eta = p - \tilde{p}$ satisfies
605 the equation

$$\begin{aligned}
& (95) \\
606 \quad & \int_0^T \int_\Omega \partial_t(p - \tilde{p})q \, d\mathbf{X}dt + \int_0^T \int_{\mathbb{R}^n} \Delta^{\frac{\beta}{4}}(p - \tilde{p})\Delta^{\frac{\beta}{4}}q \, d\mathbf{X}dt = 0 \quad \forall q \in L^2(0,T;H^{\frac{\beta}{2}}(\mathbb{R}^n)).
\end{aligned}$$

608 Substituting $q = p - \tilde{p}$ into the equation yields

$$\begin{aligned}
& (96) \\
609 \quad & \|p(\cdot, t) - \tilde{p}(\cdot, t)\|_{L^2(\Omega)}^2 + \|\Delta^{\frac{\beta}{4}}(p - \tilde{p})\|_{L^2(0,T;L^2(\mathbb{R}^n))}^2 = \|p(\cdot, 0) - \tilde{p}(\cdot, 0)\|_{L^2(\Omega)}^2 = 0,
\end{aligned}$$

611 which implies $p = \tilde{p}$ a.e. in $\mathbb{R}^n \times (0, T)$. The uniqueness is proved.

612 **Remark:** From the analysis of this section we see that, although the initial data
613 $p_0(\mathbf{X})$ physically exists in the whole space $\mathbb{R}^n$, one only needs to know its values in Ω
614 to solve the PDEs (under both Dirichlet and Neumann boundary conditions).

615 **5. Conclusion.** In the past decades, fractional PDEs become popular as the
616 effective models of characterizing Lévy flights or tempered Lévy flights. This paper
617 is trying to answer the question: What are the physically meaningful and mathe-
618 matically reasonable boundary constraints for the models? We physically introduce
619 the process of the derivation of the fractional PDEs based on the microscopic mod-
620 els describing Lévy flights or tempered Lévy flights, and demonstrate that from a
621 physical point of view when solving the fractional PDEs in a bounded domain Ω , the
622 informations of the models in $\mathbb{R}^n \setminus \Omega$ should be involved. Inspired by the deriva-
623 tion process, we specify the Dirichlet type boundary constraint of the fractional

PDEs as $p(\mathbf{X}, t)|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$ and Neumann type boundary constraints as, e.g., $(\Delta^{\beta/2} p(\mathbf{X}, t))|_{\mathbb{R}^n \setminus \Omega} = g(\mathbf{X}, t)$ for the fractional Laplacian operator, which is the usual Neumann boundary condition when $\beta \rightarrow 2$.

The tempered fractional Laplacian operator $(\Delta + \lambda)^{\beta/2}$ is physically introduced and mathematically defined. For the four specific fractional PDEs given in this paper, we prove their well-posedness with the specified Dirichlet or Neumann type boundary constraints. In fact, it can be easily checked that these fractional PDEs are not well-posed if their boundary constraints are (locally) given in the traditional way; the potential reason is that locally dealing with the boundary contradicts with the principles that the Lévy or tempered Lévy flights follow.

REFERENCES

- [1] R. ADAMS AND A. J. J. F. FOURNIER, *Sobolev Spaces*, Elsevier/Academic Press, Amsterdam, second ed., 2003.
- [2] D. APPLEBAUM, *Lévy processes and stochastic calculus*, Cambridge University Press, Cambridge, second ed., 2009, <https://doi.org/10.1017/CBO9780511809781>.
- [3] E. BARKAI, A. V. NAUMOV, Y. G. VAINER, M. BAUER, AND L. KADOR, *Lévy statistics for random single-molecule line shapes in a glass*, Phys. Rev. Lett., 91 (2003), p. 075502, <https://doi.org/10.1103/physrevlett.91.075502>.
- [4] G. BARLES, C. GEORGELIN, AND E. R. JAKOBSEN, *On Neumann and oblique derivatives boundary conditions for nonlocal elliptic equations*, J. Differential Equations, 256 (2014), pp. 1368–1394, <https://doi.org/10.1016/j.jde.2013.11.001>.
- [5] J.-P. BOUCHAUD AND A. GEORGES, *Anomalous diffusion in disordered media: Statistical mechanisms, models and physical applications*, Phys. Rep., 195 (1990), pp. 127–293, [https://doi.org/10.1016/0370-1573\(90\)90099-n](https://doi.org/10.1016/0370-1573(90)90099-n).
- [6] A.V. CHECHKIN, R. METZLER, V.Y. GONCHAR, J. KLAFTER, AND L.V. TANATAROV, *First passage and arrival time densities for Lévy flights and the failure of the method of images*, J. Phys. A: Math. Gen., 36 (2003), pp. L537–L544, <https://stacks.iop.org/JPhysA/36/L537>.
- [7] Z.-Q. CHEN AND R. SONG, *Two-sided eigenvalue estimates for subordinate processes in domains*, J. Funct. Anal., 226 (2005), pp. 90–113, <https://doi.org/10.1016/j.jfa.2005.05.004>.
- [8] W. H. DENG, X. C. WU, AND W. L. WANG, *Mean exit time and escape probability for the anomalous processes with the tempered power-law waiting times*, EPL, 117 (2017), p. 10009, <https://doi.org/10.1209/0295-5075/117/10009>.
- [9] E. DI NEZZA, G. PALATUCCI, AND E. VALDINOCI, *Hitchhiker’s guide to the fractional Sobolev spaces*, Bull. Sci. math., 136 (2012), pp. 521–573, <https://doi.org/10.1016/j.bulsci.2011.12.004>.
- [10] S. DIPIERRO, X. ROS-OTON, AND E. VALDINOCI, *Nonlocal problems with Neumann boundary conditions*, Rev. Mat. Iberoam., (2017).
- [11] Q. DU, M. GUNZBURGER, R. B. LEHOUCQ, AND K. ZHOU, *Analysis and approximation of nonlocal diffusion problems with volume constraints*, SIAM Rev., 54 (2012), pp. 667–696, <https://doi.org/10.1137/110833294>.
- [12] L. C. EVANS, *Partial differential equations*, American Mathematical Society, Providence, RI, second ed., 2010.
- [13] W. FELLER, *An introduction to probability theory and its applications*, John Wiley & Sons, Inc., New York, 1971. Vol. 2, Chap. XVI. 8, p. 525.
- [14] M. FELSINGER, M. KASSMANN, AND P. VOIGT, *The Dirichlet problem for nonlocal operators*, Math. Z., 279 (2015), pp. 779–809, <https://doi.org/10.1007/s00209-014-1394-3>.
- [15] H. C. FOGEDBY, *Langevin equations for continuous time Lévy flights*, Phys. Rev. E, 50 (1994), pp. 1657–1660, <https://doi.org/10.1103/physreve.50.1657>.
- [16] R. K. GETTOOR, *First passage times for symmetric stable processes in space*, Trans. Amer. Math. Soc., 101 (1961), pp. 75–90, <https://doi.org/10.1090/S0002-9947-1961-0137148-5>.
- [17] Q.-Y. GUAN AND Z.-M. MA, *Boundary problems for fractional Laplacians*, Stoch. Dyn., 5 (2005), pp. 385–424, <https://doi.org/10.1142/S021949370500150x>.
- [18] I. KOPONEN, *Analytic approach to the problem of convergence of truncated Lévy flights towards the Gaussian stochastic process*, Phys. Rev. E, 52 (1995), pp. 1197–1199, <https://doi.org/10.1103/physreve.52.1197>.
- [19] T. KOREN, M.A. LOMHOLT, A.V. CHECHKIN, J. KLAFTER, AND R. METZLER, *Leapover lengths*

- and first passage time statistics for Lévy flights, Phys. Rev. Lett., 99 (2007), p. 160602, <https://doi.org/10.1103/PhysRevLett.99.160602>.
- [20] P.-L. LIONS, *Mathematical Topics in Fluid Mechanics: Volume 1: Incompressible Models*, Clarendon Press, Oxford, USA., 1996.
- [21] B. B. MANDELBROT AND J. W. V. NESS, *Fractional Brownian motions, fractional noises and applications*, SIAM Rev., 10 (1968), pp. 422–437, <https://doi.org/10.1137/1010093>.
- [22] R. N. MANTEGNA AND H. E. STANLEY, *Stochastic process with ultraslow convergence to a Gaussian: The truncated Lévy flight*, Phys. Rev. Lett., 73 (1994), pp. 2946–2949, <https://doi.org/10.1103/physrevlett.73.2946>.
- [23] W. MCLEAN, *Strongly elliptic systems and boundary integral equations*, Cambridge university press, 2000.
- [24] M. M. MEERSCHAERT AND A. SIKORSKII, *Stochastic Models for Fractional Calculus*, Walter de Gruyter, Berlin, 2012.
- [25] R. METZLER AND J. KLAFTER, *The random walk's guide to anomalous diffusion: a fractional dynamics approach*, Phys. Rep., 339 (2000), pp. 1–77, [https://doi.org/10.1016/S0370-1573\(00\)00070-3](https://doi.org/10.1016/S0370-1573(00)00070-3).
- [26] E. W. MONTROLL AND G. H. WEISS, *Random walks on lattices. II*, J. Math. Phys., 6 (1965), pp. 167–181, <https://doi.org/10.1063/1.1704269>.
- [27] J. M. SANCHO, A. M. LACASTA, K. LINDENBERG, I. M. SOKOLOV, AND A. H. ROMERO, *Diffusion on a solid surface: Anomalous is normal*, Phys. Rev. Lett., 92 (2004), p. 250601, <https://doi.org/10.1103/physrevlett.92.250601>.
- [28] M. F. SHLESINGER, J. KLAFTER, AND B. J. WEST, *Levy walks with applications to turbulence and chaos*, Phys. A, 140 (1986), pp. 212–218, [https://doi.org/10.1016/0378-4371\(86\)90224-4](https://doi.org/10.1016/0378-4371(86)90224-4).
- [29] L. SILVESTRE, *Regularity of the obstacle problem for a fractional power of the Laplace operator*, Commun. Pure Appl. Math., 60 (2007), pp. 67–112, <https://doi.org/10.1002/cpa.20153>.
- [30] I. M. SOKOLOV, J. MAI, AND A. BLUMEN, *Paradoxal diffusion in chemical space for nearest-neighbor walks over polymer chains*, Phys. Rev. Lett., 79 (1997), pp. 857–860, <https://doi.org/10.1103/physrevlett.79.857>.
- [31] L. TURGEMAN, S. CARMÍ, AND E. BARKAI, *Fractional Feynman-Kac equation for non-Brownian functionals*, Phys. Rev. Lett., 103 (2009), p. 190201, <https://doi.org/10.1103/physrevlett.103.190201>.
- [32] X. C. WU, W. H. DENG, AND E. BARKAI, *Tempered fractional Feynman-Kac equation: Theory and examples*, Phys. Rev. E, 93 (2016), p. 032151, <https://doi.org/10.1103/physreve.93.032151>.
- [33] V. ZABURDAEV, S. DENISOV, AND J. KLAFTER, *Lévy walks*, Rev. Mod. Phys., 87 (2015), pp. 483–530, <https://doi.org/10.1103/revmodphys.87.483>.
- [34] E. ZEIDLER, *Nonlinear Functional Analysis and its Applications II/B: Nonlinear Monotone Operators*, Springer Science+Business Media, LLC, 1990, <https://doi.org/10.1007/978-1-4612-0981-2>. Translated by the author and by Leo F. Boron.